7.3: Institutional Distinctiveness:

Additional information and supporting document

"Nurturing excellence in education and research for sustainable innovation"

INDEX

Sr. No	File Description
Criteria 7	Institutional Value and Best Practices
7.3	7.3: Institutional Distinctiveness: Additional Information and Supporting Documents
7.3.1.1	Portray the performance of the institution in one area distinctive to its priority and thrust within 1000 words "Nourishing Pharmacy Professionals through Value Based Education"
7.3.1.2	Research Publications Books and Books Chapters published by the faculty

Since its founding, the goal of Loknete Shri Dadapatil Pharate College of Pharmacy (LSDPCOP), Mandavgan Pharata, has been to give our rural students access to cutting-edge educational resources so they may comfortably compete with the outside world. The main goal is to "Nourishing Pharmacy Professionals through Value Based Education" so that they can effectively serve the profession. In the end, we are the company that carries out our social obligations in addition to offering top-notch instruction to create competent, professional chemists. Moral education is vital, in our opinion, because it is at this age that children take their first steps towards adulthood, and it is crucial that they do so properly.

Their views, convictions, and concepts are shaped by these moral principles, which also aid in their development as morally robust and unwavering individuals. Loknete Shri Dadapatil Pharate College of Pharmacy (LSDPCOP), Mandavgan Pharata, works hard to instill in its pupil's social duties and responsibilities as well as moral and ethical ideals. This has been accomplished by creating and offering our pupils life skills classes.

Life skills are those proficiencies that enable a person to be adaptive and optimistic in the face of life's challenges. The goal is the development of one's personality through self-awareness, interpersonal relationships, introspection on both the concrete and abstract, change-leading and change-generating, and adherence to tried-and-true beliefs. By exposing students to the ideas that underpin both personal and professional success and assisting them in acquiring the skills necessary to put these ideas into practice in their daily lives and jobs, this course aims to maximise their potential and increase their employability.

In order to attract superior talent, LSDPCOP is placing greater emphasis on teaching life skills, including professional, leadership, and management abilities, as well as communication skills. Adopting life skills is, in our opinion, the secret to excellence.

The goal of the life skills course is to instil in the students a sense of self-awareness and empathy as well as professional abilities in the areas of interpersonal skills, high-performing teams, leadership potential, communication and presentation skills, problem-solving techniques, decision-making, inspiring creativity and innovation for both professional and personal excellence, stress management, time management, conflict resolution, and human values instillation.

As facilitators, mentors, and trainers, the faculty members and students must actively participate in the course. Students are motivated and enrolled in the courses they choose for this reason. Oral presentations, spontaneous speaking, quizzes, debates, case studies, creative thinking, team-building exercises, field trips, projects, psychometric analysis, and other activities have all been included into the courses. They receive a course completion certificate upon successfully finishing the course.

3.3.1. Number of papers published per teacher in the Journals notified on UGC website during the last five years.

Summary

Sr. No.	Sr. No. Parameter Academic	Year	No. of Research paper
1	**Research Paper Publications**	2018-19	42
2		2019-20	16
3		2020-21	21
4		2021-22	14
5		2022-23	20
Total Number of research papers published in the Journals notified on UGC CARE list during the last five years			**113**

3.3.1. Number of papers published per teacher in the Journals notified on UGC Website during the last five years.

Index

Sr. No.	Content	Document
1	Academic Year 2018-2019	View Document
2	Academic Year 2019-2020	View Document
3	Academic Year 2020-2021	View Document
4	Academic Year 2021-2022	View Document
5	Academic Year 2022-2023	View Document

Academic Year 2018-2019

Sr. no	Title of the paper	Name of Staff	Department	Name of the journal	ISSN number	Document	Link
1	Evaluation Of Some Natural Polymers As Pharmaceutical Excipient For Sustained Drug Delivery System	Mr. A. V. Dhobale	Pharmaceutics	International Journal Of Humanities And Social Science	E-ISSN 2348 – 1269	View Document	https://www.researchgate.net/publication/350782955_Evaluation_of_Some_Natural_Polymers_as_Pharmaceutical_Excipient_For_Sustained_Drug_Delivery_System
2	Recent Advances In Mucoadhesive Buccal Drug Delivery System And Its Marketed Scope And Opportunities	Mr. A. V. Dhobale	Pharmaceutics	International Journal Of Advanced Pharmaceutical Sciences	ISSN: 2456-8147	View Document	https://www.researchgate.net/publication/327111123_RECENT_ADVANCES_IN_MUCOADHESIVE_BUCCAL_DRUG_DELIVERY_SYSTEM_AND_ITS_MARKETED_SCOPE_AND_OPPORTUNITIES
3	Evaluation Of Antiulcer Activity Of Aconitum Heterophyllum On Experimental Animal	Mr. A. V. Dhobale	Pharmaceutics	World J Pharm Pharm Sci	ISSN 2278 – 4357	View Document	https://www.researchgate.net/publication/326837658_EVALUATION_OF_ANTIULCER_ACTIVITY_OF_ACONITUM_HETEROPHYLLU

Back to Academic Year 2018-2019

							M ON EXPERIMENTAL ANIMAL
4	Design And In-Vitro Characterization Of Gastroretentive Floating Tablets Containing Gliclazide	Mr. A. V. Dhobale	Pharmaceutics	World J Pharm Pharm Sci	ISSN 2278 – 4357	View Document	https://www.researchgate.net/publication/322307709_DESIGN_AND_IN-VITRO_CHARACTERIZATION_OF_GASTRORETENTIVE_FLOATING_TABLETS_CONTAINING_GLICLAZIDE
5	Mouth Dissolving Strips	Mr. A. V. Dhobale	Pharmaceutics	International Journal Of Advanced Pharmaceutical Sciences	ISSN: 2456-8147	View Document	https://www.researchgate.net/profile/Avinash-Dhobale-2
6	Formulation And Evaluation Of Fast Disintegrating Tablet Of Carvedilol.	Mr. A. V. Dhobale	Pharmaceutics	International Journal Of Pharmacy Pharmaceutical Sciences	ISSN 2349 7203	View Document	https://www.researchgate.net/publication/327690399_Formulation_and_Evaluation_of_Fast_Disintegrating_Tablet_of_Carvedilol
7	Evaluation Of Antiulcer Activity Of Aconitum Heterophyllum On Experimental Animal	Mr. A. V. Dhobale	Pharmaceutics	World Journal Of Pharmacy And Pharmaceutical Sciences	ISSN 2278 4357	View Document	https://www.researchgate.net/publication/326837658_EVALUATION_OF_ANTIULCER_ACTIVITY_OF_ACONITUM_HETEROPHYLLUM_ON_EXPERIMENTAL_ANIMAL

8	Formulation And Evaluation Of Atorvastatin Calcium Oral Dispersible Table	Mr. A. V. Dhobale	Pharmaceutics	International Journal Of Current Advanced Research	ISSN: O: 2319-6475, ISSN: P: 2319-6505,	View Document	https://www.researchgate.net/publication/326831851_FORMULATION_AND_EVALUATION_OF_ATORVASTATIN_CALCIUM_ORAL_DISPERSIBLE_TABLET
9	Formulation And Evaluation Of Carbamazepine Nanoemulsion For Brain Targeted Drug Delivery Via Intranasal Route	Mr. A. V. Dhobale	Pharmaceutics	Indo American Journal Of Pharmaceutical Research	ISSN NO: 2231-6876	View Document	https://www.researchgate.net/publication/326803830_FORMULATION_AND_EVALUATION_OF_CARBAMAZEPINE_NANOEMULSION_FOR_BRAIN_TARGETED_DRUG_DELIVERY_VIA_INTRANASAL_ROUTE
10	Recent Advances In Pilot Plant Scale Up Techniques	Mr. A. V. Dhobale	Pharmaceutics	Indo American Journal Of Pharmaceutical Research,	ISSN NO: 2231-6876	View Document	https://www.researchgate.net/publication/326304751_RECENT_ADVANCES_IN_PILOT_PLANT_SCALE_UP_TECHNIQUES-A_Review
11	Design, Development And Characterization Oflosartan Potassium Sustained	Mr. A. V. Dhobale	Pharmaceutics	World Journal Of Pharmacy And Pharmaceutical Scinces	ISSN 2278 – 4357	View Document	https://www.researchgate.net/publication/322308014_DESIGN_DEVELOPMENT_AND_CHARACTERIZATION_OF_LOSARTAN_POTA

	Release Matrix Tablet						SSIUM_SUSTAINED_RELEASE_MATRIX_TABLET
12	Design And In-Vitro Characterization Of Gastroretentive Floating Tablets Containing Gliclazide	Mr. A. V. Dhobale	Pharmaceutics	World Journal Of Pharmacy And Pharmaceutical Sciences	ISSN 2278 – 4357	View Document	https://www.researchgate.net/publication/322307709_DESIGN_AND_IN-VITRO_CHARACTERIZATION_OF_GASTRORETENTIVE_FLOATING_TABLETS_CONTAINING_GLICLAZIDE
14	Formulation And Evaluation Of Rosuvastatin Oral Dispersible Tablet Under A Creative Commons	Dr. V. M. Satpute	Pharmaceutics	International Journal Of Advanced Pharmaceutical Sciences	ISSN: 2456-8147	View Document	https://www.researchgate.net/publication/340262020_FORMULATION_AND_EVALUATION_OF_ROSUVASTATIN_ORAL_DISPERSIBLE_TABLET_under_a_Creative_Commons_Attribution_40
15	A Project On Human Papillomavirus-A Review Under A Creative Commons Attribution	Dr. V. M. Satpute	Pharmaceutics Pharmaceutics	International Journal Of Advanced Pharmaceutical Sciences	SSN:2456-8147	View Document	https://www.researchgate.net/publication/326571009_A_PROJECT_ON_HUMAN_PAPILLOMAVIRUS_-_A_REVIEW
16	Formulation And Evaluation Of Fast Disintegrating	Dr. V. M. Satpute	Pharmaceutics	Human Journals IJPPR	ISSN 2349 7203	View Document	https://www.researchgate.net/publication/327690399_Formulat

	Tabletof Carvedilol						ion and Evaluation of Fast Disintegrating Tablet of Carvedilol
17	Isolation And Characterization Of Phytoconstituents From Petroleum Ether Extract Of Momordica Cochinchinensis Fruits	Mr. M. R. Agrawal	Pharmacognosy	Journal Of Pharmacognosy And Phytochemistry	E-ISSN: 2278-4136 P-ISSN: 2349-8234	View Document	https://rjpponline.org/AbstractView.aspx?PID=2020-12-3-6
18	Anti-Inflammatory Activity Of Momordica Cochinchinensis And Momordica Balsamina Fruit Extracts	Mr. M. R. Agrawal	Pharmacognosy	International Journal Of Green	DOI: 10.24214/IJGHC/HC/7/4/44655.	View Document	https://www.academia.edu/42305275/Analgesic_activity_of_Momordica_cochinchinensis_and_Momordica_balsamina_fruit_extracts
19	Analgesic Activity Of Momordica Cochinchinensis And Momordica Balsamina Fruit Extracts	Mr. M. R. Agrawal	Pharmacognosy	International Journal Of Gree Pharmacy	Vol 12(4)	View Document	https://www.academia.edu/42305275/Analgesic_activity_of_Momordica_cochinchinensis_and_Momordica_balsamina_fruit_extracts
20	Ameliorative Effects of Artemisia Pallens In A Murine Model of Ovalbumin-Asthma Via Modulation of	Ms. A. A. Mukherjee	Pharmacognosy	Biomedicine & Pharmacotherapy	Https://Doi.Org/10.1016/J.Fct.2018.08.028	View Document	https://pubmed.ncbi.nlm.nih.gov/28810518/#:~:text=Conclusion%3A%20Artemisia%20pallens%20simultaneously%20orchestrate,OVA%2Dinduc

Back to Academic Year 2018-2019

	Biochemical perturbations						ed%20AHR%20in%20rats.
21	Evaluation f Health-Related Quality Of Life In Hemolytic Uraemic Syndrome Patients Treated With Eculizumab: A Systematic Evaluation On Basis Of EMPRO	Ms. A. A. Mukherjee	Pharmacognosy	Taylor & Francis	ISSN: 0886-022X (Print) 1525-6049	View Document	https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6014301/
22	Curcumin Ameliorates Vitamin A Deficiency-Induced Urolithiasis In Neonatal Rats Via Inhibition Of KIM-1/NGAL, Nrf2, And Inos Pathways	Ms. A. Kandhare	Pharmacology	Latin American Journal Of Pharmacy	ISSN 0326 2383 ISSN 2362-3853	View Document	https://www.researchgate.net/publication/331045453_Curcumin_ameliorates_vitamin_A_deficiency-induced_urolithiasis_in_neonatal_rats_via_inhibition_of_KIM-1NGAL_Nrf2_and_iNOs_pathways
23	Neuroprotective Effect Of Naringin, A Flavone Glycoside In Quinolinic Acidinduced Neurotoxicity: Possible Role Of PPAR-Γ, Bax/Bcl-2, And Caspase-3	Ms. A. Kandhare	Pharmacology	Food And Chemical Toxicology 121 (2018) 95–108	Https://Doi.Org/10.1016/J.Fct.2018.08.028	View Document	https://pubmed.ncbi.nlm.nih.gov/30130594/

24	Low Molecular Weight Galactomannans-Based Standardized Fenugreek Seed Extract Ameliorates High-Fat Diet-Induced Obesity In Mice Via Modulation Of Fasn, IL-6, Leptin, And TRIP-Br_2	Ms. A. Kandhare	Pharmacology	RSC Advances	Rsc.Li/Rsc-Advances	View Document	https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9086199/
25	Anti-Epileptic Effect Of Morin Against Experimental Pentylenetetrazol-Induced Seizures Via Modulating Brain Monoamines And Oxidative Stress	Ms. A. Kandhare	Pharmacology	Asian Pacific Journal Of Tropical Biomedicine 2018; 8(7): 352-359	ISSN 2221-1619	View Document	https://www.researchgate.net/publication/326472470_Anti-epileptic_effect_of_morin_against_experimental_pentylenetetrazol-induced_seizures_via_modulating_brain_monoamines_and_oxidative_stress
26	Add-On Therapy Of Herbal Formulation Rich In Standardized Fenugreek Seed Extract In Type 2 Diabetes Mellitus Patients With Insulin Therapy: An Efficacy And Safety Study	Ms. A. Kandhare	Pharmacology	Asian Pacific Journal Of Tropical Biomedicine 2018; 8(7): 352-359	ISSN 2221-1619	View Document	https://www.researchgate.net/publication/327910450_Add-on_therapy_of_herbal_formulation_rich_in_standardized_fenugreek_seed_extract_in_type_2_diabetes_mellitus_patients_with_insulin_therapy_An_efficacy_and_safety_study

27	Efficacy And Safety Of Herbal Formulation Rich In Standardized Fenugreek Seed Extract As Add-On Supplementation In Patients With Type 2 Diabetes Mellitus On Sulfonylurea Therapy: A 12-Week, Randomized, Double-Blind, Placebo-Controlled, Multi-Center Study			Pharmacognosy Magazine	ISSN : 0973-1296	View Document	https://www.researchgate.net/publication/327521981_Efficacy_and_safety_of_herbal_formulation_rich_in_standardized_fenugreek_seed_extract_as_add-on_supplementation_in_patients_with_type_2_diabetes_mellitus_on_sulfonylurea_therapy_A_12-week_randomized_d
28	Evaluation Of Health-Related Quality Of Life In Hemolytic Uraemic Syndrome Patients Treated With Eculizumab: A Systematic Evaluation On Basis Of EMPRO			Taylor & Francis	ISSN: 1525-6049	View Document	https://pubmed.ncbi.nlm.nih.gov/29363392/
29	Hesperidin, A Plant Flavonoid Accelerated The Cutaneous Wound Healing In Streptozotocin-Induced Diabetic Rats: Role Of			EXCLI Journal	ISSN 1611-2156	View Document	https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5962903/

	Tgf-B/Smads And Ang-1/Tie2						
30	Anti-Allergic Potential Of Fisetin In A Murine Model Of Ova-Induced Allergic Rhinitis Via Inhibition Of Gata-3 And Th2 Cytokines Zhao L			Biomedica	Vol. 34, Issue 2, Apr. – Jun., 2018	View Document	https://www.google.com/search?q=Anti-Allergic+Potential+Of+Fisetin+In+A+Murine+Model+Of+Ova-Induced+Allergic+Rhinitis+Via+Inhibition+Of+Gata-3+And+Th2+Cyt

Back to Academic Year 2018-2019

research+paper+by+amit+kandhare&rlz=1C1VDKB_en-GBIN1070IN1070&oq=Anti-Allergic+Potential+Of+Fisetin+In+A+Murine+Model+Of+Ova-Induced+Allergic+Rhinitis+Via+Inhibition+Of+Gata-3+And+Th2+Cytokines+Zhao+L+research+paper+by+amit+kandhare&gs_lcrp=EgZjaHJvbWUqBggAEEUYOzIGCAAQRRg7MgcIARAhGI8C0gEJMTAwOTNqMGo3qAIAsAIA&sourceid=chrome&ie=UTF-8#ip=1

31	Elucidation Of The Molecular Mechanism Of Tempol In Pentylenetetrazol-Induced Epilepsy In Mice: Role Of Gamma - Amino butyric Acid, Tumor Necrosis Factor-Alpha, Interleukin-1β And C-Fos			Pharmacognosy Magazine	ISSN : 0973-1296	View Document	https://ouci.dntb.gov.ua/en/works/4k3GEqn4/
32	Fisetin, A Plant Flavonoid Ameliorates Doxorubicin-Induced Cardiotoxicity In Experimental Rats: The Decisive Role OfCaspase-3, COX-II, Ctn-I, Inos And TNF			Molecular Biology Reports	ISSN 0301-4851	View Document	https://ouci.dntb.gov.ua/en/works/7P2pwv0l/
33	Pioglitazone And Risk Of Bladder Cancer In Type 2 Diabetes Mellitus Patients: A Systematic Literature Review And Meta-Analysis Of Observational Studies Using Real-World Data			Clinical Epidemiology And Global Health 6 (2018) 61–68	Http://Dx.Doi.Org/10.1016/J.Cegh.2017.08.002	View Document	https://www.sciencedirect.com/science/article/pii/S2213398417300544
34	Development And Validation Of UV-Visible Spectroscopy Method For Simultaneous	Mr. Pravin Cholke &	Pharmaceutical Chemistry	International Journal Of Research In Pharmacy	ISSN: 2455-698	View Document	https://www.pharmacyjournal.in/archives/2018/vol3/issue4/3-4-16

	Estimation Of Saxagliptin Hydrochloride And Metformin Hydrochloride In Tablet Dosage Form	Ms. Yogita Temak		And Pharmaceutical Sciences			
35	Formulation And Evaluation Of Mouth Dissolving Tablet Of Bendroflumethiazide	Ms. S. A. Nagare	Pharmaceutical Chemistry	International Journal Of Current Research	ISSN: 0975-833	View Document	https://www.researchgate.net/publication/326849498_FORMULATION_AND_EVALUATION_OF_MOUTH_DISSOLVING_TABLET_OF_BENDROFLUMETHIAZIDE
36	"Formulation And Evaluation Of Carbamazepine Nanoemulsion For Brain Targeted Drug Delivery Via Intranasal Route"	Ms. S. A. Nagare	Pharmaceutical Chemistry	Indo American Journal Of Pharmaceutical Research	ISSN NO: 2231-6876	View Document	https://www.researchgate.net/publication/326803830_FORMULATION_AND_EVALUATION_OF_CARBAMAZEPINE_NANOEMULSION_FOR_BRAIN_TARGETED_DRUG_DELIVERY_VIA_INTRANASAL_ROUTE
37	Formulation And Evaluation Of Mouth Dissolving Tablet Of Bendroflumethiazide	Ms. A. A. Najan	Pharmaceutical Chemistry	International Journal Of Current Research	ISSN: 0975-833	View Document	https://scholar.google.co.in/scholar?q=Formulation+And+Evaluation+Of+Mouth+Dissolving+Tablet+Of+Bendroflumethiazide+research+paper+by+a.a.najan&hl=en&a

							s_sdt=0&as_vis=1&oi=scholart
38	Curcumin Ameliorates Vitamin A Deficiency-Induced Urolithiasis In Neonatal Rats Via Inhibition Of KIM-1/NGAL, Nrf2, And Inos Pathways	Ms. A. A. Mukherjee	Pharmacology	Latin American Journal Of Pharmacy	ISSN 0326 2383 ISSN 2362-3853	View Document	https://www.researchgate.net/publication/331045453_Curcumin_ameliorates_vitamin_A_deficiency-induced_urolithiasis_in_neonatal_rats_via_inhibition_of_KIM-1NGAL_Nrf2_and_iNOs_pathways
39	Evaluation Of Health-Related Quality Of Life In Hemolytic Uraemic Syndrome Patients Treated With Eculizumab: A Systematic Evaluation On Basis Of EMPRO	Ms. A. A. Mukherjee		Taylor & Francis	ISSN: 1525-6049	View Document	https://pubmed.ncbi.nlm.nih.gov/29363392/
40	Hesperidin, A Plant Flavonoid Accelerated The Cutaneous Wound Healing In Streptozotocin-Induced Diabetic Rats: Role Of Tgf-B/Smads And Ang-1/Tie2			EXCLI Journal	ISSN 1611-2156	View Document	https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5962903/

Back to Academic Year 2018-2019

Evaluation of Some Natural Polymers As Pharmaceutical Excipient For Sustained Drug Delivery System

Back to Academic Year 2018-2019

"Evaluation of Some Natural Polymers as Pharmaceutical Excipient For Sustained Drug Delivery System"

Avinash V.Dhobale *[1], Dr. M. Siddaiah[2]

[1]Research Scholar, Pharmaceutical Sciences, Bhagwant University Ajmer, Rajasthan, India.

[2] Departments of Pharmaceutics, Bhagwant University Ajmer, Rajasthan, India.

ABSTRACT:

The use of natural polymer as excipients in pharmaceutical sector is expanding day by day. Low cost, safety issues, availability, bio-degradable are the main causes that make them differ from other sources. Natural sources have wide range of varieties and characteristics. Excipients facilitate the formulation design and perform a wide range of functions to obtain desired properties for the finished drug product. Polysaccharide hydrocolloids including mucilages, gums and glucans are abundant in nature and commonly found in many higher plants. Currently, various plant polysaccharides have been studied for their diverse applications as excipients like binders, granulating agents, disintegrants, emulsifiers, suspending agents, gelling agents, mucoadhesive agents, matrix-formers, release retardants, enteric resistants, these polysaccharides constitute a structurally diverse class of biological macromolecules with a broad range of physicochemical properties which are widely used for various applications in pharmacy and medicine.

Key Words: Excipients, Modification, Natural Polymer, Sustained Drug Release.Gums and Mucilages

Back to Academic Year 2018-2019

Recent Advances In Mucoadhesive Buccal Drug Delivery Sy... Opportunities

International Journal of Advanced Pharmaceutical Sciences, Volume 1, Issue 08, Page 86-104
ISSN:2456-8147

RECENT ADVANCES IN MUCOADHESIVE BUCCAL DRUG DELIVERY SYSTEM AND ITS MARKETED SCOPE AND OPPORTUNITIES

Dhobale Avinash V.*, Nikose Karishma[2], Mrunal Pharate.R[3], Mahendra Datir[4]

*,2,3,4Loknete Shree Dada Patil Pharate College of Pharmacy Mandavgan Pharata, Pune.411041 ,Maharashtra, India .

An open access journal

Supporting Information:

Received: 22 July 2018
Accepted: 31 July 2018
Published: 07 August 2018

Competing Interests:
The authors have declared that no competing interests exist.

Corresponding author address

Dhobale Avinash V
Shree Dada Patil Pharate College of Pharmacy
Mandavgan Pharata,
Pune.411041 ,
Maharashtra, India .

Copyright: © 2018 Www.ijaps.net
Published under a Creative Commons Attribution 4.0

ABSTRACT

Extensive efforts have been focused on targeting a drug or drug delivery system in a particular region of the body for extended period of time, not only for local targeting of drug but also for better compliance of systemic drug delivery. Mucoadhesive characteristics are a factor of both the bioadhesive polymer and the medium in which the polymer will reside. Buccal dosage forms can be of Matrix or Reservoir types. However, this route could become a significant means for the delivery of a range of active agents in the coming years, if the barriers to buccal drug delivery are overcome. Mucoadhesive drug delivery system prolong the residence time of the dosage form at the site of application or absorption and facilitate an intimate contact of the dosage form with the underline absorption surface and thus contribute to improved and better therapeutic performance of the drug.

Buccal delivery involves the administration of the desired drug through the buccal mucosal membrane lining of the oral cavity. The objective of this article is to review buccal drug delivery by discussing the structure and enviorment of the oral mucosa and highlighting the mechanism of drug permeation and methodology in evaluating buccal formulations. This review also highlights a brief description of advantages, limitations of buccal drug delivery and theories of bioadhesion, mucoadhesive polymer, mechanism of buccal absorption along with mucoadhesive dosage form, factors affecting mucoadhesion.

Keywords: Mucosa, Mucoadhesion, Mucoadhesive Polymers, Mucoadhesive Drug Delivery System.

Back to Academic Year 2018-2019

Evaluation Of Antiulcer Activity Of Aconitum Heteroph

WORLD JOURNAL OF PHARMACY AND PHARMACEUTICAL SCIENCES
SJIF Impact Factor 6.647
Volume 7, Issue 2, 819-839 Research Article ISSN 2278 – 4357

EVALUATION OF ANTIULCER ACTIVITY OF ACONITUM HETEROPHYLLUM ON EXPERIMENTAL ANIMAL

Vishal R. Rasve*, Vivek V. Paithankar[1], Mrunal K Shirsat[2], Avinash V. Dhobale[3]

Assist. Professor, Vidybharti college of Pharmacy, Amravati 444 602.
Principal, SVP College of Pharmacy, Hatta.
Assist. Professor, SVP College of Pharmacy, Hatta.

Article Received on
24 November 2017,
Revised on 15 Dec. 2017,
Accepted on 06 Jan. 2018
DOI: 10.20959/wjpps20182-10858

***Corresponding Author**
Vishal R. Rasve
Assist. Professor, SVP College of Pharmacy, (B.Pharm) Hatta.dist. Hingoli 431705.

ABSTRACT

In present era peptic ulcer is a worldwide problem and its prevalence is quite high in India. Some present data shows that in different part of India its occurrence is 4 to 10 % per thousand populations. The exact etiology of peptic ulcer is not known, the disease results in chronic suffering, loss of working hours and occasional fatality. Smoking habit, alcoholism, stress & spicy food make the severity of the disease which leads serious complication of ulcer. Aconitum heterophyllum is one of the most commonly use in different medical conditions has been documented. Traditionally, Aconitum heterophyllum roots or tubers are beneficial in the treatment of diseases related to lungs & airways, blood, skin, stomach, intestines, and liver. As no reports are available on the possible antiulcer effects of Aconitum heterophyllum. The present work was carried out to antiulcer activity of Aconitum heterophyllum by cold stress induced gastric ulcers in wister albino Test drug Aconitum heterophyllum (low dose) and Aconitum heterophyllum (high dose) have shown result in comparison to standard drug omeparazole and control drug. Thus study has provided documentary evidence for antiulcer property of Aconitum heterophyllum for its activity.

KEYWORDS: Aconitum heterophyllum, peptic ulcer, antiulcer property.

INTRODUCTION

Charaka made fifty groups of ten herbs each of which, according to him, would suffice an ordinary physician's need. Similarly, Sushrutha arranged 760 herbs in 7 distinct sets based on their common properties. A large portion of the Indian population even today, depends on the

Back to Academic Year 2018-2019

WORLD JOURNAL OF PHARMACY AND PHARMACEUTICAL SCIENCES
SJIF Impact Factor 6.647
Volume 7, Issue 1, 1230-1245 Research Article ISSN 2278 – 4357

DESIGN AND IN-VITRO CHARACTERIZATION OF GASTRORETENTIVE FLOATING TABLETS CONTAINING GLICLAZIDE

Deepak A. Joshi*, Dr. Sidramappa M. Chickpetty, Gunesh N. Dhembre and Avinash V. Dhobale

Dept of Pharmaceutics, SVP College of Pharmacy, Hatta-431705 Maharashtra, India.

Article Received on
11 Nov. 2017,
Revised on 30 Nov. 2017,
Accepted on 21 Dec. 2017
DOI: 10.20959/wjpps20181-10825

***Corresponding Author**
Deepak A. Joshi
Dept of Pharmaceutics, SVP College of Pharmacy, Hatta-431705 Maharashtra, India.

ABSTRACT

The current work is aimed to design a novel oral gastroretentive floating tablets of Gliclazide by using combination of HPMC, carbopole 934 (CP) and guar gum (GG) for safe and effective management of non-insulin dependent (Type II) diabetes mellitus for better control of blood glucose level by enhancing its oral bioavailability and also to improve the patient compliance. The floating tablets containing 50mg of Gliclazide was prepared by direct compression technology. All the tablet formulations were evaluated for physical properties; total buoyancy and buoyancy lag time and in vitro drug release studies in pH 1.2 buffer solution. All the tablets prepared with GG / CBP alone and their combination as matrix material fails to meet the requirements of desired floating ability, sustaining and complete drug release. Among the various ratios (5:1, 4:2 and 3:3) of HPMC:GG, the ratio 3:3 meets the desired requirement (floated for 12h with a floating lag time of 30sec) and released major portion of the drug (98.64%) in 12h. The tablet formulation containing HPMC and CBP, the ratio 4:2 was found to be suitable for designing floating tablets as it has shown less floating lag time (43sec), floated for the testing period of 12h and released 97.925 of Gliclazide. The optimized tablet batch formulations showed no change in drug content or in vitro release pattern after storage at 40° C / 75% RH for 30 days. The FTIR studies ruled out incompatibility between the drug (Gliclazide) and non drug components used in the tablets.

KEYWORDS: Gliclazide, carbopole, guar gum, FDDS, gastric residence time, diabetes mellitus.

Back to Academic Year 2018-2019

Mouth Dissolving Strips

International Journal of Advanced Pharmaceutical Sciences, Volume 1, Issue 12, Page 09– 31 eISSN: 2456-8147

Mouth dissolving strips– a review

Sneha T.Chaudhari [1]*, Gautam Girendrakumar [2], Shivaji Kanhere[3], Avinash Dhobale[4]

*Department of pharmaceutics, Bhagwant University, Ajmer (Rajasthan),India

An open access journal

Supporting Information:

Received: 05 October 2018
Accepted: 28 November 2018
Published: 12 December 2018

Competing Interests:
The authors have declared that no competing interests exist.

Corresponding author address
Sneha T.Chaudhari ,
Department of pharmaceutics,
Bhagwant University, Ajmer (Rajasthan),
India

Copyright: © 2018 Www.ijaps.net
Published under a Creative Commons Attribution 4.0

ABSTRACT

In the recent years most useful route of drug delivery system is the oral route in all types of patient. Mouth dissolving strips is the type of drug delivery system which when placed in the oral cavity, disintegrate or dissolve within few seconds without the intake of water. The advantages of mouth dissolving strips are the administration to pediatric and geriatric patient. This technology has been used for local action, rapid release products.These films have a potential to deliver the drug systemically through intragastric, sublingual or buccal route of administration and also has been used for local action. The mouth dissolving strips are available in various shapes, size and thickness.This article is an overview of mouth dissolving strips encompassing materials used in mouth dissolving strips, manufacturing process, applications, business technologies and future business prospects of this technology

Keywords: Strips, Mouth Dissolving Strips, Rapid Release. Dose, Drug delivery

Back to Academic Year 2018-2019

Formulation And Evaluation Of Fast Disintegrating Tablet Of Carvedilol

IJPPR
INTERNATIONAL JOURNAL OF PHARMACY & PHARMACEUTICAL RESEARCH
An official Publication of Human Journals
ISSN 2349-7203
HUMAN

Human Journals
Research Article
August 2018 Vol.:13, Issue:1
© All rights are reserved by Kharde Sagar N et al.

Formulation and Evaluation of Fast Disintegrating Tablet of Carvedilol

Kharde Sagar N[1*], Satpute Vivek M [2], Dhobale Avinash V [3], Wani Rakesh M[4]

Department of Quality Assurance at Pravara Rural College of Pharmacy, Pravaranagar Maharashtra, India.

Submission: 20 July 2018
Accepted: 27 July 2018
Published: 30 August 2018

www.ijppr.humanjournals.com

Keywords: FDT, Carvedilol, Disintegration, Lyophilization, Dimethylsulphoxide

ABSTRACT

The oral route of administration is considered as the most widely accepted route because of its convenience of self-administration, compactness and easy manufacturing. Fast disintegrating tablets (FDT) has enormously increased as it has significant impact on the patient compliance some novel FDT technology allows high drug loading, have an acceptable taste, offer a pleasant mouth felling, leaving minimal residue in the mouth after oral administration. FDT have been investigated for their potential in improving bioavailability of poorly soluble drug through enhancing the dissolution profile of the drug and hepatic metabolism drugs. The present study was carried out to design and evaluation of fast disintegrating tablets of carvedilol for the effective management of angina pectoris, hypertension etc. In view of substantial first pass effect and its shorter plasma half-life. The fast disintegrating tablets of carvedilol were prepared by enhancing the solubility of carvedilol using β-cyclodextrin solid dispersion technology. Solubility of carvedilol is enhanced by preparing solid dispersion with β-cyclodextrin 1:4 ratio. DSC and IR spectroscopy data showed the characterization of drug, excipient, compatibility of drug and solid dispersion with excipients, gave evidence of solid dispersion formation and UV absorption spectra shows enhancement of solubility. Various preformulation batches (f1-f10) formulated by direct compression method using different concentration of polymer such as ac-di-sol and pearlitol and it was studied for pre and post compression evaluation. formulation with AC-DI-SOL and pearlitol batch no. F9 shows excellent result with disintegration time of 105 sec, drug content of 99.35%, and greater dissolution rate 98.32%at 40 min.

Back to Academic Year 2018-2019

WORLD JOURNAL OF PHARMACY AND PHARMACEUTICAL SCIENCES
SJIF Impact Factor 6.647
Volume 7, Issue 2, 819-839 Research Article ISSN 2278 – 4357

EVALUATION OF ANTIULCER ACTIVITY OF ACONITUM HETEROPHYLLUM ON EXPERIMENTAL ANIMAL

Vishal R. Rasve*, Vivek V. Paithankar[1], Mrunal K Shirsat[2], Avinash V. Dhobale[3]

Assist. Professor, Vidybharti college of Pharmacy, Amravati 444 602.
Principal, SVP College of Pharmacy, Hatta.
Assist. Professor, SVP College of Pharmacy, Hatta.

Article Received on
24 November 2017,
Revised on 15 Dec. 2017,
Accepted on 06 Jan. 2018
DOI: 10.20959/wjpps20182-10858

***Corresponding Author**
Vishal R. Rasve
Assist. Professor, SVP College of Pharmacy, (B.Pharm) Hatta.dist. Hingoli 431705.

ABSTRACT

In present era peptic ulcer is a worldwide problem and its prevalence is quite high in India. Some present data shows that in different part of India its occurrence is 4 to 10 % per thousand populations. The exact etiology of peptic ulcer is not known, the disease results in chronic suffering, loss of working hours and occasional fatality. Smoking habit, alcoholism, stress & spicy food make the severity of the disease which leads serious complication of ulcer. Aconitum heterophyllum is one of the most commonly use in different medical conditions has been documented. Traditionally, Aconitum heterophyllum roots or tubers are beneficial in the treatment of diseases related to lungs & airways, blood, skin, stomach, intestines, and liver. As no reports are available on the possible antiulcer effects of Aconitum heterophyllum. The present work was carried out to antiulcer activity of Aconitum heterophyllum by cold stress induced gastric ulcers in wister albino Test drug Aconitum heterophyllum (low dose) and Aconitum heterophyllum (high dose) have shown result in comparison to standard drug omeparazole and control drug. Thus study has provided documentary evidence for antiulcer property of Aconitum heterophyllum for its activity.

KEYWORDS: Aconitum heterophyllum, peptic ulcer, antiulcer property.

INTRODUCTION

Charaka made fifty groups of ten herbs each of which, according to him, would suffice an ordinary physician's need. Similarly, Sushrutha arranged 760 herbs in 7 distinct sets based on their common properties. A large portion of the Indian population even today, depends on the

Back to Academic Year 2018-2019

International Journal of Current Advanced Research
ISSN: O: 2319-6475, ISSN: P: 2319-6505, Impact Factor: 6.614
Available Online at www.journalijcar.org
Volume 7; Issue 7(E); July 2018; Page No. 14170-14174
DOI: http://dx.doi.org/10.24327/ijcar.2018.14174.2561

Research Article

FORMULATION AND EVALUATION OF ATORVASTATIN CALCIUM ORAL DISPERSIBLE TABLET

Satpute Vivek M., Shirsat Mrunal K., Dhobale.Avinash V., Wani Rakesh M and Kharde Sagar N

Loknete Shri Dadapatil Pharate College of Pharmacy Mandavgan pharata, Pune

ARTICLE INFO

Article History:

Received 5th April, 2018
Received in revised form 24th May, 2018 Accepted 20th June, 2018
Published online 28th July, 2018

Key words:

Atorvastatin Calcium, oral dispersible , Crospovidone, direct compression method.

ABSTRACT

The objective of the study was to develop the oral dispersible tablets of various concentrations of polymers on in-vitro release rate from the prepared Atorvastatin calcium tablets. The effect of drug to polymer ratio on the in vitro drug release behavior was significant. Formulation F5 showed better in vitro drug release and this indicates the ideal drug, polymer and excipients combination formulations. In vitro dissolution results showed that maximum cumulative % drug release was more in formulation F5 when compared to formulation F1 to F9.Stability studies were conducted formulations F5 stored at 250C/60% RH and 400C/75% RH for 30 days. Various parameters like hardness, friability, drug content uniformity, in vitro disintegration, wetting time were analyzed at a time of interval of 10 days till a period of 30 days. Not much variation or change was observed in any parameters throughout the study period. Best –selected formulations F5 found to be stable. Hence it is concluded that The prepared tablets disintegrate in seconds without need of water and enhance the absorption, this leads to increased bioavailability of Atrovastatin calcium.

The results of stability study on Formulation F5 after a period of one month indicates that the formulation was stable. Based on result it is concluded that formulated oral dispersible tablet of atrovastatin may have wide acceptance as compared to conventional dosage form.

Copyright©2018 ***Satpute Vivek et al.*** *This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.*

INTRODUCTION

Solid medicaments may be administered orally as powders, pills, cachets, capsules or tablets. These dosage forms contain a quantity of drug which is given as a single unit and they are known collectively as solid unit dosage forms, even in the case of sustained action preparations which, technically, contain the equivalent of several normal doses of drug.

The treatment of acute or chronic illness has been achieved by delivery of drug to the patients for many years. These drug delivery systems include capsules, tablets, injectables, creams, liquids, aerosol, suspensions and ointments. The term drug delivery can be defined as technique that is used to get therapeutic agent inside the human body. [2,3]

Oral drug delivery has been known for decades as the most widely utilized route of administration among all the routes that have been explored for the systemic delivery of drugs via different dosage forms. Oral routes of drug administration have wide acceptance up to 50-60% of total dosage forms. Solid dosage forms are popular because of ease of administration, accurate dosage, self medication, pain avoidance and most importantly the patient compliance.

Corresponding author:* **Satpute Vivek
Loknete Shri Dadapatil Pharate College of Pharmacy
Mandavgan pharata, Pune

The most popular solid dosage forms are being tablets and capsules.

Atorvastatin Calcium Structure

MATERIALS AND METHODS

Gift samples of standards Atorvastatin Calcium were provided Wokhardt Pharmaceutical at, Aurangabad. Crospovidone, Croscarmellose sodium, Sodium starch glycolate, Lactose, Microcrystalline cellulose, Magnesium Stearate, were provided Lupin Pharmaceutical at, Aurangabad.

Formulation of Orally Disintegrating Tablets

Weigh all the ingredients accurately according. Mix all the ingredients geometrically except Talc and Magnesium Stearate. Then lubricate the blend with Talc, Magnesium Stearate and passed through sieve no 40 .The blend was compressed using rotary tablet machine-10 station with 6mm flat punch. Each tablet contains 10mg Atrovastatin and other pharmaceutical ingredients as in Table no 1.

Indo American Journal of Pharmaceutical Research, 2018 ISSN NO: 2231-6876

INDO AMERICAN JOURNAL OF PHARMACEUTICAL RESEARCH

"FORMULATION AND EVALUATION OF CARBAMAZEPINE NANOEMULSION FOR BRAIN TARGETED DRUG DELIVERY VIA INTRANASAL ROUTE"

Nagare Sonali .A*, Dhobale Avinash.V, Najan Anita.A

Loknete Shri Dadapatil Pharate College of Pharmacy, Mandavgan pharata,Shirur,Dist Pune,412211.

ARTICLE INFO

Article history
Received 16/07/2018
Available online
31/07/2018

Keywords
Carbamazepine,
Nanoemulsion,
Intranasal To Brain.

ABSTRACT

Intranasal drug administration is receiving increased attention as adelivery method for by-passing the blood–brain barrier and rapidly targeting therapeutics to the brain or CNS. The objective of the present study was to selected carbamazepine nanoemulsion for nose-to-brain delivery. Carbamazepine nanoemulsion (NE) formulation were successfully prepared by the spontaneous emulsification method (titration method) using Capmul MCM as the oil, Tween-80 as surfactant, and PEG-600 as co-surfactant phase on the basis of solubility studies. The nanoemulsion formulation containing 7.35% oil, 66.18% Smix ratio (3:1 Tween-80:PEG-600 ratio),26.47% (v/v) aqueous phase that displayed an optical transparency of 99.42±0.81%, globule size of 71.70±3.06 nm, and polydispersity index of 0.256±0.002. The selected Carbamazepine nanoemulsion was characterized, and the *in-vitro* drug release and *in-vivo* nasal absorption of drug from the selected formulation were evaluated in rats. *In-vitro* and ex vivo permeation studies showed an initial burst of drug release at 60 min and Carbamazepine nanoemulsion show drug release up to 5 h. In vivo pharmacokinetic studies in rats showed that Carbamazepine.

Corresponding author
Nagare Sonali
Loknete Shri Dadapatil Pharate College of Pharmacy,
Mandavgan Pharata, Shirur, Dist Pune, 412211.
sonalinagare93@gmail.com

*Please cite this article in press as **Nagare Sonali** et al. "Formulation and Evaluation of Carbamazepine Nanoemulsion for Brain Targeted Drug Delivery Via Intranasal Route". Indo American Journal of Pharmaceutical Research.2018:8(07).*

Copy right © 2018 This is an Open Access article distributed under the terms of the Indo American journal of Pharmaceutical Research, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INDO AMERICAN JOURNAL OF PHARMACEUTICAL RESEARCH

RECENT ADVANCES IN PILOT PLANT SCALE UP TECHNIQUES - A Review

AvinashV.Dhobale*[1], Arun M.Mahale[2], Mrunal Shirsat[3], Shriram Pethkar[4], Vijay Chakote[5]
[1]*LCOP Hasegaon.*
[2]*SNIOP,Pusad.*
[3]*Dr. N. JPaulbudhe College of Pharmacy Ahamadnagar.*
[4]*Latur College of Pharmacy (B.Pharm), Hasegaon.*
[5]*Atsvp College of Pharmacy Hatta.*

ARTICLE INFO
Article history
Received 21/04/2018
Available online
12/05/2018

Keywords
Pilot Plant Technique,
Solid Dosage Form,
Tablets,
Compression.

ABSTRACT
Pilot scale up techniques for solid dosage form will provide guide line forthe manufacture of large scale process and this will play a pivotal role inlarge scale manufacturing. The parameters such as granulation feed rate,compression parameters, temperature and rate of drying will have acritical role in development of any solid dosage form.Pilot plant is a relative term in the sense that pilot plants are typically smaller than full-scale production plants, but are built in a range of sizes. Also, as pilot plants are intended for learning, they typically are more flexible, possibly at the expense of economy. Some pilot plants are built in laboratories using stock lab equipment, The past two decades particular have witnessed amazing inventions and innovations in pharmaceutical research, resulting in the ability to produce new drugs faster than even before. The new drug applications (NDAs) and abbreviated new drug applications (ANDA) are all-time high. The preparation of several clinical batches in the pilot plant provides its personnel with the opportunity to perfect and validate the process.

Corresponding author
AvinashV. Dhobale
(Department of Pharmacetics)
Assistant Professorat Latur College of Pharmacy, Hasegaon
dhobaleavi@gmail.com
9604477418

*Please cite this article in press as **AvinashV. Dhobale** et al. Recent Advances in Pilot Plant Scale up Techniques - A Review. Indo American Journal of Pharmaceutical Research.2018:8(04).*

Copy right © 2018 This is an Open Access article distributed under the terms of the Indo American journal of Pharmaceutical Research, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Back to Academic Year 2018-2019

Design, Development And Characterization Oflosartan Potas

WORLD JOURNAL OF PHARMACY AND PHARMACEUTICAL SCIENCES
SJIF Impact Factor 6.647
Volume 7, Issue 1, 1349-1362 Research Article ISSN 2278 – 4357

DESIGN, DEVELOPMENT AND CHARACTERIZATION OF LOSARTAN POTASSIUM SUSTAINED RELEASE MATRIX TABLET

Avinash V. Dhobale[1*], Arun M. Mahale[1], Mrunal K. Shirsart[2], Deepak A. Joshi[3], Gunesh N. Dhembre[4] and Mahesh M. Thakare[5]

[1]Assist. Professor Sudhakarrao Naik Institute of Pharmacy, Pusad.
[2]Principal, SVP College of Pharmacy, Hatta.
[3]Assist. Professor SVP College of Pharmacy, Hatta.

Article Received on
14 November 2017,
Revised on 05 Dec. 2017,
Accepted on 26 Dec. 2017
DOI: 10.20959/wjpps20181-10845

***Corresponding Author**
Dr. Avinash V. Dhobale
Assist. Professor
Sudhakarrao Naik institute of Pharmacy, Pusad.

ABSTRACT

The aim of this research work was to formulate and develop a fixed Dose Combination product in a two different strength using same blend for both the strengths of tablet as a SR tablet formulation. In the tablet, Extended Release layer consist of Antihypertensive Drug belonging to class β-selective adrenergic blocking agent without partial agonist or membrane stabilizing properties. Extended release preparation provides sustained release and reduces the chances of tough related side effects. In selected cases of extended release preparation of this drug used in treatment of hypertension and congestive heart failure. The clinical studies have shown beneficial role of this drug as an extended release preparation. The main objective of the present study was to develop, formulate and evaluate a matrix tablet by using hydrophilic natural retardant polymers which would retard drug release in upper GI tract and should start releasing the drug when it reaches the alkaline environment of small intestine. Metolose 90 sh and xanthan gum were investigated as the model hydrophilic retardant polymers. Wet granulation method was used for preparation of sustained release matrix tablets. Nine batches of tablets were prepared. The prepared tablets were subjected for pharmacopoeial and non-pharmacopoeial evaluation parameters including loose and tapped bulk density, compressibility index, hausner ratio, angle of repose, friability, hardness, thickness, weight variation, % drug content and in-vitro drug release studies. It can be concluded that the combination of hydrophilic polymers that are retardant in nature are better suited for sustained and controlled drug delivery system than the hydrophilic polymer alone.

Back to Academic Year 2018-2019

Design And In-Vitro Characterization Of Gastro retentive Floating Tablets Containing Gliclazide

WORLD JOURNAL OF PHARMACY AND PHARMACEUTICAL SCIENCES
SJIF Impact Factor 6.647
Volume 7, Issue 1, 1230-1245 Research Article ISSN 2278 – 4357

DESIGN AND IN-VITRO CHARACTERIZATION OF GASTRORETENTIVE FLOATING TABLETS CONTAINING GLICLAZIDE

Deepak A. Joshi*, Dr. Sidramappa M. Chickpetty, Gunesh N. Dhembre and Avinash V. Dhobale

Dept of Pharmaceutics, SVP College of Pharmacy, Hatta-431705 Maharashtra, India.

Article Received on
11 Nov. 2017,
Revised on 30 Nov. 2017,
Accepted on 21 Dec. 2017
DOI: 10.20959/wjpps20181-10825

***Corresponding Author**
Deepak A. Joshi
Dept of Pharmaceutics, SVP College of Pharmacy, Hatta-431705 Maharashtra, India.

ABSTRACT

The current work is aimed to design a novel oral gastroretentive floating tablets of Gliclazide by using combination of HPMC, carbopole 934 (CP) and guar gum (GG) for safe and effective management of non-insulin dependent (Type II) diabetes mellitus for better control of blood glucose level by enhancing its oral bioavailability and also to improve the patient compliance. The floating tablets containing 50mg of Gliclazide was prepared by direct compression technology. All the tablet formulations were evaluated for physical properties; total buoyancy and buoyancy lag time and in vitro drug release studies in pH 1.2 buffer solution. All the tablets prepared with GG / CBP alone and their combination as matrix material fails to meet the requirements of desired floating ability, sustaining and complete drug release. Among the various ratios (5:1, 4:2 and 3:3) of HPMC:GG, the ratio 3:3 meets the desired requirement (floated for 12h with a floating lag time of 30sec) and released major portion of the drug (98.64%) in 12h. The tablet formulation containing HPMC and CBP, the ratio 4:2 was found to be suitable for designing floating tablets as it has shown less floating lag time (43sec), floated for the testing period of 12h and released 97.925 of Gliclazide. The optimized tablet batch formulations showed no change in drug content or in vitro release pattern after storage at 40° C / 75% RH for 30 days. The FTIR studies ruled out incompatibility between the drug (Gliclazide) and non drug components used in the tablets.

KEYWORDS: Gliclazide, carbopole, guar gum, FDDS, gastric residence time, diabetes mellitus.

Back to Academic Year 2018-2019

Formulation And Evaluation Of Rosuvastatin Oral Dispersible Tablet Under A Creative Commons

International Journal of Advanced Pharmaceutical Sciences, Volume 1, Issue 08, Page 105-121
ISSN:2456-8147

FORMULATION AND EVALUATION OF ROSUVASTATIN ORAL DISPERSIBLE TABLET

Satpute Vivek M*, K. Shirsat , Rakesh M. Wani, Avinash V. Dhobale , Sagar N. Kharde

* Loknete Shri Dadapatil Pharate College of Pharmacy Mandavgan Pharata, Pune, Maharashtra, India

An open access journal

Supporting Information:

Received: 31 July 2018
Accepted: 05 August 2018
Published: 07 August 2018

Competing Interests:
The authors have declared that no competing interests exist.

Corresponding author address

*Satpute Vivek M
(Department of Pharmaceutics)
Loknete Shri Dadapatil Pharate College of pharmacy
Mandavgan Pharata,
Pune, 412211
Maharashtra, India

Copyright: © 2018 Www.ijaps.net
Published under a Creative Commons Attribution 4.0

ABSTRACT

The objective of the study was to develop the oral dispersible tablets of various concentrations of polymers on in-vitro release rate from the prepared rosuvastatin tablets. The effect of drug to polymer ratio on the in vitro drug release behavior was significant. Formulation F5 showed better in vitro drug release and this indicates the ideal drug, polymer and excipients combination. Formulations In vitro dissolution results showed that maximum cumulative % drug release was more in formulation F5 when compared to formulation F1 to F9.Stability studies were conducted formulations F5 stored at 250C/60% RH and 400C/75% RH for 30 days. Various parameters like hardness, friability, drug content uniformity, in vitro disintegration, wetting time were analyzed at a time of interval of 10 days till a period of 30 days. Not much variation or change was observed in any parameters throughout the study period. Best –selected formulations F5 found to be stable. Hence it is concluded that the prepared tablets disintegrate in seconds without need of water and enhance the absorption, this leads to increased bioavailability of Rosuvastatin.

The results of stability study on Formulation F5 after a period of one month indicates that the formulation was stable. Based on result it is concluded that formulated oral dispersible tablet of Rosuvastatin may have wide acceptance as compared to conventional dosage form.

Key Words: Rosuvastatin, Oral dispersible, Crospovidone, Direct Compression Method.

Back to Academic Year 2018-2019

International Journal of Advanced Pharmaceutical Sciences, Volume 1, Issue 04, Page 51-66
ISSN:2456-8147

A PROJECT ON HUMAN PAPILLOMAVIRUS - A REVIEW

Satpute Vivek [1*], Dhobale Avinash [2], Wani Rakesh [3], Shirsat Mrunal[4]

*Loknete Shri Dadapatil Pharate College of Pharmacy, Mandavgan pharata, Tal- Shirur, Dist- Pune,412211 ,Maharashtra,India

An open access journal

Supporting Information:

Received: 18 July 2018
Accepted: 22 July 2018
Published: 23 July 2018

Competing Interests:
The authors have declared that no competing interests exist.

Corresponding author address

Satpute Vivek
Loknete Shri Dadapatil Pharate College of Pharmacy,
Mandavgan pharata,
Tal- Shirur, Dist Pune,412211 ,
Maharashta,India

Copyright: © 2018 Www.ijaps.net
Published under a Creative Commons Attribution 4.0

ABSTRACT

Human Papilloma Viruses (HPV) are the etiological agents of cervical and different ano-genital malignancies. Over 100 differing kinds of HPVs are familiar up to now, and each one target epithelial tissues for infection. one-third of HPV varieties specifically infect the genital tract and a group of these is that the activating agents of anogenital cancers. different HPV varieties that infect the sex organ tract induce benign hyperproliferative lesions or sex organ warts. Around eightieth of sexually active men and ladies will contract the HPV virus at some levels throughout their lifetime. Annually there are 14 million new diagnosed cases of HPV. There are additional or less 79 million men/women actively infected with the virus at any purpose in time. The productive life cycle of HPV is connected to tissue differentiation. HPV virus affects the basal cells of stratified epithelia. It then establishes their genomes as a multicopy nuclear episome. In these cells, viral DNA is replicated in conjunction with cellular chromosomes. Following biological process one among the female offspring cells migrates removed from the basal layer and undergoes differentiation. Particle production is restricted to differentiated cells. The infected basal cells can survive for up to several decades or until and unless the system clears the infection. The L1 and L2 proteins kind polyhedron capsids for offspring particle generation. The characterization of the cellular targets of these organism proteins and thus the mechanisms control the differentiation-dependent organism life cycle stay active are for the study of these important human pathogens. This review deals with numerous aspects of Human Papilloma Viruses (HPV)

Keywords: Cancer, HPV, Cervix, Proteins, Infection, Warts

Back to Academic Year 2018-2019

IJPPR
INTERNATIONAL JOURNAL OF PHARMACY & PHARMACEUTICAL RESEARCH
An official Publication of Human Journals
ISSN 2349-7203
HUMAN

Human Journals
Research Article
August 2018 Vol.:13, Issue:1
© All rights are reserved by Kharde Sagar N et al.

Formulation and Evaluation of Fast Disintegrating Tablet of Carvedilol

Kharde Sagar N[1*], Satpute Vivek M [2], Dhobale Avinash V [3], Wani Rakesh M[4]

Department of Quality Assurance at Pravara Rural College of Pharmacy, Pravaranagar Maharashtra, India.

Submission: 20 July 2018
Accepted: 27 July 2018
Published: 30 August 2018

www.ijppr.humanjournals.com

Keywords: FDT, Carvedilol, Disintegration, Lyophilization, Dimethylsulphoxide

ABSTRACT

The oral route of administration is considered as the most widely accepted route because of its convenience of self-administration, compactness and easy manufacturing. Fast disintegrating tablets (FDT) has enormously increased as it has significant impact on the patient compliance some novel FDT technology allows high drug loading, have an acceptable taste, offer a pleasant mouth felling, leaving minimal residue in the mouth after oral administration. FDT have been investigated for their potential in improving bioavailability of poorly soluble drug through enhancing the dissolution profile of the drug and hepatic metabolism drugs. The present study was carried out to design and evaluation of fast disintegrating tablets of carvedilol for the effective management of angina pectoris, hypertension etc. In view of substantial first pass effect and its shorter plasma half-life. The fast disintegrating tablets of carvedilol were prepared by enhancing the solubility of carvedilol using β-cyclodextrin solid dispersion technology. Solubility of carvedilol is enhanced by preparing solid dispersion with β-cyclodextrin 1:4 ratio. DSC and IR spectroscopy data showed the characterization of drug, excipient, compatibility of drug and solid dispersion with excipients, gave evidence of solid dispersion formation and UV absorption spectra shows enhancement of solubility. Various preformulation batches (f1-f10) formulated by direct compression method using different concentration of polymer such as ac-di-sol and pearlitol and it was studied for pre and post compression evaluation. formulation with AC-DI-SOL and pearlitol batch no. F9 shows excellent result with disintegration time of 105 sec, drug content of 99.35%, and greater dissolution rate 98.32%at 40 min.

Back to Academic Year 2018-2019

Journal of Pharmacognosy and Phytochemistry 2018; 7(6): 2489-2492

Journal of Pharmacognosy and Phytochemistry
Available online at www.phytojournal.com

E-ISSN: 2278-4136
P-ISSN: 2349-8234
JPP 2018; 7(6): 2489-2492
Received: 19-09-2018
Accepted: 21-10-2018

Mohan Agrawal
Assistant Professor, Department of Pharmacognosy, Sitabai Thite College of Pharmacy, Shirur, Pune Maharashtra, India

Dr. Anilkumar Aher
MVP's College of Pharmacy, Nashik, Maharashtra, India

Dr. Subodh Pal
MVP's College of Pharmacy, Nashik, Maharashtra, India

Dr. Deelip Derle
MVP's College of Pharmacy, Nashik, Maharashtra, India

Correspondence
Mohan Agrawal
Assistant Professor, Department of Pharmacognosy, Sitabai Thite College of Pharmacy, Shirur, Pune Maharashtra, India

Isolation and characterization of phytoconstituents from petroleum ether extract of *Momordica cochinchinensis* fruits

Mohan Agrawal, Dr. Anilkumar Aher, Dr. Subodh Pal and Dr. Deelip Derle

Abstract
The purpose of the study is to isolate and characterize the chemical constituents from fruits of *Momordica cochinchinensis*. The principal theme of the study is to develop applied chromatographic techniques for the separation, isolation and detection of the compounds. The petroleum ether extract of fruits of *Momordica cochinchinensis* was saponified and unsaponifiable matter was subjected to column chromatography and elution of column was carried out by Petroleum ether (100%) with increasing concentration of ethyl acetate for the separation of phytoconstituents. The isolated compounds were characterized and analyzed by physical characteristics, IR, NMR and Mass spectroscopy. Three known compounds lupeol, β-amyrin and β-sitosterol were determined for the first time from fruits of *Momordica cochinchinensis*. From the present study, it is concluded that the plant consists of phytosterols which can be isolated and characterized by chromatographical and spectroscopical method.

Keywords: *Momordica cochinchinensis,* phytosterols, lupeol, β-amyrin and β-sitosterol

Introduction
From thousands of years Natural products have been used by human societies. Natural sources have provided considerable value to the pharmaceutical industry over the past half century. Research studies leading to extraction, isolation and biological study of plant constituents have formed the major field of study. Various leads from plant sources were taken for discovering the new active therapeutic agents. Hence herbal medicine has played important role in managing the health conditions like diabetes, hypertension, inflammation, obesity, cancer, etc [1].
The plant *Momordica cochinchinensis* (Gac) is a Southeast Asian fruit found throughout the region from Southern China to Northeastern Australia, mostly Vietnam and throughout India. It grows on dioecious vines and usually collected from fence climbers or from wild plants [2]. It is reported that *Momordica cochinchinensis* Spreng, is one of the special fruits containing extraordinarily high levels of carotenoids, especially β -carotene (more than 16 mg/100 g) and lycopene (more than 50 mg/100 g), mainly in the red aril [3]. It is also reported to contain a protein that may inhibit the proliferation of cancer cells and also β-carotene with several phytonutrients, Vit-E, fatty acids, carbohydrates, flavonoidal glycosides [4].
Traditionally, Gac has been used as both food and medicine and promotes healthy vision by relief of dry eyes. It also possesses antioxidant, anti-microbial and antidiabetic properties. The seeds are considered to be good for cough and pains in the chest [5, 6].
But these studies are not enough for identifying and characterizing the bioactive compounds in the plant. The purpose of the study is to identify and characterize the bioactive principles from fruits of *Momordica cochinchinensis*.

Materials and Methods
Plant material: The fruits of *Momordica cochinchinensis* was collected from rural area of Kolkata. The herbarium of the *Momordica cochinchinensis* was authenticated by Botanical Survey of India, Pune Voucher specimen (MA 01) was deposited in library.

Preparation of Extracts: Fruits of *Momordica cochinchinensis* was extracted by Soxhlet extractor with Pet ether and macerated with ethanol and water successively.

Storage of Extracts: All the extracts were stored in tightly closed glass bottles in refrigerator at 2-8 ^{0}C.

Back to Academic Year 2018-2019

E-ISSN: 2278-3229

IJGHC, September 2018 – November 2018; Sec. B; Vol.7, No.4, 446-455.
DOI: 10.24214/IJGHC/HC/7/4/44655.

International Journal of Green and Herbal Chemistry

An International Peer Review E-3 Journal of Sciences
Available online at www.ijghc.com
Section B: Herbal Chemistry

Research Article

CODEN (USA): IJGHAY

Anti-inflammatory activity of *Momordica cochinchinensis* and *Momordica balsamina* fruit extracts

Mr. Mohan R Agrawal[1]*, Dr. Anilkumar N Aher[2], Dr. Subodh C Pal[2] and Dr. Deelip V Derle[2]

[1]Sitabai Thite College of Pharmacy, Shirur, Pune (District), Maharashtra-412210, India.
[2] MVP's College of Pharmacy, Nashik, Maharashtra, India

Received: 14 September 2018; **Revised:** 13 October 2018; **Accepted:** 24 October 2018

Abstract: In the present study fruit extracts of *Momordica cochinchinensis* and *Momordica balsamina (Cucurbitaceae)* were investigated for anti-inflammatory activity by Carrageenan induced hind paw edema and cotton pellet granuloma animal model. The extracts were prepared successively using powdered material with Petroleum ether, Ethanol and Water, concentrated under vacuum and were evaluated for anti-inflammatory activity at three dose level (100, 200 and 400 mg/kg). In Carrageenan-induced paw edema method, oral administration of Petroleum ether extracts of both the plants at the dose of 100 ($p < 0.05$), 200 ($p < 0.01$) and 400 mg/kg ($p < 0.001$) significantly inhibited the edema formation. The highest percentage inhibition 47.36 % and 63.06 % at sixth hour was shown by petroleum ether extract of both plants at the dose of 400 mg/kg respectively. In cotton pellet granuloma method, Pet ether extract at the dose of 100 ($p<0.05$), 200 and 400 mg/kg ($p<0.01$) of both plant material have shown significant anti-inflammatory activity and was comparable to standard drug Diclofenac (10 mg/kg) in reducing the granuloma formation. It is concluded that Pet ether extracts of both plant material have significant anti-inflammatory effect, may be attributed to

Back to Academic Year 2018-2019

Analgesic Activity of Momordica Cochinensis & Momordica Balsmina Fruit Extracts

ORIGINAL ARTICLE

Analgesic activity of *Momordica cochinchinensis* and *Momordica balsamina* fruit extracts

Mohan R. Agrawal[1]*, Anilkumar N. Aher[2], Subodh C. Pal[2], Deelip V. Derle[2]

[1]Department of Pharmacognosy, Sitabai Thite College of Pharmacy, Pune, Maharashtra, India, [2]Department of Pharmacognosy, MVP's College of Pharmacy, Nashik, Maharashtra, India

Abstract

Introduction: In the present study, fruit extracts of *Momordica cochinchinensis (Cucurbitaceae)* and *Momordica balsamina (Cucurbitaceae)* were investigated for analgesic activity by Eddy's hot plate and Tail immersion method. **Materials and Methods:** The extracts were prepared successively using powdered material with petroleum ether, ethanol, and water, and concentrated under vacuum and were evaluated for analgesic activity at three dose level (100, 200, and 400 mg/kg). **Results and Discussion:** In Eddy's hot plate method, oral administration of petroleum ether extracts of both the plants at the dose of 200 mg/kg ($P < 0.01$) and 400 mg/kg ($P < 0.001$) significantly reduced the thermal stimulation. Analgesic activity of petroleum ether extracts of both plants at the dose of 400 mg/kg after 90 min was comparable to standard drug pentazocine (10 mg/kg). In tail immersion method, petroleum ether extract at the dose of 100 mg/kg ($P < 0.05$), 200 mg/kg, and 400 mg/kg ($P < 0.01$) and alcoholic extract at the dose of 200 mg/kg and 400 mg/kg ($P < 0.05$) of both plant material has shown significant analgesic activity and was comparable to standard drug pentazocine (10 mg/kg) after 90 min. **Conclusion:** It is concluded that petroleum ether extracts of both plant material have central analgesic effects.

Key words: Analgesic activity, Eddy's hot plate, *Momordica balsamina*, *Momordica cochinchinensis*, phytosterols, tail immersion

INTRODUCTION

Momordica is a genus of about 60 species of annual or perennial climbers herbaceous or rarely small shrubs belonging to the family Cucurbitaceae, natives of tropical and subtropical Africa, Asia and Australia.[1,2]

Momordica cochinchinensis (Gac) is a Southeast Asian fruit found throughout the region from Southern China to Northeastern Australia, mostly Vietnam and throughout India. It grows on dioecious vines and is usually collected from fence climbers or wild plants. The vines can be commonly seen growing on lattices at the entrances to rural homes or in gardens. It bears fruits annually and is found in local markets. The fruit becomes a dark orange color on ripening, and is typically round or oblong, maturing to a size of about 13 cm in length and 10 cm in diameter. The exterior skin is covered in small spines, while dark red interior consists of clusters of fleshy pulp and seeds.[3] Gac fruit, *M. cochinchinensis* Spreng, is one of the special fruits containing extraordinarily high levels of carotenoids, especially β-carotene (>16 mg/100 g), and lycopene (>50 mg/100 g), mainly in the red aril.[4] Conventionally, Gac has been used as both food and medicine and promotes healthy vision by relief of dry eyes. It also possesses antioxidant, antimicrobial, and antidiabetic properties. The seeds are considered to be good for cough and pains in the chest.[5-7]

Momordica balsamina Family: Cucurbitaceae is climber with bright green leaves bears striking orange to red spindle-shaped ripe fruit. Shurb is fairly common and widespread in Malaya, Australia, West Asia, Africa, America, and India (Sind, Gujarat, and Deccan). Conventionally used as a purgative agent, purification of blood and dissipate

Address for correspondence:
Mohan R. Agrawal, Department of Pharmacognosy, Sitabai Thite College of Pharmacy, Pune, Maharashtra, India. Tel.: +91-2138-222680/9860947093.
E-mail: agrawalmohan19@gmail.com

Received: 15-05-2018
Revised: 10-08-2018
Accepted: 12-10-2018

Back to Academic Year 2018-2019

Ameliorative Effects Of Artemisia Pallens In A Murine Model of Ovalbumin-Induced Allergic Asthma Via Modulation of Biochemical perturbations

An official website of the United States government
Here's how you know

FULL TEXT LINKS
ELSEVIER FULL-TEXT ARTICLE

Biomed Pharmacother. 2017 Oct;94:880-889. doi: 10.1016/j.biopha.2017.08.017. Epub 2017 Aug 16.

Ameliorative effects of Artemisia pallens in a murine model of ovalbumin–induced allergic asthma via modulation of biochemical perturbations

Anwesha A Mukherjee [1], Amit D Kandhare [1], Supada R Rojatkar [2], Subhash L Bodhankar [3]

Affiliations

PMID: 28810518 DOI: 10.1016/j.biopha.2017.08.017

Abstract

Introduction: Asthma is a chronic, heterogeneous airway disorder characterized by airway inflammatory and remodeling. Artemisia pallens has been reported to possess antioxidant, anti-inflammatory and Anti-allergic potential.

Objective: To evaluate the anti-asthmatic effects of methanolic extract of Artemisia pallens (APME) against ovalbumin (OVA)-induced airway hyperresponsiveness (AHR) in rats.

Materials and method: AHR was induced in male Sprague-Dawley rats (180-200g) by intraperitoneal (i.p.) injection of OVA and boosted with an identical OVA solution (s.c.) on day 7. Rats were either treated orally with vehicle (10mg/kg), montelukast (10mg/kg) or APME (100, 200 and 400mg/kg) for next 28days. At the end treatments, various biochemical, molecular (RT-PCR and ELISA analysis) and histological parameters were evaluated.

Results: APME (200 and 400mg/kg) significantly attenuated ($p<0.05$) OVA-induced alteration in lung functions measured by Whole-body plethysmography. Increased Bronchoalveolar Lavage (BAL) fluid differential cell count, as well as total protein and albumin in BAL fluid and lungs, was significantly decreased ($p<0.05$) by APME. It also significantly attenuated ($p<0.05$) elevated lung oxido-nitrosative stress, myeloperoxidase, and serum IgE levels. OVA-induced down-regulation in lung Nrf2 and upregulation in TNF-α, IL-1β, IL-4, IL-6, TGF-β mRNA expression was significantly attenuated ($p<0.05$) by APME (200 and 400mg/kg) treatment. Histopathological analysis of lung tissue showed that APME treatment reduced OVA-induced inflammatory influx and fibrosis.

Conclusion: Artemisia pallens simultaneously orchestrate plethora of mechanisms viz. modulations of IgE, TGF-β, TNF-α, IL's and Nrf-2 levels to exhibit its anti-asthmatic potential in OVA-induced AHR in rats.

Keywords: Airway hyperresponsiveness; Artemisia pallens; Asthma; IL's; IgE; Nrf-2; Ovalbumin; TGF-β; TNF-α.

Copyright © 2017 Elsevier Masson SAS. All rights reserved.

PubMed Disclaimer

Related information

MedGen
PubChem Compound (MeSH Keyword)

LinkOut – more resources

Full Text Sources
Elsevier Science

Other Literature Sources
scite Smart Citations

Back to Academic Year 2018-2019

Evaluation of Health-Related Quality Of Life In Hemolytic U
With Eculizumab: A Systematic Evaluation On Basis Of EMPRO

As a library, NLM provides access to scientific literature. Inclusion in an NLM database does not imply endorsement of, or agreement with, the contents by NLM or the National Institutes of Health.
Learn more: PMC Disclaimer | PMC Copyright Notice

Renal Failure
Taylor & Francis
Taylor & Francis Group

Ren Fail. 2018; 40(1): 107–118. PMCID: PMC6014301
Published online 2018 Jan 24. doi: 10.1080/0886022X.2018.1427110 PMID: 29363392

Evaluation of health-related quality of life in hemolytic uraemic syndrome patients treated with eculizumab: a systematic evaluation on basis of EMPRO

Anwesha A. Mukherjee, Amit D. Kandhare, and Subhash L. Bodhankar

Abstract

Background: Hemolytic uraemic syndrome (HUS) is progressive renal failure disease and determination of their quality of life (QoL) on the basis of patient-reported outcomes (PROs) are becoming increasingly important in the economic evaluations for its treatment with eculizumab (ECU).

Aim: To perform the systematic evaluation of QoL in HUS patients treated with ECU on the basis of Evaluating Measures of Patient Reported Outcomes (EMPRO) tool.

Materials and methods: A systematic review was conducted in PubMed, EMBASE, the Cochrane Library, CINAHL and Google Scholar till September 2016 by two independent researchers. Each identified instrument was evaluated for its quality of performance by using the EMPRO tool for its overall score and seven attribute specific scores (range 0–100, worst to best).

Results: Five different PROs instruments were identified from 10 articles ($n = 112$) which showed eculizumab significantly improves health-related quality of life (HRQOL) in atypical HUS (aHUS) patients. Amongst five instruments viz. EuroQol five dimensions questionnaire (EQ-5 D) Functional Assessment of Chronic Illness Therapy-Fatigue (FACIT-F), Headache Impact Test-6 (HIT-6), 36-Item Short Form Health Survey (SF-36) and Visual Analogue Scale (VAS), the overal EMPRO score was higher for VAS (73.83) and EQ-5 D (73.81). Whereas, FACIT-F and HIT- 6 were just able to meet the minimal threshold of EMPRO scoring (50.24 and 59.09, respectively).

Conclusions: Evidence from present investigation support that eculizumab significantly proves HRQoL in patients with aHUS furthermore, EQ-5 D and VAS instrument should be ommended for assessing HRQoL in them. However, selection of PRO instrument for dete

Feedback
Back to Top

Back to Academic Year 2018-2019

Curcumin Ameliorates Vitamin A Deficiency-Induced Urolithiasis In Neonatal Rats Via Inhibition Of KIM-1/NGAL, Nrf2, And Inos Pathways

Recruit researchers Join for free Login



Home > Nutritional Medicine > Spices > Medicine > Food Science > Nutrition and Dietetics > Curcumin

Article

Curcumin ameliorates vitamin A deficiency-induced urolithiasis in neonatal rats via inhibition of KIM-1/NGAL, Nrf2, and iNOs pathways

January 2018 · LATIN AMERICAN JOURNAL OF PHARMACY 37(12):2502-2511

Authors:

A.A. Mukherjee · J. Zhang · Amit Kandhare, INDUS BIOTECH · Subhash Laxmanrao Bodhankar, Bharati Vidyapeeth's College of Pharmacy

Request full-text · Download citation · Copy link

To read the full-text of this research, you can request a copy directly from the authors.

Citations (6)

Abstract

The aim of present study was to investigate the effect of curcumin on vitamin A deficiency (VAD)-induced urolithiasis in neonatal rats. The weanling male Wistar rats was administered with vitamin A-deficient diet for 12 weeks and they were concomitantly treated with curcumin (20, 40, and 80 mg/kg). VAD resulted in significant alteration in urinary BUN, LDH, uric acid, creatinine, sodium, calcium, albumin, citrate, oxalate and glycosaminoglycans levels in urolithiasis control rats whereas curcumin treatment (40 and 80 mg/kg) significantly ($p < 0.001$) attenuated these alterations. Curcumin significantly ($p < 0.001$) inhibited VAD-induced elevated levels of oxido-nitrosative stress, up-regulated KIM-1, NGAL, iNOs and down-regulated Nrf2 mRNA expression. Transmission electron microscopical analysis revealed that curcumin attenuates ultrastructural alteration in the kidney induced by VAD. In conclusion, curcumin exerts it nephroprotective effect against VAD-induced urolithiasis in neonatal rats via modulation of renal function test, oxido-nitrosative stress and mRNA expressions of KIM-1, NGAL, Nrf2, and iNOs. © 2018, Colegio de Farmaceuticos de la Provincia de Buenos Aires. All rights reserved.

Discover the world's research
- 25+ million members
- 100+ million publication pages
- 2.3+ citatic Join for free

Sponsored videos

No full-text available

Back to Academic Year 2018-2019

Neuroprotective Effect of Naringin, A Flavone Glycoside In
Possible Role Of PPAR-Γ, Bax/Bcl-2, And Caspase-3

ScienceDirect

Food and Chemical Toxicology
Volume 121, November 2018, Pages 95-108

Neuroprotective effect of naringin, a flavone glycoside in quinolinic acid–induced neurotoxicity: Possible role of PPAR-γ, Bax/Bcl-2, and caspase-3

Jian Cui [a], Gang Wang [a], Amit D. Kandhare [b], Anwesha A. Mukherjee-Kandhare [b], Subhash L. Bodhankar [b]

Show more

Share Cite

https://doi.org/10.1016/j.fct.2018.08.028
Get rights and content

Highlights

- Neurotoxicity was induced in Wistar rats by intrastriatal injection of quinolinic acid.
- Naringin decreased oxido-nitrosative stress, TNF-α, IL's and NF-kB mRNA levels.
- Naringin attenuated apoptosis (Bax-Bcl-2 and Caspase-3 mRNA) level.
- It increased mitochondrial complex (I-IV) activity and PPARγ mRNA expression.
- Naringin exerts its neuroprotective effect against QA-induced neurotoxicity.

Abstract

Background

Huntington's disease (HD) is a complex multifactorial neurodegenerative disorder. Naringin, a flavanone glycoside exhibits potent anti-inflammatory and antiapoptotic effect.

Aim

To evaluate the effect of naringin in quinolinic acid (QA)-induced neurotoxicity in laboratory rats.

Methods

Neurotoxicity was induced in male Wistar rats by single intrastriatal injection of QA (300 nmol/4 μl saline) in striatum except non-treated. Rats were administered orally with either vehicle (distilled water (10 mL/kg) or naringin (20, 40 and 80 mg/kg) or pioglitazone (40 mg/kg, p.o.) or its combination for 28 days.

Low Molecular Weight Galactomannans-Based Standardized Fenugreek Seed Extract Ameliorates High Fat Diet Induced Obesity In Mice Via Modulation Of Fasn, IL-6, Leptin, And TRIP-Br_2

Back to Academic Year 2018-2019

RSC Advances

PAPER

View Article Online
View Journal | View Issue

Open Access Article. Published on 18 September 2018. Downloaded on 3/20/2024 9:00:19 AM.
This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence.

Check for updates

Cite this: *RSC Adv.*, 2018, **8**, 32401

Low molecular weight galactomannans-based standardized fenugreek seed extract ameliorates high-fat diet-induced obesity in mice *via* modulation of FASn, IL-6, leptin, and TRIP-Br2†

Amit D. Kandhare, *[a] Debasish Bandyopadhyay[b] and Prasad A. Thakurdesai[a]

Background: Obesity is a complex, chronic metabolic disorder and its prevalence is increasing throughout most of the world. Low molecular weight galactomannans-based standardized fenugreek seed extract (LMWGAL-TF) has previously shown anti-diabetic and anti-hyperlipidemic potential. *Aim*: To evaluate the efficacy and mechanism of action of LMWGAL-TF in treating high fat diet (HFD)-induced obesity and hyperlipidemia in mice. *Materials and methods*: Male C57BL/6 mice were fed the HFD for 12 weeks and were co-administered with LMWGAL-TF (10, 30 and 100 mg kg^{-1}, p.o.). Variables measured were behavioral, biochemical, molecular and histopathological. In a separate *in vitro* experiment, copper-ascorbate (Cu-As)-induced mitochondrial oxidative damage was evaluated. *Results*: The HFD-induced increase ($p < 0.001$) in body weight, fat mass, lean mass, adipose tissue (brown, mesenteric, epididymal and retroperitoneal) and liver weight was significantly attenuated ($p < 0.001$) by LMWGAL-TF (30 and 100 mg kg^{-1}). The HFD-induced elevated levels of serum lipid, interleukins (ILs)-6 and leptin were significantly decreased ($p < 0.001$) by LMWGAL-TF (30 and 100 mg kg^{-1}). Elevated fatty acid synthase (FASn), IL-6, leptin and transcriptional regulator interacting with the PHD-bromodomain 2 (TRIP-Br2) mRNA expression in brown adipose tissue (BAT), liver, and epididymal fat were significantly down-regulated ($p < 0.001$) by LMWGAL-TF (30 and 100 mg kg^{-1}). Additionally, HFD-induced histological alterations in skeletal muscle, liver, white adipose tissue (WAT) and BAT were also reduced by LMWGAL-TF. Furthermore, the Cu-As-induced alteration in mitochondria oxidative stress (lipid peroxidation, protein carbonylation, glutathione, glutathione reductase, glutathione peroxidase, isocitrate dehydrogenase and α-ketoglutarate dehydrogenase) in skeletal muscle and BAT was significantly ($p < 0.001$) ameliorated by LMWGAL-TF (2, 4 and 6 mg mL^{-1}) treatment. It also reduced the Cu-As-induced mitochondrial swelling. *Conclusion*: LMWGAL-TF showed its beneficial effect in reducing HFD-induced obesity *via* down-regulation of FASn, IL-6, leptin, and TRIP-Br2 in mice.

Received 17th June 2018
Accepted 4th September 2018

DOI: 10.1039/c8ra05204b

rsc.li/rsc-advances

1 Introduction

Type 2 diabetes mellitus (T2DM) is a complex, chronic metabolic disorder across the world. An array of conditions such as imbalanced eating habits, high-fat diet (HFD), excessive caloric intake, reduced physical activity, sedentary lifestyles and neuroendocrine factors integrated with T2DM leads to increased obesity, which is now becoming one of the most severe threats.[1] Obesity in the population has increased significantly over the past decades and has now become a major factor responsible for morbidity and mortality worldwide.[2] In 2016, there were more than 1.9 billion adults that were overweight, and of these, over 650 million were obese, whereas 41 million children were overweight or obese.[3] It has been reported that obesity is also associated with other diseases such as cardiovascular disease, dyslipidemia, cerebrovascular disease, hypertension, non-alcoholic fatty liver disease and arthritis, which diminish the quality of life.[4]

Several mechanisms have been proposed to explain T2DM-induced obesity. Researchers have established the correlation between oxido-inflammatory stress and the development of metabolic disorders associated with obesity.[5,6] White adipose tissue (WAT) plays a vital role in this oxido-inflammatory-induced obesity and dyslipidemia. Adipose tissue is well known for releasing and controlling adipokines (leptin and adiponectin) and cytokines (tumor necrosis factor-α (TNF-α) and interleukin-6 (IL-6)).[7] It has been well documented that increased oxidative stress in adipocytes causes an imbalance between the leptin and adiponectin, leading to profound metabolic alterations that

[a]*Department of Scientific Affairs, Indus Biotech Private Limited, 1, Rahul Residency, Off Salunke Vihar Road, Kondhwa, Pune 411048, Maharashtra, India. E-mail: amit.kandhare@indusbiotech.com; Tel: +91-9226164041*
[b]*Oxidative Stress and Free Radical Biology Laboratory, Department of Physiology, University of Calcutta, University College of Science and Technology, Kolkata 700 009, India*

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c8ra05204b

Anti-Epileptic Effect Of Morin Against Experimental Pentylenetetrazol-Induced Seizures Via Modulating Brain Monoamines And Oxidative Stress

Back to Academic Year 2018-2019

[Downloaded free from http://www.apjtb.org on Thursday, September 6, 2018, IP: 10.232.74.23]

Asian Pacific Journal of Tropical Biomedicine

Journal homepage: www.apjtb.org

doi: 10.4103/2221-1691.237078

©2018 by the Asian Pacific Journal of Tropical Biomedicine.

Anti-epileptic effect of morin against experimental pentylenetetrazol-induced seizures via modulating brain monoamines and oxidative stress

Amit D. Kandhare✉, Anwesha A. Mukherjee, Subhash L. Bodhankar

Department of Pharmacology, Poona College of Pharmacy, Bharati Vidyapeeth Deemed University, Erandwane, Paud Road, Pune-411 038, India

ARTICLE INFO

Article history:
Received 11 April 2018
Revision 5 May 2018
Accepted 19 June 2018
Available online 19 July 2018

Keywords:
Brain GABA
Epilepsy
Morin
Nitric oxide
Pentylenetetrazol
Oxidative stress
Xanthine oxidase

ABSTRACT

Objective: To evaluate the protective effect of morin against pentylenetetrazol (PTZ)-induced tonic-clonic convulsions in mice. **Methods:** Swiss albino mice (18-22 g) was used to induce convulsions by intraperitoneal (i.p.) administration of PTZ (90 mg/kg). Mice were either pretreated with morin (10, 20 and 40 mg/kg) or vehicle (distilled water, 10 mg/kg) 45 min before PTZ administration. Various behavioral and biochemical parameters were assessed. **Results:** PTZ administration resulted in significant production (P<0.001) of tonic-clonic conclusion and mortality in mice. PTZ-induced increase in the duration of convulsion, onset of convulsion and mortality was inhibited significantly by morin (20 and 40 mg/kg) administration. The PTZ-induced decrease in brain GABA, dopamine and Na^+K^+ATPase levels and increase in xanthine oxidase activity were inhibited significantly by morin (20 and 40 mg/kg) treatment. The increased levels of malondialdehyde and nitric oxide level were significantly decreased by morin (20 and 40 mg/kg) treatment. Also, reduced levels of superoxide dismutase and glutathione were increased significantly by morin treatment. **Conclusions:** Results of the present study indicate that morin showed its anti-convulsant effect via modulating the levels of brain GABA, Na^+K^+ATPase, and oxido-nitrosative stress. Thus, morin can be a potential candidate for further clinical evaluations as an anti-epileptic agent.

1. Introduction

Epilepsy is a chronic disease of neurobiological system, affecting approximately 50/100 000 individual while 100/100 000 individual in developed and developing countries, respectively[1]. It has been reported that epilepsy prevalence in India is about 1/18th of the world population which is about 5.5-7.9/1 000 individual[2]. The children below 7 years-of-age and person of above 55 years have the higher epilepsy incidence[2].

Research carried out over past decades indicated that there are various mechanisms responsible for triggering neuronal injury[3-5]. This mechanism includes an imbalance between the neurotransmitter system such as glutamate-GABA (gamma-Aminobutyric acid) or impairment in voltage- and receptor-gated ion channels which plays a decisive role in the neuronal cell death. This resulted in various disorders of neurobiological system such as epilepsy, Alzheimer's disease, parkinsonism and ischemia[6,7]. The increase in the consequence of epileptic seizures led to the development of many antiepileptic drugs for ameliorating the neuronal damage[8].

✉First and corresponding author: Amit D. Kandhare, Department of Pharmacology, Poona College of Pharmacy, Bharati Vidyapeeth Deemed University, Erandwane, Paud Road, Pune-411 038, India.
E-mail: amitkandhare25@gmail.com

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

©2018 Asian Pacific Journal of Tropical Biomedicine Produced by Wolters Kluwer-Medknow

How to cite this article: Kandhare AD, Mukherjee AA, Bodhankar SL. Anti-epileptic effect of morin against experimental pentylenetetrazol-induced seizures via modulating brain monoamines and oxidative stress. Asian Pac J Trop Biomed 2018; 8(7): 352-359.

Back to Academic Year 2018-2019

nulation Rich In Standardized Fenugreek Seed Extract In Type 2
nts With Insulin Therapy: An Efficacy And Safety Study

[Downloaded free from http://www.apjtb.org on Thursday, September 27, 2018, IP: 10.232.74.23]

Asian Pacific Journal of Tropical Biomedicine
Journal homepage: www.apjtb.org

doi: 10.4103/2221-1691.242288

©2018 by the Asian Pacific Journal of Tropical Biomedicine.

Add-on therapy of herbal formulation rich in standardized fenugreek seed extract in type 2 diabetes mellitus patients with insulin therapy: An efficacy and safety study

Amit Kandhare[1], Uday Phadke[2], Abhay Mane[3], Prasad Thakurdesai[1], Sunil Bhaskaran[1]

[1]Department of Scientific Affairs, Indus Biotech Private Limited, 1, Rahul Residency, Off Saiunke Vihar Road, Kondhwa, Pune - 411048, India
[2]Ruby Hall Clinic, 40, Sassoon Road, Pune 411001, India
[3]Tulip, Opp. Camp Education Society's High School, Camp, Pune-411001, India

ARTICLE INFO

Article history:
Received 2 July 2018
Revision 1 August 2018
Accepted 5 September 2018
Available online 27 September 2018

Keywords:
Add-on therapy
Dietary supplement
Glycaemic control
HbA1c
Standardized fenugreek seed extract
Type 2 diabetes

ABSTRACT

Objective: To assess the safety and efficacy of herbal formulation rich in standardized fenugreek seed extract (IND-2) add-on therapy in type 2 diabetes mellitus (T2DM) patients who were on insulin treatment in prospective, single arm, open-label, uncontrolled, multicentre trial. **Methods:** T2DM patients (n=30) with aged 18-80 years who were stabilized on insulin treatment with fasting blood sugar (FBS) level between 100-140 mg/dL received IND-2 capsules (700 mg, thrice a day) for 16 weeks. The primary endpoints were an assessment of FBS at week 2, 4, 6, 8, 12 and 16. Secondary end-points include post-prandial blood sugar level, glycosylated Hb (HbA1c), reduction in the dose of insulin and number of hypoglycemic attacks, and improvement in lipid profile at various weeks. Safety and adverse events (AEs) were also assessed during the study. **Results:** Study was completed in twenty T2DM patients, and there was no significant reduction in FBS and post-prandial blood sugar level after add-on therapy of IND-2. However, add-on therapy of IND-2 significantly reduced (P<0.01) the HbA1c values, requirements of insulin and hypoglycemic events as compared with baseline. Total cholesterol, high-density lipoproteins-cholesterol, and low-density lipoprotein-cholesterol levels were significantly increased (P<0.01) after IND-2 add-on therapy. Body weight and safety outcomes did not differ significantly in IND-2 add-on therapy group at week 16. Additionally, add-on therapy of IND-2 did not produce any serious adverse events. **Conclusions:** The results of present investigation suggest that add-on therapy of IND-2 with insulin in T2DM patients improves glycaemic control through a decrease in levels of HbA1c and number of insulin doses needed per day without an increase in body weight and risk of hypoglycemia. Thus, IND-2 may provide a safe and well-tolerated add-on therapy option for the management of T2DM.

1. Introduction

Type 2 diabetes mellitus (T2DM) is a complex, chronic progressive disease which is most common amongst the various forms of diabetes. It is manifested by hyperglycemia, disturbance in the metabolism of carbohydrate, fat, and protein which could be the result from deficiency secretion of insulin or its actions[1]. It is a long-term metabolic disorder which is associated with multiple comorbidities as well as microvascular and macrovascular

*First and corresponding author: Amit Kandhare, Department of Scientific Affairs, Indus Biotech Private Limited, Rahul Residency, Off Saiunke Vihar Road, Kondhwa, Pune - 411 048, Maharashtra, India.
Tel: +020 2685 1239; 020 2685 2139
E-mail: amit.kandhare@indusbiotech.com
Foundation project: This work was funded by Indus Biotech Pvt. Ltd., Pune.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.
For reprints contact: reprints@medknow.com
©2018 Asian Pacific Journal of Tropical Biomedicine Produced by Wolters Kluwer-Medknow
How to cite this article: Kandhare A, Phadke U, Mane A, Thakurdesai P, Bhaskaran S. Add-on therapy of herbal formulation rich in standardized fenugreek seed extract in type 2 diabetes mellitus patients with insulin therapy: An efficacy and safety study. Asian Pac J Trop Biomed 2018; 8(9): 446-455.

Efficacy And Safety Of Herbal Formulation Rich In Standardized Fenugreek Seed Extract As Add-On [illegible] With Type 2 Diabetes Mellitus On Sulfonylurea Therapy: A 12-Week, Double-Blind, Placebo-Controlled, Multi-Center Study

Back to Academic Year 2018-2019

Pharmacogn. Mag.
A multifaceted peer reviewed journal in the field of Pharmacognosy and Natural Products
www.phcog.com | www.phcog.net

ORIGINAL ARTICLE

Efficacy and Safety of Herbal Formulation Rich in Standardized Fenugreek Seed Extract as Add-on Supplementation in Patients with Type 2 Diabetes Mellitus on Sulfonylurea Therapy: A 12-week, Randomized, Double-blind, Placebo-controlled, Multi-center Study

Amit D. Kandhare[1], Nadeem Rais[2], Nivedita Moulick[3], Alaka Deshpande[4], Prasad Thakurdesai[1], Sunil Bhaskaran[1]

[1]Department of Scientific Affairs, Indus Biotech Private Limited, Pune, [2]Endocrine and Metabolic Section, Chowpatty Medical Centre, Babulnath Road, [3]Department of Medicine, Lokmanya Tilak Memorial Medical College and Lokmanya Tilak General Hospital, Sion, [4]Department of Medicine, Grant Medical College and Sir JJ Group of Hospitals, Byculla, Mumbai, Maharashtra, India

Submitted: 16-05-2018 Revised: 21-05-2018 Published: 10-09-2018

ABSTRACT

Background: Type 2 diabetes mellitus (T2DM) is a chronic, complex, and progressive illness that often needs combination therapy for better glycemic control. IDM1, an herbal formulation which is rich in standardized fenugreek seed extract. **Aim:** The aim of this study is to evaluate the efficacy and safety of add-on therapy of IDM1 in T2DM patients inadequately controlled on sulfonylurea monotherapy. **Materials and Methods:** In this 12-week, randomized, double-blind, placebo-controlled, multi-centric study, T2DM patients which inadequate glycemic control with background stable dose medication of sulfonylurea was screened (n = 120). The patients were randomized 1:1 to add-on therapy of IDM1 and placebo, 700 mg three times daily for 12 weeks. **Results:** A total of 119 patients were randomized and included in the efficacy analysis (IDM1, n = 60; placebo, n = 59). At week 12, adjusted fasting plasma glucose (FPG) (20 mg%), postprandial plasma glucose (PPPG) (26 mg%), and glycated hemoglobin (HbA1c) (0.9 mg%) was reduced significantly ($P < 0.05$) from baseline as compared to placebo group (FPG: 7 mg%; PPPG: 4 mg% and HbA1c: 0.4 mg%). These beneficial effects were seen as early as 1 month after consumption of IDM1 and continued until at least 15 days after withdrawal of IDM1. Hypoglycemic events were mostly mild, and none required emergency treatment. There were no major changes in body weight, hematology, and biochemistry at week 12 as compared to baseline. Overall AEs rates were similar in both groups. **Conclusions:** IDM1 is a safe, effective, and well-tolerated add-on oral medication therapy that supports healthy blood sugar levels and glycosylated hemoglobin levels in T2DM patients inadequately controlled with a sulfonylurea.

Key words: Add-on therapy, glycemic control, glycated hemoglobin, standardized fenugreek seed extract, sulfonylureas, type 2 diabetes

SUMMARY

- IDM1, an herbal formulation which is rich in standardized fenugreek seed extract
- In this 12-week, randomized, double-blind, placebo-controlled, multi-centric study, T2DM patients which inadequate glycemic control with a stable dose of sulfonylurea was received add-on therapy of IDM1 (700 mg, three times daily) for 12 weeks
- At week 12, add-on therapy of IDM1 showed significant ($P < 0.05$) reduction in adjusted FPG (20 mg%), PPPG (26 mg%), and HbA1c (0.9 mg%) from baseline as compared to placebo group
- Thus, IDM1 is a safe, effective, and well-tolerated add-on oral medication therapy that supports healthy blood sugar levels and glycosylated hemoglobin levels in T2DM patients inadequately controlled with a sulfonylurea.



Abbreviation used: AEs: Adverse events; ALP: Alkaline phosphatase; ALT: Alanine transaminase; AST: Aspartate transaminase; BMI: Body mass index; BUN: Blood urea nitrogen; FPG: Fasting plasma glucose; GGT: Gamma glutamyl transferase; HbA1c: Glycated hemoglobin; HDL: High-density lipoproteins; LDH: Lactate dehydrogenase; LDL: Low-density lipoproteins.

Correspondence:
Dr. Amit D. Kandhare,
Department of Scientific Affairs, Indus Biotech Private Limited, 1, Rahul Residency, Off Salunke Vihar Road, Kondhwa, Pune - 411 048, Maharashtra, India.
E-mail: amit.kandhare@indusbiotech.com

DOI: 10.4103/pm.pm_260_18

Access this article online
Website: www.phcog.com
Quick Response Code:

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Cite this article as: Kandhare AD, Rais N, Moulick N, Deshpande A, Thakurdesai P, Bhaskaran S. Efficacy and safety of herbal formulation rich in standardized fenugreek seed extract as add-on supplementation in patients with type 2 diabetes mellitus on sulfonylurea therapy: A 12-week, randomized, double-blind, placebo-controlled, multi-center study. Phcog Mag 2018;14:S393-402.

Back to Academic Year 2018-2019

ed Quality Of Life In Hemolytic Uraemic Syndrome Patients Treated With
zumab: A Systematic Evaluation On Basis Of EMPRO

As a library, NLM provides access to scientific literature. Inclusion in an NLM database does not imply endorsement of, or agreement with, the contents by NLM or the National Institutes of Health.
Learn more: PMC Disclaimer | PMC Copyright Notice

Renal Failure

Ren Fail. 2018; 40(1): 107–118.
Published online 2018 Jan 24. doi: 10.1080/0886022X.2018.1427110

PMCID: PMC6014301
PMID: 29363392

Evaluation of health-related quality of life in hemolytic uraemic syndrome patients treated with eculizumab: a systematic evaluation on basis of EMPRO

Anwesha A. Mukherjee, Amit D. Kandhare, and Subhash L. Bodhankar

Abstract

Background: Hemolytic uraemic syndrome (HUS) is progressive renal failure disease and determination of their quality of life (QoL) on the basis of patient-reported outcomes (PROs) are becoming increasingly important in the economic evaluations for its treatment with eculizumab (ECU).

Aim: To perform the systematic evaluation of QoL in HUS patients treated with ECU on the basis of Evaluating Measures of Patient Reported Outcomes (EMPRO) tool.

Materials and methods: A systematic review was conducted in PubMed, EMBASE, the Cochrane Library, CINAHL and Google Scholar till September 2016 by two independent researchers. Each identified instrument was evaluated for its quality of performance by using the EMPRO tool for its overall score and seven attribute specific scores (range 0–100, worst to best).

Results: Five different PROs instruments were identified from 10 articles ($n = 112$) which showed eculizumab significantly improves health-related quality of life (HRQOL) in atypical HUS (aHUS) patients. Amongst five instruments viz. EuroQol five dimensions questionnaire (EQ-5 D) Functional Assessment of Chronic Illness Therapy-Fatigue (FACIT-F), Headache Impact Test-6 (HIT-6), 36-Item Short Form Health Survey (SF-36) and Visual Analogue Scale (VAS), the overal EMPRO score was higher for VAS (73.83) and EQ-5 D (73.81). Whereas, FACIT-F and HIT- 6 were just able to meet the minimal threshold of EMPRO scoring (50.24 and 59.09, respectively)

Conclusions: Evidence from present investigation support that eculizumab significantly proves HRQoL in patients with aHUS furthermore, EQ-5 D and VAS instrument should be ommended for assessing HRQoL in them. However, selection of PRO instrument for dete

Feedback
Back to Top

Hesperidin, A Plant Flavonoid Accelerated The Cutaneous Wound Healing In Streptozotocin-Induced Diabetic Rats: Role Of Tgf-B/Smads And Ang-1/Tie2

Back to Academic Year 2018-2019

EXCLI Journal 2018;17:399-419 – ISSN 1611-2156
Received: January 07, 2018, accepted: March 28, 2018, published: May 04, 2018

Supplementary data to:

HESPERIDIN, A PLANT FLAVONOID ACCELERATED THE CUTANEOUS WOUND HEALING IN STREPTOZOTOCIN-INDUCED DIABETIC RATS: ROLE OF TGF-B/SMADS AND ANG-1/TIE2 SIGNALING PATHWAYS

Wenbin Li[1], Amit D. Kandhare[2,3]*, Anwesha A. Mukherjee[2], Subhash L. Bodhankar[2]

[1] Department of Dermatology, Shaanxi Traditional Chinese Medicine Hospital, Xi'an, Shaanxi, 710003, China
[2] Department of Pharmacology, Poona College of Pharmacy, Bharati Vidyapeeth Deemed University, Erandwane, Paud Road, Pune-411 038, India
[3] Jalan Universiti Bandar Barat, 31900, Kampar, Perak, Malaysia

* Corresponding author: Dr. Amit D. Kandhare, Department of Pharmacology, Poona College of Pharmacy, Bharati Vidyapeeth Deemed University, Erandwane, Paud Road, Pune-411 038, India, E-mail: amitkandhare25@gmail.com

http://dx.doi.org/10.17179/excli2018-1036

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0/).

Effect of hesperidin and insulin treatment on body weight (in gm) of diabetic rats

Rat No.	Normal	NWC	DWC	I (10)	H (25)	H (50)	H (100)
1	188.2	205.0	159.5	196.9	166.0	192.0	188.9
2	214.8	178.0	162.0	206.2	171.0	171.0	206.1
3	202.0	180.0	168.0	202.4	171.0	196.0	200.0
4	199.0	185.0	162.0	200.0	179.5	201.0	200.0
5	187.6	184.0	164.0	210.1	177.0	196.0	189.1
6	183.4	181.0	175.5	185.4	175.5	200.0	189.9

NWC: Normal wound control group; DWC: Diabetic wound control group; I (10): Insulin (10 IU/kg, s.c.) treated group; H (25): Hesperidin (25 mg/kg, p.o.) treated group; H (50): Hesperidin (50 mg/kg, p.o.) treated group; H (100): Hesperidin (100 mg/kg, p.o.) treated group

Back to Academic Year 2018-2019

Anti-Allergic Potential Of Fisetin In A Murine Model Of Ova-Induced Allergic Rhinitis Via Inhibition Of Gata-3 And Th2 Cytokines Zhao L

D:\Biomedica Vol. 34, Issue 2, Apr. – Jun., 2018\Bio-8.Doc Fig. 2-4 Color P. 88 – 101 (KC) IV

ORIGINAL ARTICLE

ANTI-ALLERGIC POTENTIAL OF FISETIN IN A MURINE MODEL OF OVA-INDUCED ALLERGIC RHINITIS VIA INHIBITION OF GATA-3 AND TH2 CYTOKINES

ZHAO L.,[1] KANDHARE A.,[2] MUKHERJEE A.[2] AND SUBHASH BODHANKAR S.[2]
[1]*Department of Otolaryngology, Head and Neck Surgery, The First Affiliated Hospital of Xi'an Medical University Feng Hao West Road, Lianhu District, Xi'an, Shaanxi-710077, China*
[2]*Department of Pharmacology, Poona College of Pharmacy, Bharati Vidyapeeth Deemed University Erandwane, Paud Road, Pune-411038, India*

ABSTRACT

Background and Objectives: Allergic rhinitis (AR) is an IgE mediated immune-inflammatory disease, characterized by sneezing, rhinorrhea, lacrimation, etc. Fisetin is a plant flavonoid, reported to have anti-allergic potential via inhibition of Th2 response. Objective of this study was to evaluate the anti-allergic potential of fisetin against ovalbumin (OVA)-induced experimental allergic rhinitis in BALB/c mice.

Methodology: Mice were sensitization by OVA (500 µl, i.p.) on days 1, 3, 5, 7, 9, 11, 13 and treated with fisetin (10, 20 and 40 mg/kg, p.o.) for 21 days, followed by OVA (5 µl per nostril) challenged on the 21st day.

Results: Administration of fisetin (20 and 40 mg/kg) significantly attenuated ($p < 0.05$) OVA-induced nasal rubbing, sneezing and discharge, histamine-induced rubbing and sneezing, and hematological parameters as compared to AR control mice. OVA-induced elevated levels of serum histamine, β-hexosaminidase, IgE and IgG1 as well as Th2 cytokines (IL-4, IL-5, IL-13, and IL-17) in nasal lavage fluid was significantly ($p < 0.05$) attenuated by fisetin. It also significantly inhibit ($p < 0.05$) up-regulated mRNA expressions of GATA3, IL-4, IL-5 and IL-13 in spleen tissue. Fisetin administration significantly reduced ($p < 0.05$) histological aberrations induced by OVA in nasal mucosa, spleen, and lungs.

Conclusion: The findings of the present study showed that fisetin exerts its anti-allergic potential via modulation of GATA3 pathway to inhibit the release of Th2 cytokines (IL-4, IL-5, IL-13) and IgE, thus reducing OVA-induced nasal rubbing and sneezing during allergic rhinitis.

Keywords: Allergic rhinitis, Fisetin, GATA3, lgE, Interleukins, Ovalbumin, TNF-α.

INTRODUCTION

Allergic rhinitis (AR) is a chronic immune-inflammatory disorder which is increasing, significantly, worldwide resulting in important social and medical problems.[1] Rhinitis is associated with widespread morbidity, significant treatment costs, impaired work productivity and quality of life. The characteristic features of AR include nasal congestion, sneezing, rhinorrhea, and pruritus of nosea long with eyes. Additionally, it is also associated with various complications including headache, fatigue, sinusitis, sleep disturbance, eustachian tube dysfunction, and cognitive impairment.[2] According to a report by European Community Respiratory Health Survey (ECRHS), the prevalence of AR is 5-22%. While, more than 35% of the European and Australian population suffer from AR, about 26% of Indians are affected with it.[3]

Researchers have well documented that exposure to an array of mediators such as indoor allergens (such as dust mites and stuffed furniture), outdoor allergens (like molds and pollen of grains, grass, trees, weeds, etc.), chemical irritants, tobacco smoke, air pollution, and food cause hypersensitive response which results in Allergic rhinitis.[4]Cumulative data obtained from animal studies have suggested that imbalance in T-helper type 1 (Th1) and T-helper type 2 (Th2) responses cause inflammation and remodelling in nasal mucosa that result in the progress of AR.[5,6] Th2 responses are up-regulated due to elevated production of cytokines including tumour necrosis factor-alpha (TNF-α), interleukins (IL's) (IL-1β, IL-4, IL-5 and IL-6), release of nitric oxide from macrophages, production of reactive oxygen species (ROS), immunoglobulin (IgE), and mast cell.[5,6] These vicious molecules are responsible for synthesis of IgE and its cross-linking with high-affinity IgE receptors (FcεRI) present on mast cell sur-

88 Corresponding Author: Dr. S. L. Bodhankar
Dept. of Pharmacology, Poona College of Pharmacy, Bharati Vidyapeeth Deemed University, Erandwane, Pune-411038, Maharashtra, India. Email: drslbodh@gmail.com

Elucidation of The Molecular Mechanism of Tempol In Pentylenetetrazol-Induced Epilepsy In Mice: Role of Gamma - Aminobutyric Acid, Tumor Necrosis Factor-Alpha, Interleukin-1β And C-Fos

Back to Academic Year 2018-2019

Elucidation of the molecular mechanism of tempol in pentylenetetrazol-induced epilepsy in mice: Role of gamma-aminobutyric acid, tumor necrosis factor-alpha, interleukin-1β and c-Fos

https://doi.org/10.4103/pm.pm_263_18

Journal: Pharmacognosy Magazine, 2018, Nº 59, p. 520

Publisher: EManuscript Technologies

Authors:

1. SubhashL Bodhankar
2. Leguo Zhang
3. Tao Wu
4. Amit Kandhare
5. Anwesha Mukherjee
6. Gang Guo

List of references

1. Fisher, Epilepsia, Epileptic seizures and epilepsy: Definitions proposed by the International League Against Epilepsy (ILAE) and The International Bureau for Epilepsy (IBE), c. 470
2. Saberi, Neurosci Lett, The antiepileptic activity of Vitex agnus castus extract on amygdala kindled seizures in male rats, c. 193
3. Nag, Neurol India, Gender and epilepsy: A Clinician's experience, c. 99
4. World Health Organization. Neurological Disorders: Public Health Challenges. Geneva, Switzerland: World Health Organization; 2006.
5. Amudhan, Ann Indian Acad Neurol, Epilepsy in India I: Epidemiology and public health, c. 263
6. Uttara, Curr Neuropharmacol, Oxidative stress and neurodegenerative diseases: A review of upstream and downstream antioxidant therapeutic options, c. 65
7. Achliya, Indian J Pharmacol, Evaluation of CNS activity of Bramhi Ghrita, c. 33
8. Pitkänen, Epilepsy Res, Efficacy of current antiepileptics to prevent neurodegeneration in epilepsy models, c. 141
9. Costa, Epilepsia, Clinical comparability of the new antiepileptic drugs in refractory partial epilepsy: A systematic review and meta-analysis, c. 1280
10. Gupta, Indian J Physiol Pharmacol, Antiepileptic drug therapy in the twenty first century, c. 8
11. de, Seizure, Side-effects of antiepileptic drugs: The economic burden, c. 184
12. Kandhare, Int J Pharm Biol Arc, Patentability of animal models: India and the globe, c. 1024
13. Fisher, Curr Neurol Neurosci Rep, The new classification of seizures by the international league against epilepsy 2017, c. 48
14. Patil, Orient Pharm Exp Med, Determination of role of GABA and nitric oxide in anticonvulsant activity of Fragaria vesca L.ethanolic extract in chemically induced epilepsy in laboratory animals, c. 255
15. Patki, PLoS One, Tempol treatment reduces anxiety-like behaviors induced by multiple anxiogenic drugs in rats, c. e0117498
16. Shahidi, Avicenna J Med Biochem, The effects of the synthetic antioxidant, tempol, on serum glucose and lipid profile of diabetic and non-diabetic rats, c. e31043
17. Stopka, Czech J Food Sci, Antioxidant activity of wines and related matters studied by EPR spectroscopy, c. 49
18. Santos, J Braz Chem Soc, Antioxidant properties of plant extracts: An EPR and DFT comparative study of the reaction with DPPH, TEMPOL and spin trap DMPO, c. 1483
19. Laight, Environ Toxicol Pharmacol, Microassay of superoxide anion scavenging activity in vitro, c. 65
20. Chatterjee, Kidney Int, Tempol, a membrane-permeable radical scavenger, reduces oxidant stress-mediated renal dysfunction and injury in the rat, c. 658
21. Leker, Exp Neurol, The nitroxide antioxidant tempol is cerebroprotective against focal cerebral ischemia in spontaneously hypertensive rats, c. 355
22. Samaiya, J Neurol Sci, Tempol (4 hydroxy-tempo) inhibits anoxia-induced progression of mitochondrial dysfunction and associated neurobehavioral impairment in neonatal rats, c. 58
23. Adil, Ren Fail, Ameliorative effect of naringin in acetaminophen-induced hepatic and renal toxicity in laboratory rats: Role of FXR and KIM-1, c. 1007
24. Kandhare, Anat Cell Biol, Ameliorative effects of type-A procyanidins polyphenols from cinnamon bark in compound 48/80-induced mast cell degranulation, c. 275
25. Kandhare, Chem Biol Interact, Naringin, a flavanone glycoside, promotes angiogenesis and inhibits endothelial apoptosis through modulation of inflammatory and growth factor expression in diabetic foot ulcer in rats, c. 101
26. Li, PLoS One, The anticonvulsant effects of SR 57227 on pentylenetetrazole-induced seizure in mice, c. e93158
27. Kurian, Lancet Neurol, , Clayton PT.The monoamine neurotransmitter disorders: An expanding range of neurological syndromes, c. 721
28. Yang, Eur J Pharmacol, Involvement of 5-HT7 receptors in the pathogenesis of temporal lobe epilepsy, c. 52
29. Kourdougli, Ann Neurol, Depolarizing γ-aminobutyric acid contributes to glutamatergic network rewiring in epilepsy, c. 251
30. Mahendran, Phytomedicine, Anticonvulsant activity of embelin isolated from Embelia ribes, c. 186
31. Honmore, Pharm Biol, Artemisia pallens alleviates acetaminophen induced toxicity via modulation of endogenous biomarkers, c. 571

About this publication

Number of citations	5
Number of works in the list of references	52
Journal indexed in Scopus	Yes
Journal indexed in Web of Science	Yes

ASJC classification

3002 Drug Discovery
3003 Pharmaceutical Science

Citations

Citation Indexes: 8

Captures

Readers: 8

PLUMX View details >

Back to Academic Year 2018-2019

iorates Doxorubicin-Induced Cardiotoxicity In Experimental
e Role OfCaspase-3, COX-II, Ctn-I, Inos And TNF

An official website of the United States government
Here's how you know

FULL TEXT LINKS
SpringerLink

Mol Biol Rep. 2019 Feb;46(1):105-118. doi: 10.1007/s11033-018-4450-y. Epub 2018 Oct 25.

Fisetin, a plant flavonoid ameliorates doxorubicin-induced cardiotoxicity in experimental rats: the decisive role of caspase-3, COX-II, cTn-I, iNOs and TNF-α

Tao Ma [1], Amit D Kandhare [2], Anwesha A Mukherjee-Kandhare [2], Subhash L Bodhankar [3]

Affiliations
PMID: 30362071 DOI: 10.1007/s11033-018-4450-y

Abstract

Doxorubicin (DOX) is a widely used anthracycline antibiotic for the management of carcinoma. However, it is associated with cardiotoxicity. Fisetin is a plant flavonoid reported to have anti-inflammatory and antiapoptotic potential. To evaluate the cardioprotective potential of fisetin in DOX-induced cardiotoxicity in experimental rats. Sprague-Dawley rats were pre-treated with either fisetin (10, 20 and 40 mg/kg) or sitagliptin (10 mg/kg, p.o.) for 7 days. Cardiac toxicity was induced in rats (except the normal group) by doxorubicin (15 mg/kg i.p.) on 8th day. Various behavioral, biochemical, molecular and histological parameters were assessed in cardiac tissue. DOX-induced alterations in electrocardiographic, hemodynamic and left ventricular function were significantly ($p < 0.05$) inhibited by fisetin (20 and 40 mg/kg) treatment. Fisetin significantly decrease ($p < 0.05$) DOX-induced elevated serum CK-MB, LDH, AST, ALT and ALP levels. DOX-induced elevated cardiac oxido-nitrosative (SOD, GSH, MDA and NO) was significantly inhibited ($p < 0.05$) by fisetin. Up-regulated cardiac caspase-3, COX-II, cTn-I, iNOs, TNF-α, and IL-1β mRNA, as well as protein expressions were significantly decreased ($p < 0.05$) by fisetin treatment. It also significantly ($p < 0.05$) attenuated DOX-induced histopathological alterations in cardiac tissue. In conclusion, the fisetin exerts its cardioprotective potential against DOX-induced toxicity via inhibition of multiple pathways including oxidative stress (SOD, GSH, MDA and NO), inflammation (COX-II, TNF-α, and IL-1β), and apoptosis (Caspase-3). Therefore, fisetin can be considered as a potential cardioprotective agent during the management of carcinoma using doxorubicin anthracyclines.

Keywords: COX-II; Cardiotoxic; Caspase-3; Doxorubicin; Fisetin; cTn-I.

PubMed Disclaimer

Related information

PubChem Compound (MeSH Keyword)

LinkOut - more resources

Full Text Sources
Springer

Research Materials
NCI CPTC Antibody Characterization Program

Miscellaneous
NCI CPTAC Assay Portal

Back to Academic Year 2018-2019

Pioglitazone And Risk Of Bladder Cancer In Type 2 Diabetes Mellitus Patients: A Systematic Literature Review And Meta-Analysis Of Observational Studies Using Real-World Data

Clinical Epidemiology and Global Health 6 (2018) 61–68

Contents lists available at ScienceDirect

Clinical Epidemiology and Global Health

journal homepage: www.elsevier.com/locate/cegh

Original article

Pioglitazone and risk of bladder cancer in type 2 diabetes mellitus patients: A systematic literature review and meta-analysis of observational studies using real-world data

Mohammad Adil[a], Rashid Ali Khan[a], Pinaki Ghosh[b], Shiva Kumar Venkata[b], Amit Dattatraya Kandhare[b], Manju Sharma[a,*]

[a] Department of Pharmacology, School of Pharmaceutical Education and Research, Jamia Hamdard (Hamdard University), New Delhi, 110062, India
[b] Department of Pharmacology, Poona College of Pharmacy, Bharati Vidyapeeth Deemed University, Pune, 411038, Maharashtra, India

ARTICLE INFO

Article history:
Received 16 June 2017
Accepted 16 August 2017
Available online 30 August 2017

Keywords:
Bladder cancer
Pioglitazone
Thiazolidinedione
Systematic review
Meta-analysis

ABSTRACT

Objectives: Patients with type 2 diabetes mellitus (T2DM) have a higher incidence of bladder cancer (BC); however, the evidence underlining the association between pioglitazone use and BC risk remains inconclusive. We conducted a systematic review and meta-analysis of observational studies to investigate the effect of pioglitazone on risk of BC in T2DM patients.
Methods: We searched all publications regarding risk of BC with pioglitazone use through PubMed, Web of Science and Cochrane library databases from inception to March, 2017. Pooled hazard ratio (HR) and 95% confidence interval (CI) were calculated using a random-effect, generic inverse variance method.
Results: Total 15 observational (9 cohort and 6 case-control) studies were meta-analyzed. The pooled results showed a significant association between risk of BC and pioglitazone use (HR 1.20, 95%CI 1.09–1.31; $P < 0.0001$; $I^2 = 4\%$). In subgroup analysis, cumulative dose of pioglitazone (– and >mg) showed a significant association with risk of BC (HR 1.27; 95%CI 1.05–1.54; $P = 0.01$; $I^2 = 0\%$ and HR 1.68, 95%CI 1.36–2.08; $P < 0$ $I^2 = 0\%$ respectively). In addition, a significant association was seen with risk of BC and pioglitazone treatment duration (12–24 months and >24 months) (HR 1.43; 95%CI 1.19–1.71; $P = 0.0001$; $I^2 = 0\%$ and HR 1.58; 95%CI 1.27–1.97; $P < 0.0001$; $I^2 = 29\%$ respectively). Meta-analysis of pioglitazone *vs.* rosiglitazone use, showed a significant association (HR 1.34; 95%CI 1.05–1.71; $P = 0.02$; $I^2 = 0\%$) with BC risk and pioglitazone use.
Conclusion: Pioglitazone use is associated with risk of BC in T2DM patients. Risk of bladder cancer appears to be associated with higher dose and longer duration of pioglitazone use.
© 2017 Published by Elsevier, a division of RELX India, Pvt. Ltd on behalf of INDIACLEN.

1. Introduction

Thiazolidinediones (TZDs), pioglitazone and rosiglitazone are oral hypoglycaemic drugs used for the management of type 2 diabetes mellitus (T2DM). TZDs are insulin sensitizers, widely used for the management of T2DM. Pioglitazone belongs to the TZD group, target PPAR-γ protein, a key transcription factor for adipogenesis and glucose homeostasis.[1] On the other hand, patients with diabetes mellitus have a higher incidence of bladder cancer,[2] but the association between TZD use and bladder cancer still a matter of debate.[3,4] According to American Cancer Society, new cases of bladder cancer and deaths from bladder cancer are expected in 2016 in the United States (US).[5] Incidence of bladder cancer first time observed in a PROactive study, where 14 cases of bladder cancer were found in the pioglitazone treatment arm compared with 5 cases in the placebo group.[6] Preclinical studies reported that pioglitazone and rosiglitazone treatment could develop bladder cancer in diabetic rats.[7,8] It has been also found that pioglitazone exposure caused bladder cancer in male rats, while not observed in mice of either sex. However, mechanism behind the pioglitazone-induced urinary bladder cancer might not be due to PPAR-γ interactions because of differential variation in PPAR-γ expression subsist between the species and sex.[9] Results from a 2 year nonclinical carcinogenicity studies suggested that monotherapy of rosiglitazone were not associated with the bladder cancer.[10] Conversely, another study reveals that rosiglitazone remarkably prop up the bladder neoplasm occurrence in rats pre-treated with hydroxybutyl (butyl) nitrosamine.[7] An independent re-analysis of the RECORD trial data confirmed the original

* Corresponding author.
E-mail addresses: manjusharma72@gmail.com, msharma@jamiahamdard.ac.in (M. Sharma).

http://dx.doi.org/10.1016/j.cegh.2017.08.002
2213-3984/© 2017 Published by Elsevier, a division of RELX India, Pvt. Ltd on behalf of INDIACLEN.

Back to Academic Year 2018-2019

Development And Validation Of UV-Visible Spectroscopy Method For Simultaneous Estimation Of Saxagliptin Hydrochloride And Metformin Hydrochloride In Tablet Dosage Form

International Journal of Research in Pharmacy and Pharmaceutical Sciences

International Journal of Research in Pharmacy and Pharmaceutical Sciences
ISSN: 2455-698X
Impact Factor: RJIF 5.22
www.pharmacyjournal.in
Volume 3; Issue 4; July 2018; Page No. 31-34

Development and validation of UV-Visible spectroscopy method for simultaneous estimation of Saxagliptin hydrochloride and metformin hydrochloride in tablet dosage form

Pravin Cholke[1*], Dr. Mrunal Shirsath[2], Yogita Temak[3], Aditee Kagde[4], Rutuja Lagad[5]
[1, 3] HOD, Department of Pharmaceutical Analysis and Pharmaceutical Chemistry, Loknete Shri Dadapatil Pharate College of Pharmacy, Pune, Maharashtra, India
[2] Principal of Loknete Shri Dadapatil Pharate College of Pharmacy, Pune, Maharashtra, India
[4, 5] Students of 3rd year B.Pharmacy, Loknete Shri Dadapatil Pharate College of Pharmacy, Pune, Maharashtra, India

Abstract
A simple, rapid and validated analytical method has been developed for estimation of Saxagliptin and Metformin in tablet dosage form. The optimum conditions for the analysis of the drug were established. The maximum wavelength (λ_{max}) of Saxagliptin were found to be 274nm and Metformin 231nm. The percentage recovery of Saxagliptin(API) 100.10% and Metformin(API) 99.98%. Beer's laws were obeyed in the concentration range 50-90 µg/ml for Saxagliptin and 2-10µg/ml for Metformin. The linear equation of Saxagliptin and Metformin was calculated. The method was validated in terms of linearity, accuracy, precision, specificity, limit of detection and limit of quantitation. Validation was performed as per ICH guidelines. The proposed method was successfully applied for the quantitative determination of Metformin and Saxagliptin in tablet dosage form (percentage label claim 99.60% and 100% respectively).

Keywords: Saxagliptin, metformin, simultaneous estimation

Introduction

Gliptin
Dipeptidyl peptidase-4 inhibitor (DPP-4) they are prescribed to type-2 Diabetes patients. DPP-4 inhibitor may help with weight loss as well as decrease blood glucose level.
Derivatives of Gliptins are
a. Alogliptin
b. Linagliptin
c. Saxagliptin
d. Sitagliptin.

Saxagliptin
This is new oral hypoglycemic of the new DPP-4 inhibitor class of drugs. Saxagliptin is chemically identify as (1S, 3S, 5S)-2-[(2S)-2-Amino-2-(3 Hydroxytricyclo[3.3.1.13,7]dec-1-yl)acetyl]-2 azabicyclo[3.1.0]hexane-3-carbonitrile previously identified as BMS-477118.
Empirical formula is $C_{18}H_{25}N_3O_2$, H_2O and molecular weight 333.43. It is sparingly soluble in water and soluble in methanol.
Structural formula of Saxagliptin is as follows



Saxagliptin is used for treatment of type-2 Diabetes Mellitus and also used to improve glycaemic control in type-2 Diabetes Mellitus patients. Saxagliptin is used with Metformin, Sulphonylurea or Pioglytazone, when blood sugar level is not controlled by one of these agents alone. Literature survey reveals that the drug can be estimated only by LC-MS/MS, Spectrophotometric method.

Metformin Hydrochloride
It is water soluble compound and the anti-diabetic drug from the biguanide class of oral hypoglycemic agent. It chemically identify as 1, 1-dimehtylbiguanidine monohydrochloride and empirical formula is $C_4H_{12}N_5CI$, molecular weight is 165.625 gm/mol. Metformin hydrochloride increase glucose transport across the cell membrane in skeletal muscle. Spectroflourimetry, RP-HPLC, HPTLC, LC-MS/MS and UV-Visible Spectroscopy methods was reported for determination of Metformin.
Structural formula of Metformin:



Literature survey reveals that various methods were reported for single estimation of Saxagliptin and Metformin, but no spectroscopic method has been reported for the analysis of

Back to Academic Year 2018-2019

ISSN: 0975-833X

Available online at http://www.journalcra.com

International Journal of Current Research
Vol. 10, Issue, 07, pp.71025-71031, July, 2018

INTERNATIONAL JOURNAL OF CURRENT RESEARCH

RESEARCH ARTICLE

FORMULATION AND EVALUATION OF MOUTH DISSOLVING TABLET OF BENDROFLUMETHIAZIDE

[1]Sonali A. Nagare, [1,*]Anita A. Najan, [2]Mrunal K, Shirsat and [3]Avinash V. Dhobale

[1]Assistant Professor, Pharmaceutical Chemistry, Loknete Shri Dadapatil Pharate college of Pharmacy, Mandavgan Pharata, Tq-Shirur Dist-Pune
[2]Loknete Shri Dadapatil Pharate college of Pharmacy, Mandavgan Pharata, Tq-Shirur Dist-Pune
[3]Assistant Professor, Pharmaceutics, Loknete Shri Dadapatil Pharate college of Pharmacy, Mandavgan Pharata, Tq-Shirur Dist-Pune

ARTICLE INFO

Article History:
Received 28th April, 2018
Received in revised form
07th May, 2018
Accepted 14th June, 2018
Published online 30th July, 2018

Key Words:
Mouth Dissolving Tablets,
Bendroflumethiazide,
Super-Disintegrates,
Croscarmellose Sodium,
Sodium Starch Glycolate.

ABSTRACT

In the present research work, mouth dissolving tablets of bendroflumethiazide were prepared by direct compression method with a view to enhance patient compliance. Mouth dissolving tablets have many numbers of advantages over the conventional tablets like rapid disintegration, faster dissolution, Ease of administration and quick onset of action etc. Two super-disintegrants, viz., croscarmellose sodium and sodium starch glycolate in different ratios with microcrystalline cellulose along with mannitol to enhance mouth feel. Many pharmaceutical dosages are administered in the form of pills, granules, powders, and liquids. Generally, a pill is designed for swallowing intact or chewing to deliver a precise dosage of medication to patients. The drug and excipient interaction study was carried out by taking Infrared Spectrum of Pure drug and optimized formulation (B2). There was no change in the prominent functional groups of indicating drug is in intact form. The prepared batches of tablets were evaluated for hardness, friability, drug content, wetting time, water absorption ratio, disintegration time and dissolution study.

Copyright © 2018, Sonali A. Nagare et al. *This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.*

***Citation: Sonali A. Nagare, Anita A. Najan, Mrunal K, Shirsat and Avinash V. Dhobale*, 2018.** "Formulation and evaluation of mouth dissolving tablet of bendroflumethiazide", *International Journal of Current Research*, 10, (07), 71025-71031.

INTRODUCTION

The most commonly used dosage form for pharmaceutical preparations is currently the tablet, available in various forms and administered orally. The advantages of this dosage form are manifold: tablets are cost effective to manufacture, convenient to dispense and store, easy for the patient to administer and they provide a versatile means of delivering the drug. Release of drug from the tablet can be controlled by altering the design and content of the formulation. Also, since this is a dry dosage form, tablets provide a supportive environment for drug stability and generally have a relatively long shelf life (Bharat Parashar *et al.*, 2012). The properties of the tablet (e.g. mechanical strength, disintegration time and drug release characteristics) are affected by both the properties of the constituent materials and the manufacturing process. Excipients such as diluents, binders and lubricants are generally needed in a formulation in order to facilitate the manufacturing process, but also to ensure that the resulting tablets have the desired properties. For instance, tablets should be sufficiently strong to withstand handling during manufacturing and usage, but should also disintegrate and release the drug in a predictable and reproducible manner (Rudnic, ?). Drug Delivery Systems (DDS) are a strategic tool for expanding markets/indications, extending product life cycles and generating opportunities. DDS make a significant contribution to global pharmaceutical sales through market segmentation and are moving rapidly. Drug delivery systems are becoming increasingly sophisticated as scientists acquire a better understanding of the physicochemical parameters pertinent to their performance (Chein, 1992). Despite of tremendous advancements in drug delivery, the oral route remains the perfect route for the administration of therapeutic agents because of low cost of therapy, ease of administration, self-medication, leading to high levels of patient compliance. Tablets and capsules are the most popular dosage forms [4]. It is a tablet that disintegrates and dissolves rapidly in the saliva, within a few seconds without the need of drinking water or chewing. A mouth dissolving tablet usually dissolves in the oral cavity within 15 second to 3min.

Corresponding author: Assistant Professor, Pharmaceutical Chemistry, Loknete Shri Dadapatil Pharate college of Pharmacy, Mandavgan Pharata, Tq-Shirur Dist-Pune
DOI: https://doi.org/10.24941/ijcr.31484.07.2018

Back to Academic Year 2018-2019

Indo American Journal of Pharmaceutical Research, 2018 ISSN NO: 2231-6876

INDO AMERICAN JOURNAL OF PHARMACEUTICAL RESEARCH

"FORMULATION AND EVALUATION OF CARBAMAZEPINE NANOEMULSION FOR BRAIN TARGETED DRUG DELIVERY VIA INTRANASAL ROUTE"

Nagare Sonali .A*, Dhobale Avinash.V, Najan Anita.A

Loknete Shri Dadapatil Pharate College of Pharmacy, Mandavgan pharata,Shirur,Dist Pune,412211.

ARTICLE INFO

Article history
Received 16/07/2018
Available online
31/07/2018

Keywords
Carbamazepine,
Nanoemulsion,
Intranasal To Brain.

ABSTRACT

Intranasal drug administration is receiving increased attention as adelivery method for by-passing the blood–brain barrier and rapidly targeting therapeutics to the brain or CNS. The objective of the present study was to selected carbamazepine nanoemulsion for nose-to-brain delivery. Carbamazepine nanoemulsion (NE) formulation were successfully prepared by the spontaneous emulsification method (titration method) using Capmul MCM as the oil, Tween-80 as surfactant, and PEG-600 as co-surfactant phase on the basis of solubility studies. The nanoemulsion formulation containing 7.35% oil, 66.18% Smix ratio (3:1 Tween-80:PEG-600 ratio),26.47% (v/v) aqueous phase that displayed an optical transparency of 99.42±0.81%, globule size of 71.70±3.06 nm, and polydispersity index of 0.256±0.002. The selected Carbamazepine nanoemulsion was characterized, and the *in-vitro* drug release and *in-vivo* nasal absorption of drug from the selected formulation were evaluated in rats. *In-vitro* and ex vivo permeation studies showed an initial burst of drug release at 60 min and Carbamazepine nanoemulsion show drug release up to 5 h. In vivo pharmacokinetic studies in rats showed that Carbamazepine.

Corresponding author
Nagare Sonali
Loknete Shri Dadapatil Pharate College of Pharmacy,
Mandavgan Pharata, Shirur, Dist Pune, 412211.
sonalinagare93@gmail.com

*Please cite this article in press as **Nagare Sonali** et al. "Formulation and Evaluation of Carbamazepine Nanoemulsion for Brain Targeted Drug Delivery Via Intranasal Route". Indo American Journal of Pharmaceutical Research.2018:8(07).*

Copy right © 2018 This is an Open Access article distributed under the terms of the Indo American journal of Pharmaceutical Research, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

ISSN: 0975-833X

Available online at http://www.journalcra.com

International Journal of Current Research
Vol. 10, Issue, 07, pp.71025-71031, July, 2018

INTERNATIONAL JOURNAL OF CURRENT RESEARCH

RESEARCH ARTICLE

FORMULATION AND EVALUATION OF MOUTH DISSOLVING TABLET OF BENDROFLUMETHIAZIDE

[1]Sonali A. Nagare, [1,*]Anita A. Najan, [2]Mrunal K, Shirsat and [3]Avinash V. Dhobale

[1]Assistant Professor, Pharmaceutical Chemistry, Loknete Shri Dadapatil Pharate college of Pharmacy, Mandavgan Pharata, Tq-Shirur Dist-Pune
[2]Loknete Shri Dadapatil Pharate college of Pharmacy, Mandavgan Pharata, Tq-Shirur Dist-Pune
[3]Assistant Professor, Pharmaceutics, Loknete Shri Dadapatil Pharate college of Pharmacy, Mandavgan Pharata, Tq-Shirur Dist-Pune

ARTICLE INFO

Article History:
Received 28th April, 2018
Received in revised form
07th May, 2018
Accepted 14th June, 2018
Published online 30th July, 2018

Key Words:
Mouth Dissolving Tablets,
Bendroflumethiazide,
Super-Disintegrates,
Croscarmellose Sodium,
Sodium Starch Glycolate.

ABSTRACT

In the present research work, mouth dissolving tablets of bendroflumethiazide were prepared by direct compression method with a view to enhance patient compliance. Mouth dissolving tablets have many numbers of advantages over the conventional tablets like rapid disintegration, faster dissolution, Ease of administration and quick onset of action etc. Two super-disintegrants, viz., croscarmellose sodium and sodium starch glycolate in different ratios with microcrystalline cellulose along with mannitol to enhance mouth feel. Many pharmaceutical dosages are administered in the form of pills, granules, powders, and liquids. Generally, a pill is designed for swallowing intact or chewing to deliver a precise dosage of medication to patients. The drug and excipient interaction study was carried out by taking Infrared Spectrum of Pure drug and optimized formulation (B2). There was no change in the prominent functional groups of indicating drug is in intact form. The prepared batches of tablets were evaluated for hardness, friability, drug content, wetting time, water absorption ratio, disintegration time and dissolution study.

Copyright © 2018, Sonali A. Nagare et al. *This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.*

***Citation: Sonali A. Nagare, Anita A. Najan, Mrunal K, Shirsat and Avinash V. Dhobale*, 2018.** "Formulation and evaluation of mouth dissolving tablet of bendroflumethiazide", *International Journal of Current Research*, 10, (07), 71025-71031.

INTRODUCTION

The most commonly used dosage form for pharmaceutical preparations is currently the tablet, available in various forms and administered orally. The advantages of this dosage form are manifold: tablets are cost effective to manufacture, convenient to dispense and store, easy for the patient to administer and they provide a versatile means of delivering the drug. Release of drug from the tablet can be controlled by altering the design and content of the formulation. Also, since this is a dry dosage form, tablets provide a supportive environment for drug stability and generally have a relatively long shelf life (Bharat Parashar *et al*., 2012). The properties of the tablet (e.g. mechanical strength, disintegration time and drug release characteristics) are affected by both the properties of the constituent materials and the manufacturing process. Excipients such as diluents, binders and lubricants are generally needed in a formulation in order to facilitate the manufacturing process, but also to ensure that the resulting tablets have the desired properties. For instance, tablets should be sufficiently strong to withstand handling during manufacturing and usage, but should also disintegrate and release the drug in a predictable and reproducible manner (Rudnic, ?). Drug Delivery Systems (DDS) are a strategic tool for expanding markets/indications, extending product life cycles and generating opportunities. DDS make a significant contribution to global pharmaceutical sales through market segmentation and are moving rapidly. Drug delivery systems are becoming increasingly sophisticated as scientists acquire a better understanding of the physicochemical parameters pertinent to their performance (Chein, 1992). Despite of tremendous advancements in drug delivery, the oral route remains the perfect route for the administration of therapeutic agents because of low cost of therapy, ease of administration, self-medication, leading to high levels of patient compliance. Tablets and capsules are the most popular dosage forms [(4)]. It is a tablet that disintegrates and dissolves rapidly in the saliva, within a few seconds without the need of drinking water or chewing. A mouth dissolving tablet usually dissolves in the oral cavity within 15 second to 3min.

Corresponding author: Assistant Professor, Pharmaceutical Chemistry, Loknete Shri Dadapatil Pharate college of Pharmacy, Mandavgan Pharata, Tq-Shirur Dist-Pune
DOI: https://doi.org/10.24941/ijcr.31484.07.2018

Back to Academic Year 2018-2019

Academic Year 2019

Sr. no	Title of the paper	Name of Staff	Department	Name of the journal	ISSN number	Document	Link

1	Ferulic acid ameliorates doxorubicin-induced cardiac toxicity in rats	Dr. A. D. Kandhare	Pharmacology	Naunyn-Schmiedeberg's Arch Pharmacol	https://doi.org/10.1007/s00210-019-01623-4	View Document	https://www.researchgate.net/publication/330967015_Ferulic_acid_ameliorates_doxorubicin-induced_cardiac_toxicity_in_rats
2	Diarylheptanoid, a constituent isolated from methanol extract of Alpinia officinarum attenuates TNF-α level in Freund's complete adjuvant-induced arthritis in rats	Dr. A. D. Kandhare	Pharmacology	Journal of Ethnopharmacology	https://doi.org/10.1016/j.jep.2018.10.019	View Document	https://www.researchgate.net/publication/328300349_Diarylheptanoid_a_constituent_isolated_from_methanol_extract_of_Alpinia_officinarum_attenuates_TNF-a_level_in_Freund's_complete_adjuvant-induced_arthritis_in_rats
3	Efficacy of Antioxidant Supplements on Prevention and Ameliorati on of Cisplatin-Induced Nephrotoxicity: A Systematic Review and Meta-analysis of Randomize	Dr. A. D. Kandhare	Pharmacology	Jundishapur J Nat Pharm	Jundishapur J Nat Pharm Prod	View Document	https://www.researchgate.net/publication/334747602_Efficacy_of_Antioxidant_Supplements_on_Prevention_and_Amelioration_of_Cisplatin-Induced_Nephrotoxicity_A_Systematic_Review_and_Meta-analysis_of_Randomized_Controlled_Trials

	d Controlled Trials						
4	A systematic literature review of fenugreek seed toxicity by using ToxRTool: evidence from preclinical and clinical studies	Dr. A. D. Kandhare	Pharmacology	Elsevier Ltd	https://doi.org/10.1016/j.heliyon.2019.e01536	View Document	https://www.researchgate.net/publication/332619750_A_systematic_literature_review_of_fenugreek_seed_toxicity_by_using_ToxRTool_evidence_from_preclinical_and_clinical_studies
5	Therapeutic Potential of Morin in Ovalbumin-induced Allergic Asthma Via Modulation of SUMF2/IL-13 and BLT2/NF-kB Signaling Pathway	Dr. A. D. Kandhare	Pharmacology	Current Molecular Pharmacology	Current Molecular Pharmacology, 2019, Vol. 12,	View Document	https://www.researchgate.net/publication/330122445_Therapeutic_Potential_of_Morin_in_Ovalbumin-induced_Allergic_Asthma_Via_Modulation_of_SUMF2IL-13_and_BLT2NF-kB_Signaling_Pathway
6	Hesperidin Ameliorates Bleomycin-Induced Experimental Pulmonary Fibrosis Via Inhibition Of	Dr. A. D. Kandhare	Pharmacology	EXCLI Journal	ISSN 1611-2156	View Document	https://www.researchgate.net/publication/336589331_Hesperidin_ameliorates_bleomycin-induced_experimental_pulmonary_fibrosis_via_inhibition_of_TGF-beta1Smad3AMPK_and_IkappaBalphaNF-kappaB_pathways

	Tgf-B1/Smad3/Ampk And Iκbα/Nf-Kb Pathways						
7	Acute and sub-chronic oral toxicity studies of hesperidin isolated from orange peel extract in Sprague Dawley rats	Dr. A. D. Kandhare	Pharmacology	Regulatory Toxicology and Pharmacology	www.elsevier.com/locate/yrtph	View Document	https://www.researchgate.net/publication/332404120_Acute_and_sub-chronic_oral_toxicity_studies_of_hesperidin_isolated_from_orange_peel_extract_in_Sprague_Dawley_rats
8	Development and Validation of RP-HPLC and UV Spectrophotometric Absorptivity Method for Simultaneous Estimation of Cyclobenzaprine hydrochloride and Aceclofenac in Pharmaceutical dosage form	Mr. A. S. Gaikwad	Pharmaceutical Chemistry	International Journal of Scientific Development and Research	ISSN: 2455-2631	View Document	https://www.researchgate.net/publication/374813338_Development_and_Validation_of_RP-HPLC_and_UV-_Spectrophotometric_Absorptivity_Method_for_Simultaneous_Estimation_of_Cyclobenzaprine_hydrochloride_and_Aceclofenac_in_Pharmaceutical_dosage_form

10	A Review - Corona Virus (Covid-19)	Mr. A. S. Gaikwad	Pharmaceutical Chemistry	International Journal for Research Trends and Innovation	ISSN: 2456-3315	View Document	https://scholar.google.com/citations?view_op=view_citation&hl=en&user=ORlyYt4AAAAJ&citation_for_view=ORlyYt4AAAAJ:YOwf2qJgpHMC
11	Targeted Drug Delivery – From Magic Bullet to Nanomedicine	Mr. A. S. Gaikwad	Pharmaceutical Chemistry	Journal of Xidian University	ISSN No:1001-240	View Document	https://scholar.google.com/citations?view_op=view_citation&hl=en&user=ORlyYt4AAAAJ&citation_for_view=ORlyYt4AAAAJ:kNdYIx-mwKoC
12	Cancer Chemotherapy with Peptides & Peptidomimetics drug & Peptide based Vaccine	Ms. S. A. Nagare	Pharmaceutical Chemistry	International Research Journal of Pharmacy and Medical Sciences	ISSN2581-3277	View Document	https://www.researchgate.net/publication/330957007_Cancer_Chemotherapy_with_Peptides_and_Peptidomimetics_Drug_and_Peptide_Based-Vaccines
13	A study to assess the knowledge regarding hand hygiene practice among secondary school going children in selected schools of Pune city	Ms. S. B. Kshirsagar	Pharmaceutical Chemistry	International Journal of Applied Research 2020; 6(12): 27-29	ISSN : 2394-7500	View Document	https://www.researchgate.net/publication/362517268_A_study_to_assess_the_knowledge_regarding_hand_hygiene_practice_among_secondary_school_going_children_in_selected_schools_of_Pune_city

14	Efficacy of Antioxidant Supplements on Prevention and Amelioration of Cisplatin-Induced Nephrotoxicity: A Systematic Review and Meta-analysis of Randomized Controlled Trials	Ms. A. A. Mukharjee	Pharmacology	Jundishapur J Nat Pharm	Jundishapur J Nat Pharm Prod	View Document	https://www.researchgate.net/publication/334747602_Efficacy_of_Antioxidant_Supplements_on_Prevention_and_Amelioration_of_Cisplatin-Induced_Nephrotoxicity_A_Systematic_Review_and_Meta-analysis_of_Randomized_Controlled_Trials
15	Therapeutic Potential of Morin in Ovalbumin-induced Allergic Asthma Via Modulation of SUMF2/IL-13 and BLT2/NF-kB Signaling Pathway	Ms. A. A. Mukharjee	Pharmacology	Current Molecular Pharmacology	Current Molecular Pharmacology, 2019, Vol. 12,	View Document	https://www.researchgate.net/publication/330122445_Therapeutic_Potential_of_Morin_in_Ovalbumin-induced_Allergic_Asthma_Via_Modulation_of_SUMF2IL-13_and_BLT2NF-kB_Signaling_Pathway
16	Acute and sub-chronic oral toxicity studies of hesperidin isolated from orange peel	Ms. A. A. Mukharjee	Pharmacology	Regulatory Toxicology and Pharmacology	www.elsevier.com/locate/yrtph	View Document	https://www.researchgate.net/publication/332404120_Acute_and_sub-chronic_oral_toxicity_studies_of_hesperidin_isolated_from_orange_peel_extract_in_Sprague_Dawley_rats

	extract in Sprague Dawley rats						

Back to Academic Year 2019-2020

Ferulic acid ameliorates doxorubicin-induced cardiac toxicity in rats

Back to Academic Year 2019-2020

armacology
23-4

CrossMark

Ferulic acid ameliorates doxorubicin-induced cardiac toxicity in rats

Urmila Aswar[1] · Umesh Mahajan[2] · Amit Kandhare[1] · Manoj Aswar[2]

Received: 28 August 2018 / Accepted: 28 January 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Abstract
Ferulic acid (FA) is a phenolic compound with potent antioxidant activity. The objective of the study was to study the protective effects of FA on doxorubicin (Dox)-induced myocardial toxicity in rats. Wistar rats received vehicle (control) or Dox (20 mg/kg, i.p.) or telmisartan (Tel; 10 mg/kg, p.o.) or ferulic acid (20 mg/kg and 40 mg/kg, p.o.) for 7 days followed by treatment with Dox (20) on the fifth day of treatment, except the control group. On day 8, electrocardiographic parameters were recorded followed by blood withdrawal and then the animals were sacrificed for histopathology. Administration of Dox showed prolonged RR, QTc interval, and QRS complex. The levels of serum CK-MB, LDH, IL-1β, and IL-6 were significantly increased ($p < 0.01$). Similarly, levels of Ca^{+2}, Mg^{+2} ATPase, and Ca^{+2} ATPase and expression of ANP and BNP were significantly higher as compared to the control. In the FA-treated group, ECG was normal. The serum levels of CK-MB, LDH, IL-1β, and IL-6 were not elevated. Heart tissue Ca^{+2}, Mg^{+2} ATPase, and Ca^{+2} ATPase did not show a statistical difference compared to the control group. The FA treatment attenuated the expression of ANP and BNP. FA (20 and 40) augmented myocardial GSH and Na^{+}/K^{+} ATPase. Histopathology of the heart confirmed the cardioprotective effect of FA.

Keywords Anti-oxidant · Cardiac toxicity · Doxorubicin · Ferulic acid · Rats

Introduction

Doxorubicin (Dox) is an anthracycline derivative, clinically used for treatment of a variety of cancerous growth such as breast cancer, lung cancer, and acute leukemias (Hortobagyi 1997; Pugazhendhi et al. 2018). Cardiotoxicity is the severe adverse effect associated with its use. It is speculated to be mediated via mitochondrial dysfunction (Singal and Iliskovic 1998). Dox-induced cardiac toxicity is a multifactorial process that includes oxidative stress, thereby stimulating lipid peroxidation (Vallejo et al. 2017), iron metabolism, inflammation, ER stress, mitochondrial permeability transition, loss of mitochondrial integrity, and function which is essential for Ca homeostasis and signaling, as well as activation of the renin-angiotensin system (Yu et al. 2018). These pathological consequences restrict clinical use of Dox due to its dose-dependent cardiotoxicity leading to the electrocardiographic irregularity, arrhythmias, cardiomyopathy, and congestive heart failure (Pugazhendhi et al. 2018). It has also been reported that approximately 10% of cancer survivors treated with Dox or its derivatives may develop cardiac complications, even when chemotherapy is stopped (Octavia et al. 2012).

Preclinical studies have evaluated several molecules, such as β-receptor blockers, angiotensin receptor blockers, amifostine, dexrazoxane, mesna, leucovorin, and erythropoietin, as cardio-protective adjuvants (Ludke et al. 2009; Alimoradi et al. 2012). A variety of plant-derived bioactive also have been proved to attenuate Dox-induced cardiac toxicity such as p-coumaric acid (Abdel-Wahab et al. 2003), curcumin, green tea, garlic or L-carnitine (Radwan et al. 2012), thymoquinone (Nagi and Almakki 2009), alpha-lipoic acid (Al-Majed et al. 2002), and resveratrol (Oktem et al. 2012). However, no molecule is approved clinically due to severe adverse effects and complex pharmacokinetics of natural compounds in the human body. Ferulic acid [FA (3-(4-hydroxy-3-methoxy-phenyl) prop-2-enoic acid)] is a phenolic compound present in most of the plants (Maurya and Devasagayam 2010). It readily forms resonance stabilized phenoxy radical that might be responsible for its strong anti-oxidant activity. FA absorbs UV radiation and forms a stable

✉ Urmila Aswar
aswarurmila@gmail.com

[1] Department of Pharmacology, Poona College of Pharmacy, Bharati Vidyapeeth Deemed University, Erandwane, Pune, Maharashtra 411038, India

[2] Department of Pharmacology, Sinhgad Institute of Pharmacy, Narhe, Pune, Maharashtra 411041, India

Published online: 08 February 2019

Back to Academic Year 2019-2020

Diarylheptanoid, a constituent isolated from methanol extract of Alpinia officinarum attenuates TNF-α level in Freund's complete adjuvant-induced arthritis in rats

Journal of Ethnopharmacology 229 (2019) 233–245

Contents lists available at ScienceDirect

Journal of Ethnopharmacology

journal homepage: www.elsevier.com/locate/jethpharm

Diarylheptanoid, a constituent isolated from methanol extract of *Alpinia officinarum* attenuates TNF-α level in Freund's complete adjuvant-induced arthritis in rats

Varsha S. Honmore[a,1], Amit D. Kandhare[b], Parag P. Kadam[b,1], Vijay M. Khedkar[c], Arun D. Natu[a,*], Supada R. Rojatkar[d,*], Subhash L. Bodhankar[b,*]

[a] Post Graduate and Research Center, Department of Chemistry, MES Abasaheb Garware College, Pune, Maharashtra, India
[b] Department of Pharmacology, Poona College of Pharmacy, Bharati Vidyapeeth Deemed University, Pune, Maharashtra, India
[c] Department of Pharmaceutical Chemistry, SVKM's Institute of Pharmacy, Survey No. 499, Plot No, 03, Mumbai - Agra National Highway, Maharashtra 424001, India
[d] R&D Centre in Pharmaceutical Sciences and Applied Chemistry, Poona College of Pharmacy, Bharati Vidyapeeth Deemed University, Pune, Maharashtra, India

ARTICLE INFO

Keywords:
1-phenyl-5-hydroxy-7- (4″-hydroxy-3″-methoxyphenyl) heptane-3-one (i.e. 5-HPH)
Alpinia officinarum (Zingiberaceae)
Complete Freund's adjuvant-induced arthritis
Interleukin (IL)-1β
Molecular docking
Oxidative stress
Tumor necrosis factor (TNF)-α

ABSTRACT

Ethnopharmacological relevance: Rheumatoid arthritis (RA) is a chronic inflammatory and destructive joint disease that affects the worldwide population. *Alpinia officinarum* Hance (Zingiberaceae), rhizomes are widely used ethnobotanically as an anti-inflammatory, analgesic, and antioxidant agent in traditional medicine.
Aim: To investigate the efficacy and possible mechanism of isolated phytoconstituent from the methanol extract of *A. officinarum* (MEAO) rhizomes against Freund's complete adjuvant (FCA)-induced arthritis in rats. Furthermore, molecular docking was performed to study the binding mode of this compound into the active site of TNF-α.
Materials and methods: Diarylheptanoid was isolated from MEAO, well characterized (HPTLC, ^{1}H NMR, ^{13}C NMR, and ESI-MS) and evaluated for its antiarthritic activity in female Wistar rats (170–200 g). Diarylheptanoid (5, 10 and 20 mg/kg, p.o.) was administered starting from day 12. Various behavioral, biochemical, molecular and histopathology parameters were evaluated. Molecular docking study was performed using Glide module integrated into Schrodinger molecular modeling software.
Results: The structure and molecular weight of the isolated compound (diarylheptanoid) were confirmed by 1D and mass spectral data and characterized as 1-phenyl-5-hydroxy-7- (4″-hydroxy-3″-methoxyphenyl) heptane-3-one (i.e., 5-HPH) with molecular formula $C_{20}H_{24}O_4$. Administration of 5-HPH (10 and 20 mg/kg) significantly inhibited ($p < 0.05$) FCA induced increases in paw volume, joint diameter, thermal hyperalgesia and tactile allodynia. It also significantly decreased oxido-inflammatory markers (SOD, GSH, MDA, and TNF-α). FCA induced a histological alteration in ankle joint also attenuated by 5-HPH. Its Glide docking score was found to be −9.702 with binding energy (Glide energy) of −37.033 kcal/mol.
Conclusion: 5-HPH may exhibit its anti-arthritic potential via inhibition of elevated oxido-inflammatory markers thus restoring the elevated hyperalgesia, allodynia and reducing destruction in synovial membrane and cartilage. Therefore, 5-HPH is a potential moiety bearing antioxidant and with anti-inflammatory properties to inhibit FCA-induced arthritis in rats. The results of the present investigation should enable the design of potent small-molecule inhibitors that inactivate TNF-α with high affinity and specificity.

Abbreviations: AIA, Adjuvant-Induced Arthritis; ALP, Alkaline Phosphatase; ALT, Alanine Aminotransferase; AST, Aspartate Aminotransferase; ESI-MS, Electrospray Ionization-Mass Spectrometry; ELISA, Enzyme-Linked Immunosorbent Assay; FCA, Freund's Complete Adjuvant; GSH, Reduced Glutathione; 5-HPH, 1-phenyl-5-hydroxy-7- (4″-hydroxy-3″-methoxyphenyl) heptane-3-one; Malonadehyde MDA, Lipid Peroxidation; MEAO, Methanol Extract of A. officinarum; ROS, Reactive Oxygen Species; SOD, Superoxide Dismutase; TNF-α, Tumor necrosis factor-alpha; IL's, Interleukin's

* Corresponding authors.
E-mail address: subhash.bodhankar@bharatividyapeeth.edu (S.L. Bodhankar).
[1] Authors contributed equally.

https://doi.org/10.1016/j.jep.2018.10.019
Received 11 May 2017; Received in revised form 17 June 2018; Accepted 12 October 2018
Available online 15 October 2018
0378-8741/ © 2018 Elsevier B.V. All rights reserved.

Back to Academic Year 2019-2020

Efficacy of Antioxidant Supplements on Prevention and Amelioration of Cisplatin-Induced Nephrotoxicity: A Systematic Review and Meta-analysis of Randomized Controlled Trials

Corrected Proof

Jundishapur J Nat Pharm Prod. 2019 August; 14(3):e61527. doi: 10.5812/jjnpp.61527.
Published online 2019 July 29. **Review Article**

Efficacy of Antioxidant Supplements on Prevention and Amelioration of Cisplatin-Induced Nephrotoxicity: A Systematic Review and Meta-analysis of Randomized Controlled Trials

Amit D. Kandhare [1], Anwesha Mukherjee [1] and Subhash L. Bodhankar [1, *]

[1] Department of Pharmacology, Poona College of Pharmacy, Bharati Vidyapeeth Deemed University, Erandwane, India

[*] *Corresponding author:* Department of Pharmacology, Poona College of Pharmacy, Bharati Vidyapeeth Deemed University, Erandwane, Paud Road, Pune-411 038, Maharashtra, India. Tel: +91-2025437237, Fax: +91-2025439383, Email: drslbodh@gmail.com

Received 2017 September 10; **Accepted** 2018 April 22.

Abstract

Context: Cisplatin is a widely used antineoplastic agent in the treatment of a wide range of malignancies although it is associated with nephrotoxicity. Much clinical evidence supports the use of antioxidant supplements in the prevention of cisplatin-induced nephrotoxicity (CIN). However, conflicting evidence makes us unable to provide any robust results for antioxidants use against CIN.
Objectives: The study aimed to investigate the efficacy of antioxidant supplements on CIN through a comprehensive meta-analysis of randomized controlled trials.
Data Sources: A systematic literature search was conducted in PubMed, EMBASE, Cochrane Library, CPCI-S (Conference Proceedings Citation Index-Science), ICTRP (International Clinical Trials Registry Platform), and Google Scholar until February 2017 by two independent researchers. Various outcomes such as serum creatinine, estimated glomerular filtration rate (eGFR), blood urea nitrogen (BUN), creatinine clearance, and incidence of CIN were assessed. All statistical analyses were performed using RevMan V.5.3
Results: Overall, 672 patients were identified from 10 studies of whom 330 (49.10%) patients received antioxidant treatment. Antioxidant treatment showed a significant reduction in serum creatinine (SMD: -3.40, 95% CI: -5.47 to -1.33, P = 0.001), BUN (SMD = -5.96, 95% CI: -10.07 to -1.86, P = 0.004), and eGFR (SMD = -3.77, 95% CI: -6.16 to -1.38; P = 0.002) when compared to the control group.
Conclusions: Antioxidant treatment is associated with a reduced risk of CIN. It also has important clinical implications for CIN patients who are not responding to other therapies such as hydration, diuresis, or magnesium supplementation.

Keywords: Antioxidant, Cisplatin, Meta-Analysis, Nephrotoxicity, Systematic Review

1. Context

Cisplatin (cis-diamminedichloroplatinum [II]; CDDP; Platinol) is a platinum-based antineoplastic agent that serves a highly effective treatment regimen for an array of malignancies such as head and neck cancer, cervical cancer, soft-tissue neoplasms, squamous cell cancer, non-small cell lung cancer, gastric cancer, testicular cancer, bladder cancer, and ovarian cancer (1, 2). Despite its effectiveness, the clinical use of cisplatin is compromised in up to 85% of cases due to severe side effects including ototoxicity, nephrotoxicity, bone marrow toxicity, gastrointestinal toxicity, and peripheral neuropathy (3). The prevalence of cisplatin nephrotoxicity was reported as 34.1% among various cancer patients in a study (4) and the incidence of cisplatin-induced nephrotoxicity was around 30% - 40% in another study (5), which was dose-dependent and usually reversible. It has been reported that CDDP, as a platinum-based alkylating compound, has an ability to interact with DNA to form interstrand cross-links and intrastrand bifunctional N-7 DNA adducts at d(GpG) and d(ApG) (6). The formation of these adducts can result in DNA damage, oxidative stress, protein synthesis inhibition, and mitochondrial dysfunction (7).

The current treatment strategy for cisplatin-induced nephrotoxicity (CIN) manifestations primarily includes supportive care with sodium chloride or bicarbonate volume expansion, metformin withdrawal, administration of various agents such as nonsteroidal anti-inflammatory drugs, angiotensin-converting enzyme inhibitors, angiotensin II-receptor blockers and statins, and reduced volume of contrast media (8). CIN can be ameliorated using oral hydration therapy (9) or magnesium supplementation (7); however, it is reversible and does not completely prevent CIN. Dialysis has been successfully implicated in the management of various features of CIN

Copyright © 2019, Jundishapur Journal of Natural Pharmaceutical Products. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.

Back to Academic Year 2019-2020

A Systematic literature review of fenugreek seed by using ToxRTool;evidence from preclinical & clinical studies

Heliyon

ELSEVIER

Received:
13 July 2018
Revised:
20 November 2018
Accepted:
15 April 2019

Cite as: Amit D. Kandhare, Prasad A. Thakurdesai, Pralhad Wangikar, Subhash L. Bodhankar. A systematic literature review of fenugreek seed toxicity by using ToxRTool: evidence from preclinical and clinical studies. Heliyon 5 (2019) e01536. doi: 10.1016/j.heliyon.2019.e01536

A systematic literature review of fenugreek seed toxicity by using ToxRTool: evidence from preclinical and clinical studies

Amit D. Kandhare[a,*], **Prasad A. Thakurdesai**[*,a], **Pralhad Wangikar**[b], **Subhash L. Bodhankar**[c]

[a] *Indus Biotech Private Limited, Pune, India*

[b] *Prado-Preclinical Research and Development Organization Pvt. Ltd., Pune, India*

[c] *Department of Pharmacology, Poona College of Pharmacy, Bharati Vidyapeeth Deemed University, Pune, India*

[*] Corresponding authors.

E-mail addresses: amit.kandhare@indusbiotech.com (A.D. Kandhare), prasad@indusbiotech.com (P.A. Thakurdesai).

Abstract

Fenugreek (*Trigonella foenum graecum*) seed extract is a bioactive ingredient of many food supplements. Hence, there is a need for systematic assessment of the quality of published toxicological studies for its use in human health, hazard consideration, and risk assessment. The aim of the present investigation was to determine the reliability of published toxicological studies of fenugreek seed by using ToxRTool (Toxicological data reliability assessment tool). A comprehensive systematic literature search was conducted in PubMed, EMBASE, Cochrane Library, CPCI-S, ICTRP, Ovid, and Google Scholar till October 2018. Each identified study was evaluated for its quality using the ToxRTool with outcomes such as combined score, weighted score, and reliability category by three independent raters. Correlations of various criteria groups with the combined score were evaluated by Pearson correlation and Kendall rank correlation coefficient. Inter-rater consistency was measured by Cronbach's alpha coefficient. The database searches initially yielded 436 results, of which 391

https://doi.org/10.1016/j.heliyon.2019.e01536
2405-8440/© 2019 Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

Back to Academic Year 2019-2020

Therapeutic Potential of Morin in Ovalbumin-induced Allergic Asthma Via Modulation of SUMF2/IL-13 and BLT2/NF - kB Signalling Pathway

Send Orders for Reprints to reprints@benthamscience.net

Current Molecular Pharmacology, 2019, *12*, 000-000

RESEARCH ARTICLE

Therapeutic Potential of Morin in Ovalbumin-induced Allergic Asthma Via Modulation of SUMF2/IL-13 and BLT2/NF-kB Signaling Pathway

Amit D. Kandhare[1], Zihao Liu[2], Anwesha A. Mukherjee[1] and Subhash L. Bodhankar[1,*]

[1]*Department of Pharmacology, Poona College of Pharmacy, Bharati Vidyapeeth Deemed University, Erandwane, Paud Road, Pune-411 038, India;* [2]*Jiangxi medical college, Nanchang University, 330006, China*

ARTICLE HISTORY

Received: October 03, 2018
Revised: November 30, 2018
Accepted: December 17, 2018

DOI:
10.2174/1874467212666190102105052

Abstract: ***Background***: Allergic asthma is a chronic immune-inflammatory disorder, characterized by airway inflammation and airway hyperresponsiveness (AHR). Morin is a natural flavonoid reported to exhibit inhibitory action against IgE-mediated allergic response.

Aim: To determine the efficacy of murine model of ovalbumin (OVA)-induced AHR inhibition by morin and decipher the molecular mechanism involved.

Materials and Methods: Sprague-Dawley rats were sensitized and challenged with OVA to induce AHR. Rats received treatment with morin (10, 30 and 100 mg/kg, p.o.) for the next 28 days.

Results: Morin (30 and 100 mg/kg) significantly and dose-dependently attenuated ($p < 0.01$ and $p < 0.001$) OVA-induced alterations in pulse oxy and lung function test, increased bronchoalveolar lavage fluid cell counts, elevated total protein and albumin levels in serum, BALF, and lungs, increased serum total and OVA-specific IgE levels and, elevated oxidative stress levels in the lung. RT-PCR analysis revealed that morin treatment (30 and 100 mg/kg) significantly ($p < 0.001$) up-regulated SUMF2 mRNA expression in lungs whereas mRNA expressions of BLT2, NF-κB, and Th2-cytokine (TNF-α, IL-1β, IL-4, IL-6, and IL-13) were down-regulated significantly and dose-dependently ($p < 0.01$ and $p < 0.001$). Also, histologic and ultrastructural studies showed that morin significantly inhibited ($p < 0.001$) OVA-induced perivascular and peribranchial inflammatory infiltration and interstitial fibrosis.

Conclusion: Morin exhibited inhibitory effect against OVA-induced allergic asthma by activation of SUMF2 which impeded IL-13 expression and in turn attenuated Th2-cytokines, BLT2, NF-κB, and IgE levels to ameliorate AHR. Thus, our findings suggested that morin could be considered as a potential alternative therapeutic agent for the management of allergic asthma.

Keywords: Airway hyperresponsiveness, asthma, leukotriene B4 receptor 2, Morin, NF-kB, sulfate-modifying factor 2, Th2 cytokines.

1. INTRODUCTION

Allergic asthma is the most prevalent chronic, immune-inflammatory disorder characterized by airway remodeling, inflammation and, airway hyperresponsiveness (AHR) [1]. According to World Health Organization, approximately 300 million people suffer from asthma worldwide and, the mortality is 180,000 deaths a year [2, 3]. Moreover, the prevalence of asthma is increasing in both developed and developing countries. In a year, the mean cost for asthma management is reckoned at $3,100 per patient (Boonpiyathad *et al*., 2016) which poses a significant economic burden on the healthcare systems calling for urgent implementation of cost-effective strategies.

Numerous evidence has suggested that imbalance in responses of T-helper type 1 (Th1)/T-helper type 2 (Th2) are classical factors for occurrence and progress of asthma [4-7]. It has been associated with an elevation in the production of Th2 cytokine (including interleukin (IL)-4, IL-5, and IL-13) from various Th2 cells (such as eosinophils, B-cells, and mast cells) and decrease in the production of Th1 cytokine [3]. IL-4 and IL-13 play important roles in the activation of B cell for synthesis of immunoglobulin (Ig)-E [8]. The crosslinking of IgE to high-affinity IgE receptors (FcεRI) on the surface of mast cells result in its activation and degranulation. This is followed by the release of an array of inflammatory mediators including histamine, cytokines, and leukotrienes, which lead to immediate bronchoconstriction [9].

*Address correspondence to this author at the Department of Pharmacology, Poona College of Pharmacy, Bharati Vidyapeeth University, Erandwane, Pune, 411 038, India; Tel: + 91-20-24537237 (Ext. 29); Fax: + 91-20-25439386; E-mail: drslbodh@gmail.com

1874-4672/19 $58.00+.00 © 2019 Bentham Science Publishers

Back to Academic Year 2019-2020

EXCLI Journal 2019;18:723-745 – ISSN 1611-2156
Received: June 27, 2019, accepted: August 26, 2019, published: August 29, 2019

Original article:

HESPERIDIN AMELIORATES BLEOMYCIN-INDUCED EXPERIMENTAL PULMONARY FIBROSIS VIA INHIBITION OF TGF-B1/SMAD3/AMPK AND IKBA/NF-KB PATHWAYS

Zheng Zhou[1], Amit D. Kandhare[2], Anwesha A. Kandhare[2], Subhash L. Bodhankar[2]*

[1] Department of Respiratory Medicine, the Second Affiliated Hospital of Zhengzhou University, Zhengzhou City, Henan Province, 450014, China
[2] Department of Pharmacology, Center for Advanced Research in Pharmaceutical Sciences, Bharati Vidyapeeth Deemed University, Poona College of Pharmacy, Pune-411 038, India

* **Corresponding author:** Dr. Subhash L. Bodhankar, Professor and Head, Department of Pharmacology, Center for Advanced Research in Pharmaceutical Sciences, Bharati Vidyapeeth Deemed University, Poona College of Pharmacy, Pune-411 038, India. Phone: +91-20-25437237, Fax: +91-20-25439383; E-mail: drslbodh@gmail.com

http://dx.doi.org/10.17179/excli2019-1094

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0/).

ABSTRACT

Bleomycin (BLM) is a chemotherapeutic agent which is associated with Idiopathic pulmonary fibrosis (IPF) due to its chronic administration. Hesperidin, a bioflavonoid has been reported to possess antioxidant, anti-inflammatory, wound healing, and antiapoptotic potential. To evaluate the therapeutic potential of hesperidin against BLM-induced pulmonary fibrosis and decipher its possible mechanism of action. Intraperitoneal administration of BLM (6 IU/kg) caused induction of IPF in Sprague-Dawley rats. Rats were treated with hesperidin (25, 50, and 100 mg/kg, p.o.) for 28 days, followed by estimation of various parameters in bronchoalveolar lavage fluid (BALF) and lung. Hesperidin (50 and 100 mg/kg) administration significantly ameliorated ($p < 0.05$) alterations induced by BLM in lung index, percent oxygen saturation, serum ALP and LDH levels, BALF differential cell count, and lung function test. Elevated levels of oxido-nitrosative stress, hydroxyproline, and myeloperoxidase levels in BALF and lung were significantly decreased by hesperidin on day 14. Hesperidin significantly inhibited BLM-induced down-regulated lung Nrf2 and HO-1 as well as up-regulated TNF-α, IL-1β, IL-6, collagen-1, TGF-β, and Smad-3 mRNA expressions. Western blot analysis showed that alteration in lung NF-κB, IκBα, AMPK, and PP2C-α protein expressions were ameliorated by hesperidin on day 28. Furthermore, BLM induced histological and ultrastructural aberrations in the lung which were attenuated by hesperidin treatment. Hesperidin alleviates BLM-induced IPF via inhibition of TGF-β1/Smad3/AMPK and IκBα/NF-κB pathways which in turn ameliorate the modulation of oxido-inflammatory markers (Nrf2 and HO-1) and pro-inflammatory markers (TNF-α, IL-1β, and IL-6) to reduce collagen deposition during pulmonary fibrosis.

Keywords: AMPK, bleomycin, hesperidin, IκBα, NF-κB, Nrf2, pulmonary fibrosis, Smad3, TGF-β1

Abbreviations:
Adenosine monophosphate-activated protein kinase (AMPK), Alkaline Phosphatase (ALP), Bleomycin (BLM), Bronchoalveolar Lavage Fluid (BALF), Enhanced Pause (Penh), Expired Volume (EV), Frequency of breathing (f), Glyceraldehyde 3-phosphate dehydrogenase (GAPDH), Glutathione (GSH), Heme oxygenase 1 (HO-1), Hydroxyproline (HP), Idiopathic pulmonary fibrosis (IPF), Interleukins (IL's), Lactate Dehydrogenase (LDH), Malondialdehyde (MDA), Mothers against decapentaplegic homolog-3 (Smad-3), Myeloperoxidase (MPO), Nitric Oxide (NO), Non-phosphorylated AMPK (PP2C-α), Nuclear factor E2-related factor 2 (Nrf2), Nuclear factor-Kappa B (NF-κB), Nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor-alpha (IκBα), Transforming Growth Factor-β (TGF-β), Transmission Electron Microscopy (TEM), Tumor Necrosis Factor-alpha (TNF-α).

Back to Academic Year 2019-2020

Acute and sub-chronic oral toxicity studies of hesperidin isolated from orange peel extract in Sprague Dawley rats

Regulatory Toxicology and Pharmacology 105 (2019) 77–85

Contents lists available at ScienceDirect

Regulatory Toxicology and Pharmacology

journal homepage: www.elsevier.com/locate/yrtph

Yongsheng Li[a], Amit D. Kandhare[b], Anwesha A. Mukherjee[b], Subhash L. Bodhankar[b,*]

[a] Pharmacy Department, Honghui Hospital, Xi'an Jiaotong University, Shaanxi Province, 710054, China
[b] Department of Pharmacology, Poona College of Pharmacy, Bharati Vidyapeeth Deemed University, Erandwane, Paud Road, Pune, 411038, India

ARTICLE INFO

Keywords:
Acute toxicity test
Hesperidin
LD_{50}
OECD
Orange peel
Sub-chronic toxicity tests

ABSTRACT

Citrus sinensis contains glycoside hesperetin-7-rhamnoglucoside (hesperidin) which harbor an array of therapeutic potentials including antioxidant, anticancer, and anti-inflammatory. However, a systematic examination of safety is needed before its utilization. Hence, the present investigation is aimed to evaluate acute and sub-chronic toxicity of hesperidin isolated from the citrus fruit. Hesperidin (73%) was isolated from a methanolic extract of dried peel of the citrus fruit, characterized using FTIR, and standardized by HPLC. Its acute oral toxicity (AOT) and sub-chronic toxicity studies were carried out in Sprague-Dawley rats. Hesperidin (5000 mg/kg) showed 10% mortality in AOT. In sub-chronic toxicity study, hesperidin (250 and 500 mg/kg) did not induce any abnormalities in body weight, food consumption, clinical signs, ophthalmological and neurological observations, urine analysis, hematology, clinical chemistry, organ weights, and gross pathology. However, hesperidin (1000 mg/kg) showed significant ($p < 0.05$) alterations in body and organ weights, hematology, clinical chemistry, and tissue histopathology. To conclude, hesperidin has median lethal dose (LD_{50}) of 4837.5 mg/kg, and Low Observed Adverse Effect Level (LOAEL) at 1000 mg/kg for both male and female Sprague-Dawley rats. Thus, hesperidin isolated from citrus fruit showed a good safety profile in animal study.

1. Introduction

Natural products are an essential source of drugs for management of various diseases. Numerous evidence have suggested that herbal therapeutic moieties are better in terms of safety and efficacy compared to synthetic chemicals (Aniagu et al., 2004; Delaney, 2007; Joshua Allan et al., 2007; Saad et al., 2006). Therefore, with the development of new medical technology, research on isolated compounds from natural and biological resources have garnered much attention. Various natural substances have wide applications in an array of fields including medicine, functional health food, and home remedies (Jakkula et al., 2004; World Health Organization, 2002; Yang et al., 2009).

In developing countries, majority of the population rely on folk medicine obtained from natural sources for the treatment of various diseases. However, these plants contain diverse bioactive compounds that bear the potential to cause an adverse effect (Bent and Ko, 2004). In the Indian subcontinent, 'Ayurveda' has been widely used as an 'alternative medicine' for the past 3000 years or so. Recently, Ayurveda has attained more focus in the field of medicine due to its safety and frequency of success. Moreover, it has been shown to provide excellent clinical results with lower adverse effects than Western medicine (Corns, 2003; Mashour et al., 1998; Yin et al., 2013). But before these traditional medicines can be transformed into modern drugs, significant toxicological information is needed. However, in recent years, the toxicological studies of natural products have rarely been reported. Therefore, there is a need to conduct and document systematic safety studies which examine these substances for their possible toxic effects.

Citrus sinensis (L.) Osbeck, (family: Rutaceae), commonly known as Sweet orange, is a citrus fruit which is largely cultivated in India and various countries around the world. It serves as a major source of vitamin C, potassium, folic acid, and pectin. It is widely consumed as fresh fruit, and juice while the peel is often discarded as waste. According to a report by the United States Department of Agriculture, the global production of this citrus fruit is 82 million tons per year (United States Department of Agriculture, 2018) of which 34% is used for juice production which yields about 44% peel as a by-product (Rafiq et al., 2016). It is to be noted that these peels possess an array of secondary components including polyphenols, vitamins, amino acids, minerals, dietary fibers, essential oils, pectin, flavonoids, vitamin C, and carotenoids. Thus, these have a broad range of activities including

* Corresponding author. Department of Pharmacology, Poona College of Pharmacy, Bharati Vidyapeeth Deemed University, Erandwane, Paud Road, Pune, 411038, Maharashtra, India.
E-mail address: drslbodh@gmail.com (S.L. Bodhankar).

https://doi.org/10.1016/j.yrtph.2019.04.001
Received 1 February 2019; Received in revised form 22 March 2019; Accepted 1 April 2019
Available online 13 April 2019
0273-2300/ © 2019 Elsevier Inc. All rights reserved.

Back to Academic Year 2019-2020

Development and Validation of RP-HPLC and UV Spectrophotometric Absorptivity Method for Simultaneous Estimation of Cyclobenzaprine hydrochloride and Aceclofenac in Pharmaceutical dosage form

ISSN: 2455-2631 © April 2020 IJSDR | Volume 5, Issue 4

Development and Validation of RP-HPLC and UV-Spectrophotometric Absorptivity Method for Simultaneous Estimation of Cyclobenzaprine hydrochloride and Aceclofenac in Pharmaceutical dosage form

Annasaheb S Gaikwad*[1], Monika N Madibone[2], Jeevan R Rajguru[3], Prashant B Jadhav[4]

[1]Department of Quality Assurance Techniques, Pravara Rural College of Pharmacy, Sinnar, Dist- Nashik, M.S. India, 422103,
[2,4]Department of Pharmaceutics, Srinath College of Pharmacy, Bajajnagar, Dist-Aurangabad, M.S. India- 431136
[3]Department of Pharmaceutics, AAEMFS Delight Institute of Pharmacy, Koregaon Bhima, Dist- Pune, M.S. India, 422103,

***Abstract*: The objective of present research is to develop a simple, sensitive, linear, precise and accurate RP-HPLC and UV-Spectrophotometric method for simultaneous estimation of Cyclobenzaprine hydrochloride and Aceclofenac in bulk and tablet formulation as developed and validated. UV-Spectrophotometric method Calibration plot were linear R^2=0.9996 over the concentration range 1-5µg/ml for Cyclobenzaprine hydrochloride, R^2= 0.9995 for the Aceclofenac 13-65µg/ml. And Chromatographic conditions used are stationary phase Inertsil ODS column (250 × 4.6 mm×5µ) (5µm particle size) the mobile phase Methanol: 10mm KH_2PO_4 Buffer (Ph-3) (70:30) and flow rate was maintained 0.9ml/min, detection wavelength was 220nm. The retention times were 4.7min and 2.9 min for Cyclobenzaprine hydrochloride and Aceclofenac respectively. Calibration plot were linear R^2 = 0.9996 over the concentration range 3-15µg/ml for Cyclobenzaprine hydrochloride, R^2 = 0.9994 for the Aceclofenac 40-200µg/ml. No interference from any component of pharmaceutical dosage form was observed. The proposed method has been validated as per ICH guidelines, validation studies revealed that method id specific, rapid, reliable and reproducible. The developed method successfully employed for routine quality control analysis in the combined pharmaceutical dosage form.**

***Keywords*: Cyclobenzaprine hydrochloride and Aceclofenac, UV- Spectrophotometric, RP-HPLC Method.**

I. INTRODUCTION:-

Cyclobenzaprine hydrochloride is a once daily extended release skeletal muscle relaxant which relieves muscle spasm of local origin without interfering with muscle function. The exact mechanism of action is unknown. The IUPAC name is 3-(5H-Dibenzo [A, D] Cyclohepten-5-Ylidene) N, N-Dimethyl-1-Propanamine Absorption: Immediate-release (IR) mean oral bioavailability ranges from 33% to 55. Distribution: About 93% is plasma protein-bound. Metabolism: During first pass through GI tract and liver, drug and metabolites undergo enterohepatic recycling. Cyclobenzaprine is eliminated quite slowly, with an effective half-life of 18 hours (range 8-37 hours; n=18); plasma clearance is 0.7 L/min. Metabolites excreted in the urine are likely water-soluble glucuronide conjugates.



Cyclobenzaprine Hydrochloride

Aceclofenac is a non-steroidal agent with marked anti-inflammatory and analgesic properties. The mode of action of aceclofenac is largely based on the inhibition to prostaglandin synthesis. Aceclofenac is a potent inhibitor of the enzyme cyclo-oxygenase, which is involved in the production of prostaglandins. The IUPAC name is [(2, 6-dichlorophenyl) amino] phenylacetoxyacetic acid. After oral administration, aceclofenac is rapidly absorbed and the bioavailability is almost 100%. Peak plasma concentrations are reached approximately1.25 to 3 hours. Aceclofenac is highly protein-bound (> 99.7%). Aceclofenac penetrates into the synovial fluid where the concentrations reach approximately 60%of those in plasma. Aceclofenac is probably metabolized via CYP2C9 to the main metabolite 4-hydroxyaceclofenac. The mean plasma elimination half-life is 4-4.3 hours.Approximately two-thirds of the administered dose is excreted via the urine, analysis conjugated hydroxymetabolites.

Back to Academic Year 2019-2020

A Review - Corona Virus (Covid-19)

A REVIEW - CORONA VIRUS (COVID-19)

Jeevan R. Rajguru, Ms. Snehal D. Jadhav, Annasaheb S. Gaikwad, Dr. Sampat D. Navale

Prof. Jeevan R. Rajguru (Department of Pharmaceutics, AAEMF'S Delight Institute of Pharmacy, Koregaon Bhima, Dist-Pune M.S. India)

Ms. Snehal D. Jadhav (Department of Quality Assurance, AAEMF'S Delight College of Pharmacy, Koregaon Bhima, Dist-Pune M.S. India)

Annasaheb S. Gaikwad (Department of Quality Assurance, Pravara Rural Education Society's College Of Pharmacy, Sinnar Dist-Nashik M.S. India)

Dr. Sampat D. Navale (Department of Pharmacognosy, AAEMF'S Delight Institute of Pharmacy, Koregaon Bhima, Dist-Pune M.S. India)

***Abstract*: An acute respiratory disease, caused by a novel coronavirus (SARS-CoV-2, previously known as 2019-nCoV), the coronavirus disease 2019 (COVID-19) has spread throughout China and received worldwide attention. On 30 January 2020, World Health Organization (WHO) officially declared the COVID-19 epidemic as a public health emergency of international concern. The emergence of SARS-CoV-2, since the severe acute respiratory syndrome coronavirus (SARS-CoV) in 2002 and Middle East respiratory syndrome coronavirus (MERS-CoV) in 2012, marked the third introduction of a highly pathogenic and large-scale epidemic coronavirus into the human population in the twenty-first century. As of 1 March 2020, a total of 87,137 confirmed cases globally, 79,968 confirmed in China and 7169 outside of China, with 2977 deaths (3.4%) had been reported by WHO. Meanwhile, several independent research groups have identified that SARS-CoV-2 belongs to β-coronavirus, with highly identical genome to bat coronavirus, pointing to bat as the natural host. The novel coronavirus uses the same receptor, angiotensin-converting enzyme 2 (ACE2) as that for SARS-CoV, and mainly spreads through the respiratory tract. Importantly, increasingly evidence showed sustained human-to-human transmission, along with many exported cases across the globe. The clinical symptoms of COVID-19 patients include fever, cough, fatigue and a small population of patients appeared gastrointestinal infection symptoms. The elderly and people with underlying diseases are susceptible to infection and prone to serious outcomes, which may be associated with acute respiratory distress syndrome (ARDS) and cytokine storm.Currently, there are few specific antiviral strategies, but several potent candidates of antivirals and repurposed drugs are under urgent investigation. In this review, we summarized the latest research progress of the epidemiology, pathogenesis, and clinical characteristics of COVID-19, and discussed the current treatment and scientific advancements to combat the epidemic novel coronavirus.**

***Keywords*: COVID-19, β-coronavirus, SARS-CoV, canine coronavirus (CCoV).**

BACKGROUND:

In December 2019, a cluster of pneumonia cases, caused by a newly identified β-coronavirus, occurred in Wuhan, China. This coronavirus, was initially named as the 2019- novel coronavirus (2019-nCoV) on 12 January 2020 by World Health Organization (WHO). WHO officially named the disease as coronavirus disease 2019 (COVID- 19) and Coronavirus Study Group (CSG) of the International Committee proposed to name the new coronavirus as SARS-CoV-2, both issued on 11 February 2020. The Chinese scientists rapidly isolated a SARS-CoV-2 from a patient within a short time on 7 January 2020 and came out to genome sequencing of the SARS-CoV-2 [6]. As of 29 March 2020, a total of 81,439 cases of COVID-19 have been confirmed in mainland China including 3,300 deaths [7]. Studies estimated the basic reproduction number (R0) of SARS-CoV-2 to be around 2.2 [8], or even more (range from 1.4 to 6.5) [9], and familial clusters of pneumonia [10] outbreaks add to evidence of the epidemic COVID-19 steadily growing by human-to-human transmission.

INTRODUCTION:

Coronavirus disease 2019 (COVID-19) is a respiratory illness that can spread from person to person. The virus that causes COVID-19 is a novel coronavirus that was first identified during an investigation into an outbreak in Wuhan, China. [1]

CORONAVIRUSES:

Coronaviruses are a group of related viruses that cause diseases in mammals and birds. In humans, coronaviruses cause respiratory tract infections that can be mild, such as some cases of the common cold (among other possible causes, predominantly rhinoviruses), and others that can be lethal, such as SARS, MERS, and COVID-19. Symptoms in other species vary: in chickens, they cause an upper respiratory tract disease, while in cows and pigs they cause diarrhea. There are yet to be vaccines or antiviral drugs to prevent or treat human coronavirus infections.[10] Coronaviruses constitute the subfamily ***Orthocoronavirinae***, in the family *Coronaviridae*, order *Nidovirales*, and realm *Riboviria*. They are enveloped viruses with a positive-sense single-stranded RNA genome and a nucleocapsid of helical symmetry. The genome size of coronaviruses ranges from approximately 27 to 34 kilobases, the largest among known RNA viruses. The name *coronavirus* is derived from the Latin *corona*, meaning "crown" or "halo", which refers to the characteristic appearance reminiscent of a crown or

Back to Academic Year 2019-2020

Targeted Drug Delivery – From Magic Bullet to Nanomedicine

Journal of Xidian University DOI:10.37896/jxu17.8/128 ISSN No:1001-2400

Targeted Drug Delivery – From Magic Bullet to Nanomedicine

Annasaheb S Gaikwad[1], Mayur R Waje[2], Vaishanavi S Tile[3], Pallavi Patharkar[4]

[1]Jaihind College of Pharmacy, Vadgaon Sahani, Pune (MS), India.

[3] Asian Institute of Pharmacy, Pathardi Phata, Nashik, (MS), India

[2,4] Mangaldeep Institute of Pharmacy, Nipani, Chap. Sambhajinagar (MS), India.

Corresponding Author;

Annasaheb S Gaikwad

Abstract:

Targeted medication delivery has in fact substantially improved from conventional techniques to more complex strategies like the magic pill and nanomedicine. The idea of delivering a medicine to a specific target in the body while limiting its effects on healthy tissues is known as the "magic bullet" theory. This idea can target therapeutic effects only to the desired site while differentiating between healthy and diseased cells. Targeted drug distribution is one of the applications of nanotechnology in nanomedicine. Controlled drug release provided by nanoparticles ensures long-lasting therapeutic levels and lowers systemic toxicity. The blood-brain barrier is just one example of a biological barrier that they can get through, making it possible to deliver drugs to previously unreachable places. The review article highlights targeted drug delivery's uses in treating cancer, neurological disorders, and cardiovascular ailments, among other diseases. It emphasizes how increasing therapeutic results and reducing off-target consequences might improve illness treatment through targeted medication delivery strategies.

Keywords: Magic Mullet, Blood Brain Barrier, Inaccessible Area, Revolutionize Disease, Off-Target Effect.

Back to Academic Year 2019-2020

Cancer Chemotherapy with Peptides & Peptidomimetics drug & Peptide based Vaccine

International Research Journal of Pharmacy and Medical Sciences
ISSN (Online): 2581-3277

Cancer Chemotherapy with Peptides and Peptidomimetics Drug and Peptide Based-Vaccines

Jeevan R. Rajguru*[1], Sonali A. Nagare[2], Ashish A. Gawai[3], Amol G. Jadhao[4], Mrunal K. Shirsat[5]
[1, 4]Master of Pharmacy in Pharmaceutics at Anuradha College of Pharmacy, Chikhli, Dist-Buldana M.S, India
[2]Department of Pharmaceutical Chemistry Loknete Shri Dadapatil Pharate College of pharmacy, Mandavgan Pharata, Shirur Pune, M.S, India
[3]Associate Professor, Department of Pharmaceutical Chemistry at Anuradha College of pharmacy, Chikhli, Dist- Buldana M.S, India
[5]Department of Pharmacognosy and Principal of Loknete Shri Dadapatil Pharate College of pharmacy, Mandavgan Pharata, Shirur Pune, M.S, India
Corresponding Author Email ID: jeevanrajguru97@gmail.com
Co-Author Email ID: sonalinagare93@gmail.com

Abstract— *A summary of the current status of the application of peptidomimetics in cancer therapeutics as an alternative to peptide drugs is provided. Only compounds that are used in therapy or at least under clinical trials are discussed, using inhibitors of farnesyltransferase, proteasome and matrix metalloproteinases as examples. The design and synthesis of peptidomimetics are most important because of the dominant position peptide and protein-protein interactions play in molecular recognition and signalling, especially in living systems. The design of peptidomimetics can be viewed from several different perspectives and peptidomimetics can be categorized in a number of different ways. Study of the vast literature would suggest that medicinal and organic chemists, who deal with peptide mimics, utilize these methods in many different ways. Conventional methods used to treat cancer, from non-specific chemotherapy to modern molecularly targeted drugs have generated limited results due to the complexity of the disease as well as lack of molecular classes that can be developed into treatments rapidly, easily and economically. Peptidomimetics that are easy to synthesize and optimize and has been studied in different oncology applications as great biologically amenable compounds and can be considered as a promising alternative molecular class for anticancer drug developments. Peptide can be utilized directly as a cytotoxic agent through various mechanisms or can act as a carrier of cytotoxic agents and radioisotopes by specifically targeting cancer cells. Peptide-based hormonal therapy has been extensively studied and utilized for the treatment of breast and prostate cancers. Tremendous amount of clinical data is currently available attesting to the efficiency of peptide-based cancer vaccines.*

Keywords— *Farnesyltransferase inhibitors, Ras, Metal-loproteinase inhibitors, Angiogenesis, Proteasome inhibitors.*

I. Introduction

Mortality from cancer is about to surpass that from cardiovascular diseases in near future. About 7 million people die from cancer-related cases per year, and it is estimated that there will be more than 16 million new cancer cases every year by 2020 [1, 2]. Cancer is characterized by uncontrolled division of cells and the ability of these cells to invade other tissues leading to the formation of tumor mass, vascularization, and metastasis (spread of cancer to other parts of the body) [3]. Though angiogenesis (growth of new blood vessels from pre-existing vessels) is a normal and vital process in growth and development, it is also a fundamental step in the transition of b tumors from adormant state to a malignant one [4]. Chemotherapy is one of the major approaches to treat cancer by delivering a cytotoxic agent to the cancer cells. The main problem with the conventional chemotherapy is the inability to deliver the correct amount of drug directly to cancer cells without affecting normal cells [5]. Drug resistance, altered bio distribution, biotransformation, and drug clearance are also common problems [5]. Targeted Mortality from cancer is about to surpass that from cardiovascular diseases in near future. About 7 million people die from cancer-related cases per year, and it is estimated that there will be more than 16 million new cancer cases every year by 2020 [1, 2]. Cancer is characterized by uncontrolled division of cells and the ability of these cells to invade other tissues leading to the formation of tumor mass, vascularization, and metastasis (spread of cancer to other parts of the body) [3]. Though angiogenesis (growth of new blood vessels from pre-existing vessels) is a normal and vital process in growth and development, it is also a fundamental step in the transition of tumors from a dormant state to a malignant one [4]. The "biologics" treatment option against cancer includes the use of proteins, monoclonal antibodies, and peptides. The monoclonal antibodies (mAbs) and large protein ligands have two major limitations compared to peptides: poor delivery to tumors due to their large size and dose-limiting toxicity to the liver and bone marrow due to nonspecific uptake into the reticulo endothelial system. The use of such macromolecules has therefore been restricted to either vascular targets present on the luminal side of tumor vessel endothelium or hematological malignancies [6–11]. Peptides possess many advantages such as small size, ease of synthesis and modification, tumor penetrating ability, and good biocompatibility [12, 13]. Peptide degradation by proteolysis can be prevented by chemical modifications such as incorporation of D-amino acids or cyclization [14]. Over the years peptides have been evolved as promising therapeutic agents in the treatment of cancer, diabetes, and cardiovascular diseases, and application of peptides in a variety of other therapeutic areas is growing rapidly. Currently there are about

Jeevan R. Rajguru, Sonali A. Nagare, Ashish A. Gawai, Amol G. Jadhao, and Mrunal K. Shirsat, "Cancer Chemotherapy with Peptides and Peptidomimetics Drug and Peptide Based-Vaccines," *International Research Journal of Pharmacy and Medical Sciences (IRJPMS)*, Volume 2, Issue 2, pp. 21-30, 2019.

A study to assess the knowledge regarding hand hygi [Back to Academic Year 2019-2020] g children in selected schools of Pune city

International Journal of Applied Research 2020; 6(12): 27-29

ISSN Print: 2394-7500
ISSN Online: 2394-5869
Impact Factor: 8.4
IJAR 2020; 6(12): 27-29
www.allresearchjournal.com
Received: 15-10-2020
Accepted: 18-11-2020

Sneha Sankpal Desai
Bharati Vidyapeeth (Deemed to be) University College of Nursing, Pune, Maharashtra, India

Sampurna Kar
Bharati Vidyapeeth (Deemed to be) University College of Nursing, Pune, Maharashtra, India

Shital Kshirsagar
Bharati Vidyapeeth (Deemed to be) University College of Nursing, Pune, Maharashtra, India

Shinthoi Chanu
Bharati Vidyapeeth (Deemed to be) University College of Nursing, Pune, Maharashtra, India

Prajakta Bansode
Bharati Vidyapeeth (Deemed to be) University College of Nursing, Pune, Maharashtra, India

Kunal Pandya
Bharati Vidyapeeth (Deemed to be) University College of Nursing, Pune, Maharashtra, India

Corresponding Author:
Sneha Sankpal Desai
Bharati Vidyapeeth (Deemed to be) University College of Nursing, Pune, Maharashtra, India

A study to assess the knowledge regarding hand hygiene practice among secondary school going children in selected schools of Pune city

Sneha Sankpal Desai, Sampurna Kar, Shital Kshirsagar, Shinthoi Chanu, Prajakta Bansode and Kunal Pandya

Abstract
Hand hygiene practice is the single most effective way to prevent the spread of germs. Good hand hygiene reduces the risk of flu, food poisoning and health care associated infections passed from person to person. Although maintaining hand hygiene practice is very important for all of us but it is more important to teach the children about hand hygiene practice because they are at increased risk for getting infection as they are grouped together and their immunity system is not fully developed.
A cross-sectional survey was conducted in Pune (West of India). 150 samples were collected from schools for the study. The questionnaire consist of 30 items and two section (demographic information and knowledge score about hand hygiene practice).
Demographic data of the sample was statistically analysed by using frequency and percentage it was observed that (2.67%) children are having poor knowledge, (8%) children are having average knowledge, (21.33%) children are having good knowledge and (68%) children are having excellent knowledge of hand hygiene practice.
There is no association between knowledge and selected demographic variables (Age, Gender, Mother's education, Father's education, Mother's occupation, Father's occupation, from where did children learn about hand hygiene practice? How many times a day children wash their hands?)

Keywords: Hand hygiene, risk of flu, Pune

Introduction
Hand washing or hand hygiene is the act of cleaning one's hands with or without the use of water or another liquid, or with use of soap for the purpose of removing soil, dirt or micro-organisms. Hand washing is the single most effective way to prevent the spread of germs or micro-organisms which prevent communicable diseases. It is easy to learn how to control the spread of infection by washing the germs away.
The aim of hand washing is to remove micro-organisms from the hands preventing their potential transfer. It is known that organism survives and multiply on human hands, creating the opportunity to infect the others or the host.
Hand washing reduces the number of transient organisms on the skin surface. Although hands cannot be sterilized, most transient organisms can be removed by thirty seconds of proper scrubbing with soap and water.

Need of the study
The significance of hand washing in patients care was conceptualized in the early 19th century. Labarraque provided the first evidence that hand decontamination can markedly reduce the incidence of puerperal fever and maternal mortality.
Semmelweis worked in the Great hospital in Vienna in the 1840s. There was two maternity clinics in the hospital. The first clinic was attended by medical students, who moved straight from autopsy rooms to the delivery suit and had an average maternal mortality rate due to puerperal fever of about 10 percent. The second clinic, attended by midwives had a maternal mortality of only 2 percent. He instituted a policy of washing hands with chlorinated lime for those leaving the autopsy room, following which the rate of maternal mortality dropped ten-folds, comparable to the second clinic.

Back to Academic Year 2019-2020

Academic Year 2

Sr. no	Title of the paper	Name of Staff	Department	Name of the journal	ISSN number	Document	Link

1	Drug-Excipient Interactions Study	Dr. H. V. Kamble	Pharmacology	World Journal of Pharmacy and Pharmaceutical Sciences	ISSN 2277–7105	View Document	https://www.ijnrd.org/viewpaperforall?paper=IJNRD2302162
2	Elucidation of Molecular Mechanism Involved in Nephroprotective Potential of Naringin in Ethylene Glycol-Induced Urolithiasis in Experimental Uninephrectomized Hypertensive Rats	Dr. A. D. Kandhare	Pharmacology	Latin American Journal of Pharmacy - 39 (5): 991-9 (2020)	ISSN 2362-3853	View Document	https://www.google.com/search?q=Elucidation+of+Molecular+Mechanism+Involved+in+Nephroprotective+Potential+of+Naringin+in+Ethylene+Glycol-Induced+Urolithiasis+in+Experimental+Uninephrectomized+Hypertensive+Rats+a.d.kandhare&rlz=1C1VDKB_en-GBIN1070IN1070&oq=Elucidation+of+Molecular+Mechanism+Involved+in+Nephroprotective+Potential+of+Naringin+in+Ethylene+Glycol-Induced+Urolithiasis+in+Experimental+Uninephrectomized+Hypertensive+Rats+a.d.kandhare&gs_lcrp=EgZjaHJvbWUyBggAEEUYOdIBCTQ1ODIxajBqNKgCALACAA&sourceid=chrome&ie=UTF-8
3	Herbal Medicines and Screening Models for Hepatoprotective Agents	Ms. P. P. Kothari	Pharmacology	World Journal of Pharmaceutical Research	ISSN 2277–7105	View Document	https://www.google.com/search?q=Herbal+Medicines+and+Screening+Models+for+Hepatoprotective+Agents+research+paper+by+pratiksha+kothari&rlz=1C1VDKB_en-GBIN1070IN1070&oq=Herbal+Medicines+and+Screening+Models+for

							+Hepatoprotective+Agents+research+paper+by+pratiksha+kothari&gslcrp=EgZjaHJvbWUyBggAEEUYOTIGCAEQRRhAMgYIAhBFGDvSAQkyMDkxNWowajeoAgCwAgA&sourceid=chrome&ie=UTF-8
4	Herbal Medicines and Screening Models for Hepatoprotective Agents	Dr. S. A. Waghmare	Pharmacology	World Journal of Pharmaceutical Research	ISSN 2277–7105	View Document	https://www.google.com/search?q=Herbal+Medicines+and+Screening+Models+for+Hepatoprotective+Agents+research+paper+by+pratiksha+kothari&rlz=1C1VDKB_en-GBIN1070IN1070&oq=Herbal+Medicines+and+Screening+Models+for+Hepatoprotective+Agents+research+paper+by+pratiksha+kothari&gslcrp=EgZjaHJvbWUyBggAEEUYOTIGCAEQRRhAMgYIAhBFGDvSAQkyMDkxNWowajeoAgCwAgA&sourceid=chrome&ie=UTF-8
5	Herbal Medicines and Screening Models for Hepatoprotective Agents	Dr. H. V. Kamble	Pharmacology	World Journal of Pharmaceutical Research	ISSN 2277–7105	View Document	https://www.google.com/search?q=Herbal+Medicines+and+Screening+Models+for+Hepatoprotective+Agents+research+paper+by+pratiksha+kothari&rlz=1C1VDKB_en-GBIN1070IN1070&oq=Herbal+Medicines+and+Screening+Models+for+Hepatoprotective+Agents+research+paper+by

							+pratiksha+kothari&gslcrp=EgZjaHJvbWUyBggAEEUYOTIGCAEQRRhAMgYIAhBFGDvSAQkyMDkxNWowajeoAgCwAgA&sourceid=chrome&ie=UTF-8
8	Development and Validation of RP-HPLC and UV-Spectrophotometric Method for Simultaneous Estimation of Teneligliptin Hydrobromide Hydrate and Metformin Hydrochloride in Pharmaceutical dosage form	Mr. A. S. Gaikwad	Pharmacology	International Journal of Creative Research Thoughts	ISSN: 2320-2882	View Document	https://www.researchgate.net/publication/374813445_Development_and_Validation_of_RP-HPLC_and_UV-Spectrophotometric_Method_for_Simultaneous_Estimation_of_Teneligliptin_Hydrobromide_Hydrate_and_Metformin_Hydrochloride_in_Pharmaceutical_dosage_form
9	High Performance Liquid Chromatography-A Review	Mr. A. S. Gaikwad	Pharmacology	World Journal Of Pharmacy And Pharmaceutical Sciences	ISSN 2278 – 4357	View Document	https://scholar.google.co.in/scholar?q=High+Performance+Liquid+Chromatography-A+Review+by+a.s.gaikwad&hl=en&as_sdt=0&as_vis=1&oi=scholart

Back to Academic Year 2020-21

10	Development and Validation of RP-HPLC and UV Spectrophotometric Absorptivity Method for Simultaneous Estimation of Cyclobenzaprine hydrochloride and Aceclofenac in Pharmaceutical dosage form	Mr. A. S. Gaikwad	Pharmacology	International Journal of Scientific Development and Research	ISSN: 2455-2631	View Document	https://www.researchgate.net/publication/374813338_Development_and_Validation_of_RP-HPLC_and_UV-_Spectrophotometric_Absorptivity_Method_for_Simultaneous_Estimation_of_Cyclobenzaprine_hydrochloride_and_Aceclofenac_in_Pharmaceutical_dosage_form
11	Synthesis, Characterization and Antidiabetic Evaluation of Sulfonamide in Corporated with 1,3,4-Oxadiazole Derivatives	Ms. M. T. Salve	Pharmacology	Indian Journal of Pharmaceutical Education and Research	Vol 55 Issue 4 1145	View Document	https://www.google.com/search?q=Synthesis%2C+Characterization+and+Antidiabetic+Evaluation+of+Sulfonamide+in+Corporated+with%0D%0A1%2C3%2C4-Oxadiazole+Derivatives%0D%0A+research+paper+by+m.t.salve&sca_esv=9c9edb63dbab71a4&sca_upv=1&rlz=1C1VDKB_en-GBIN1070IN1070&sxsrf=ACQVn0_YILL1hUj1r65PNTu3cPUJlg_93A%3A1713677958243&ei=hqYkZr-zDq3XseMP_pmd0A4&ved=0ahUKEwi_jtfvy9KFAxWta2wGHf5MB-oQ4dUDCBA&uact=5&o

							q=Synthesis%2C+Characterization+and+Antidiabetic+Evaluation+of+Sulfonamide+in+Corporated+with%0D%0A1%2C3%2C4-Oxadiazole+Derivatives%0D%0A+research+paper+by+m.t.salve&gs_lp=Egxnd3Mtd2l6LXNlcnAikwFTeW50aGVzaXMsIENoYXJhY3Rlcml6YXRpb24gYW5kIEFudGlkaWFiZXRpYyBFdmFsdWF0aW9uIG9mIFN1bGZvbmFtaWRlIGluIENvcnBvcmF0ZWQgd2l0aAoxLDMsNC1PeGFkaWF6b2xlIERlcml2YXRpdmVzCiByZXNlYXJjaCBwYXBlciBieSBtLnQuIHNhbHZlMgAkAEAmAEAoAEAqgEAuAEDyAEA-AEC-AEBmAIAoAIAmAMAkgcAoAcA&sclient=gws-wiz-serp
13	Synthesis, Characterization, and Evaluation of in vitro Antidiabetic Activity of Novel Pyrrolidine Sulphonamide Derivative	Ms. M. T. Salve	Pharmacology Pharmaceutical Chemistry	International Journal of Pharmaceutical Investigation	Vol 11, Issue 4	View Document	https://www.google.com/search?q=Synthesis%2C+Characterization%2C+and+Evaluation+of+in+vitro%0D%0AAntidiabetic+Activity+of+Novel+Pyrrolidine+Sulphonamide%0D%0ADerivative%0D%0A+research+paper+by+m.t.salve&sca_esv=9c9edb63dbab71a4&sca_upv=1&rlz=1C1VDKB_en-GBIN1070IN1070&sxsrf=ACQVn09IiPSXqpa0597xagN66VXIm4RFtw%3A1713678080795&ei=AKckZsqRMLaNseM

							PkbGW6AQ&ved=0ahUKEwjKkY-qzNKFAxW2RmwGHZGYBU0Q4dUDCBA&uact=5&oq=Synthesis%2C+Characterization%2C+and+Evaluation+of+in+vitro%0D%0AAntidiabetic+Activity+of+Novel+Pyrrolidine+Sulphonamide%0D%0ADerivative%0D%0A+research+paper+by+m.t.salve&gs_lp=Egxnd3Mtd2l6LXNlcnAilwFTeW50aGVzaXMsIENoYXJhY3Rlcml6YXRpb24sIGFuZCBFdmFsdWF0aW9uIG9mIGluIHZpdHJvCkFudGlkaWFiZXRpYyBBY3Rpdml0eSBvZiBOb3ZlbCBQeXJyb2xpZGluZSBTdWxwaG9uYW1pZGUKRGVyaXZhdGl2ZQogcmVzZWFyY2gg cGFwZXIgYnkgbS50LnNhbHZlMggQABiABBiiBEjWGFAAWMgQcAB4AJABAJgBpwGgAacBqgEDMC4xuAEDyAEA-AEBwAECmAIBoAKuAZgDAJIHAzAuMaAH2wI&sclient=gws-wiz-serp
14	Quality By Design Approaches To Analytical Method Development: A Review	Ms. S. S. Shingne & Mr. A. E Bhujbal		World Journal of Pharmaceutical Research World Journal of Pharmaceutical Research	ISSN 2277–7105	View Document	https://wjpr.net/public/abstract_show/18121

15	Genotoxicity: Malignancy, Damage To DNA	Ms. S. S. Shingne & Mr. A. E Bhujbal		World Journal of Pharmaceutical Research	ISSN 2277–7105	View Document	https://wjpr.net/abstract_show/18131
16	Isolation and characterization of phytoconstituents from petroleum ether extract of Momordica cochinchinensis fruits	Dr. M. R. Agrawal	Pharmacognosy	Journal of Pharmacognosy and Phytochemistry	E-ISSN: 2278-4136 P-ISSN: 2349-8234	View Document	https://scholar.google.co.in/scholar?q=Isolation+and+characterization+of+phytoconstituents+from+petroleum+ether+extract+of+Momordica+cochinchinensis+fruits+research+paper+by+m.+r.+agrawal&hl=en&as_sdt=0&as_vis=1&oi=scholart
17	Analgesic activity of Momordica cochinchinensis and Momordica balsamina fruit extracts	Dr. M. R. Agrawal	Pharmacognosy	International Journal of Green and Herbal Chemistry	Vol.7	View Document	https://scholar.google.co.in/scholar?q=Analgesic+activity+of+Momordica%0D%0Acochinchinensis+and+Momordica+balsamina+fruit+extracts%0D%0A+research+paper+by+m.+r.+agrawal&hl=en&as_sdt=0&as_vis=1&oi=scholart
18	Isolation, Characterization of Phytoconstituents from petroleum ether extract of Momordica balsamina fruits and its	Dr. M. R. Agrawal	Pharmacognosy	Research Journal of Pharmacognosy and Phytochemistry	ISSN 0975-4385	View Document	https://rjpponline.org/AbstractView.aspx?PID=2020-12-3-6

	quantitative analysis by HPTLC						
19	Quality By Design Approaches To Analytical Method Development: A Review	Mr. S. A. Waghmare	Pharmaceutical Chemistry	World Journal of Pharmaceutical Research World Journal of Pharmaceutical Research	ISSN 2277–7105	View Document	https://wjpr.net/public/abstract_show/18121
20	Genotoxicity: Malignancy, Damage To DNA	**Dr. H. V. Kamble**	Pharmacology	World Journal of Pharmaceutical Research	ISSN 2277–7105	View Document	https://wjpr.net/abstract_show/18131
21	Genotoxicity: Malignancy, Damage To DNA	Mr. S. A. Waghmare	Pharmaceutical Chemistry	World Journal of Pharmaceutical Research	ISSN 2277–7105	View Document	https://wjpr.net/abstract_show/18131

Back to Academic Year 2020-21

Drug- Excipient Interactions Study

WhatsApp editor@ijnrd.org

INTERNATIONAL JOURNAL OF NOVEL RESEARCH AND DEVELOPMENT
International Peer Reviewed & Refereed Journals, Open Access Journal
ISSN Approved Journal No: 2456-4184 | Impact factor: 8.76 | ESTD Year: 2016
Scholarly open access journals, Peer-reviewed, and Refereed Journals, Impact factor 8.76 (Calculate by google scholar and Semantic Scholar | AI-Powered Research Tool) , Multidisciplinary, Monthly, Indexing in all major database & Metadata, Citation Generator, Digital Object Identifier(DOI)

Submit Paper Online | Login to Author Home

Home IJNRD Editioral Board For Author Current Issue Archive Conference All Policy FAQ Contact Us

Call For Paper
Issue: March 2024
Volume 9 | Issue 3
Submit Paper Online
Last date: 31-Mar-2024
Review Result and Publication of Paper within : 2-3 days
Click Here For more Details

For Authors
Submit Paper Online
Publication Guidelines
Publication Charges
Pay Publication Fees Online
Track Paper
Research Area
All Policy

Forms / Download
Copyright Form
Undertaking Form
Paper Format
Sample Certificate
Sample Publication Letter
Sample Hard Copy of Journal

Published Issue Details
Current Issue
Archive
Conference Proposal
Recent Conference Details

Editorial Board

Published Paper Details

Paper Title:	**A REVIEW: DRUG- EXCIPIENT INTERACTIONS STUDY**
Authors Name:	Mr. Shiddesh D. Gawari , Ms.Sonali B. Jadhav , Dr. Hemant V. Kambale , Dr. Vivek M. Satpute , Santosh A. Waghmare
Download E-Certificate:	Download
Author Reg. ID:	IJNRD_187576
Published Paper Id:	IJNRD2302162
Published In:	Volume 8 Issue 2, February-2023
DOI:	
Abstract:	Excipients are included in dosage forms to aid manufacture, administration or absorption. Although considered pharmacologically inert, excipients can initiate, propagate or participate in chemical or physical interactions with drug compounds, which may compromise the effectiveness of a medication. Exicipients are not exquisitely pure. Even for the most commonly used excipients, it is necessary to understand the context of their manufacture in order to identify potential active pharmaceutical ingredients interactions with trace components. Chemical interactions can lead to degradation of the active ingredient, thereby reducing the amount available for therapeutic effect. Physical interactions can affect rate of dissolution, uniformity of dose or ease of administration. Understanding the chemical and physical nature of excipients, the impurities or residues associated with them and how they may interact with other materials, or with each other, forewarns the pharmaceutical technologist of possibilities for undesirable developments. The potential physical and chemical interactions between drugs and excipients can affect the chemical, physical, therapeutical properties and stability of the dosage form. The present review contains a basic mode of drug degradation, mechanism of drug- excipient interaction like physical, chemical and biopharmaceutical. Different Thermal and Non-thermal method of analysis, Tools and software for incompatibility is also discussed. Once the type of interaction is determined we can take further steps to improve the stability of drug and dosage form. From review, we conclude that consequent use of thermal and non-thermal method provide data for drug- excipient interaction which can further help in selection of excipient for the development of stable dosage form.
Keywords:	Excipient, Drug, Interaction, Physical, Chemical
Cite Article:	**"A REVIEW: DRUG- EXCIPIENT INTERACTIONS STUDY"**, International Journal of Novel Research and Development (www.ijnrd.org), ISSN:2456-4184, Vol.8, Issue 2, page no.b641-b651, February-2023, Available :http://www.ijnrd.org/papers/IJNRD2302162.pdf
Downloads:	000118744
ISSN:	2456-4184 | IMPACT FACTOR: 8.76 Calculated By Google Scholar| ESTD YEAR: 2016 An International Scholarly Open Access Journal, Peer-Reviewed, Refereed Journal

Submit Paper Online
Check Paper Status
Call for Paper
Join As reviewer
All Policy

ISSN Details
INTERNATIONAL STANDARD SERIAL NUMBER
ISSN 2456-4184
9 772456 418401
OPEN ACCESS

ISSN: 2456-4184
Impact Factor: 8.76 and ISSN APPROVED Journal Starting Year (ESTD) : 2016

DOI (A digital object identifier)
Providing A digital object identifier by DOI
How to Get DOI? DOI

Conference
Conference Proposal
Recent Conference Details

Open Access License Policy
This work is licensed under a Creative Commons Atribution-

Back to Academic Year 2020-21

Herbal Medicines and Screening Models for Hepatoprotective Agents

WORLD JOURNAL OF PHARMACEUTICAL RESEARCH
SJIF Impact Factor 8.084
Volume 10, Issue 8, 1373-1386. Review Article ISSN 2277– 7105

A REVIEW: HERBAL MEDICINES AND SCREENING MODELS FOR HEPATOPROTECTIVE AGENTS

Pratiksha Kothari*, Ashvini Andhale and Santosh Waghmare

Loknete Shri Dadasaheb Pharate Patil College of Pharmacy, Mandavgoan Pharata, Tal-Shirur, Dist.-Pune. Maharashtra.

Article Received on
21 May 2021,
Revised on 11 June 2021,
Accepted on 01 July 2021
DOI: 10.20959/wjpr20218-20983

***Corresponding Author**
Pratiksha Kothari
Loknete Shri Dadasaheb Pharate Patil College of Pharmacy, Mandavgoan Pharata, Tal- Shirur, Dist.-Pune. Maharashtra.

ABSTRACT

Liver is the largest gland present in human body. The function of this vital organ includes storage of glycogen, vitamins and minerals, enzyme activation, metabolism of carbohydrate, proteins and detoxification as well as purification process. Liver regulates chemical levels in blood and forms excretory material that is bile. All the blood leaving the stomach and intestine passes through liver. Chronic alcohol consumption, exposure to toxic chemicals like paracetamol, tetracycline, anti-TB drugs, NSAIDS, chemotherapeutic agents damage the hepatocytes. Liver diseases are major health and challenging to health care professions and pharmaceutical industry. Therefore there is need to study the hepatoprotective agents. Modern medicine have provided many medicaments that cure liver diseases but compared to this herbal medicine is preferred due to its cost effectiveness as well as minimal side effects and safe approach. The aim of this review is compiling information of different medicinal plants with their hepatoprotective activity on various models of hepatotoxicity.

KEYWORDS: Liver, Hepatotoxic agents, Medicinal herbs as hepatoprotective agents, Screening models.

INTRODUCTION

The Liver is the largest gland in a human body, situated in the right side of upper abdominal cavity. The cells of the liver called hepatocytes plays vital functions like;

1. Synthesis of proteins, biles,
2. Stores glycogen, vitamins, iron,
3. Metabolizes toxic chemicals and drugs

Back to Academic Year 2020-21

Elucidation of Molecular Mechanism Involved in Nephroprotective Potential of Naringin in Ethylene Glycol-Induced Urolithiasis in Experimental Uninephrectomized Hypertensive Rats

Latin American Journal of Pharmacy
(formerly *Acta Farmacéutica Bonaerense*)
Lat. Am. J. Pharm. **39** (5): 991-9 (2020)

Received: December 31, 2019
Accepted: January 30, 2020

Elucidation of Molecular Mechanism Involved in Nephroprotective Potential of Naringin in Ethylene Glycol-Induced Urolithiasis in Experimental Uninephrectomized Hypertensive Rats

Jing LIU [1], Xian Cheng HAN [2], Dan WANG [3], Amit D. KANDHARE [4], Anwesha A. MUKHERJEE-KANDHARE [4], Subhash L. BODHANKAR [4] & Kun Ming WANG [5] *

[1] *Deparment of Pathology, Weifang Traditional Chinese Medicine Hospital, Weifang Shandong, 261041, China*
[2] *Department of Urology, Affiliated Hospital of Weifang Medical University, Weifang, Shandong, 261041, China*
3 Department of Emergency, Sunshine Union Hospital, Weifang, Shandong, 261041, China
[4] *Department of Pharmacology, Poona College of Pharmacy, Bharati Vidyapeeth Deemed University, Pune, Maharashtra, 411038, India*
[5] *Department of Urology, Sunshine Union Hospital, Weifang Shandong, 261041, China*

SUMMARY. **The objective of present investigation was to evaluate the nephroprotective potential of naringin against ethylene glycol (EG)-induced urolithiasis in experimental uninephrectomized hypertensive rats. EG (0.75% in drinking water) was used to induce urolithiasis in uninephrectomized hypertensive rats. Rats were treated with either naringin (20, 40, and 80 mg/kg, p.o.) for 28 days. Chronic administration of EG resulted in significant alterations ($p < 0.05$) in serum and urinary parameters (urea nitrogen, uric acid, creatinine, sodium, calcium, and LDH) whereas, administration of naringin (40 and 80 mg/kg) significantly attenuated ($p < 0.05$) these alterations. Naringin also significantly attenuated ($p < 0.05$) EG-induced hemodynamic and electrocardiographic abnormalities. Elevated levels of cardiac and renal MDA and nitric oxide and decreased levels of SOD and GSH were significantly restored ($p < 0.05$) by naringin treatment. RT-PCR analysis revealed that naringin significantly inhibited ($p < 0.05$) EG-induced upregulated mRNA expressions of renal KIM-1, NGAL, bikunin, and iNOs as well as down-regulated mRNA expressions of eNOs and OPN. Histological aberrations induced in renal and cardiac tissue after chronic administration of EG was significantly decreased ($p < 0.05$) by naringin. In conclusion, naringin exerts its nephroprotective effect against EG-induced via modulation of elevated oxidative stress and altered renal KIM-1, NGAL, bikunin, iNOs, eNOs, and OPN mRNA expressions.**

RESUMEN. El objetivo de la presente investigación fue evaluar el potencial nefroprotector de la naringina contra la urolitiasis inducida por etilenglicol (EG) en ratas hipertensas no nefrectomizadas experimentales. Se usó EG (0,75% en agua potable) para inducir urolitiasis en ratas hipertensas no nefrectomizadas. Las ratas se trataron con naringina (20, 40 y 80 mg/kg, p.o.) durante 28 días. La administración crónica de EG resultó en alteraciones significativas ($p < 0.05$) en los parámetros séricos y urinarios (nitrógeno ureico, ácido úrico, creatinina, sodio, calcio y LDH) mientras que la administración de naringina (40 y 80 mg/kg) atenuó significativamente ($p < 0.05$) estas alteraciones. La naringina también inhibió significativamente ($p < 0.05$) las alteraciones inducidas por EG en las anomalías hemodinámicas y electrocardiográficas. Los niveles elevados de MDA cardíaca y renal y óxido nítrico y la disminución de los niveles de SOD y GSH se restablecieron significativamente ($p < 0,05$) mediante el tratamiento con naringina. El análisis de RT-PCR reveló que la naringina inhibió significativamente ($p < 0.05$) las expresiones de ARNm reguladas por EG inducidas por KIM-1, NGAL, bikunina e iNOs, así como las expresiones de ARNm reguladas por disminución de eNO y OPN. Las aberraciones histológicas inducidas en el tejido renal y cardíaco después de la administración crónica de EG disminuyeron significativamente ($p < 0.05$) por acción de naringina. En conclusión, naringina ejerce su efecto nefroprotector contra la EG inducida por la modulación del estrés oxidativo elevado y la alteración de la expresión renal de KIM-1, NGAL, bikunina, iNOs, eNOs y ARNm de OPN.

INTRODUCTION

Nephrolithiasis or urolithiasis or urinary stone disease, which is the third most prevalent disorder of the urinary tract system affecting humankind since ancient times [1]. It has been well documented that calcium oxalate (CaOx) contributed approximately 80% to this urinary calculi disease which may result in hydronephrosis, obstruction, hemorrhage, and infection in the urinary system [2,3]. Albeit an array of sophisticated and comprehensive studies, the underlying pathogenesis of urolithiasis is still not well

KEY WORDS: bikunin, eNOs, ethylene glycol-induced urolithiasis, iNOs, kidney stone, naringin, osteopontin

* Author to whom correspondence should be addressed. *E-mail:* wangkunming_123@sina.com

ISSN 0326 2383 (printed ed.)
ISSN 2362-3853 (on line ed.)

Back to Academic Year 2020-21

www.ijcrt.org © 2021 IJCRT | Volume 9, Issue 10 October 2021 | ISSN: 2320-2882

IJCRT.ORG ISSN : 2320-2882

INTERNATIONAL JOURNAL OF CREATIVE RESEARCH THOUGHTS (IJCRT)

An International Open Access, Peer-reviewed, Refereed Journal

"Development and Validation of RP-HPLC and UV-Spectrophotometric Method for Simultaneous Estimation of Teneligliptin Hydrobromide Hydrate and Metformin Hydrochloride in Pharmaceutical dosage form"

Annasaheb S Gaikwad*[1], Swapnil R Raut[2], Goraksh S Bargaje[3]
Department of Quality Assurance Techniques,

1, 3 -Mangaldeep Institute of Pharmacy, Aurangabad. DBATU, M.S. India.
2- Rashtrasant Janardhan Swami College of Pharmacy, Kokamthan, Kopargaon. DBATU, M.S. India.

Abstract:

The objective of present research is to develop a simple, sensitive, linear, precise and accurate RP-HPLC and UV-Spectrophotometric method for simultaneous estimation of Teneligliptin Hydrobromide Hydrate and Metformin Hydrochloride in bulk and tablet formulation as developed and validated. UV-Spectrophotometric method Calibration plot were linear R^2 =0.9982 over the concentration range 0.8-1.6µg/ml for Teneligliptin Hydrobromide Hydrate, R^2 = 0.997 for the Metformin Hydrochloride 20-40µg/ml. And Chromatographic conditions used are stationary phase Grace C^{18} column (250mm × 4.6mm, 5µ particle size. The mobile phase Methanol: 10mm KH_2PO_4 Buffer (Ph-3) (70:30) and flow rate was maintained 0.8ml/min, detection wavelength was 240nm. The retention times were 4.5 min and 2.8 min for Teneligliptin Hydrobromide Hydrate and Metformin Hydrochloride respectively. Calibration plot were linear R^2 =0.9982 over the concentration range 1-5µg/ml for Teneligliptin Hydrobromide Hydrate, R^2 = 0.9981 for the Metformin Hydrochloride 25-125µg/ml. No interference from any component of pharmaceutical dosage form was observed. The proposed method has been validated as per ICH guidelines, validation studies revealed that method id specific, rapid, reliable and reproducible. The developed method successfully employed for routine quality control analysis in the combined pharmaceutical dosage form.

Keywords: Teneligliptin Hydrobromide, Metformin Hydrochloride, UV- Spectrophotometric, RP-HPLC Method, ICH Guideline.

Back to Academic Year 2020-21

WORLD JOURNAL OF PHARMACY AND PHARMACEUTICAL SCIENCES
SJIF Impact Factor 7.632
Volume 10, Issue 1, XXX-XXX Review Article ISSN 2278 – 4357

HIGH PERFORMANCE LIQUID CHROMATOGRAPHY-A REVIEW

Bharati R. Hole*[1], Annasaheb S. Gaikwad[2], Ankita A. Giramkar[2] and Dr. Dipak G. Umalkar[3]

[1]Department of Quality Assurance Technique.
[2]Saikrupa Institute of Pharmacy, Ghargaon, DBATU, M.S. India.
[3]Mangaldeep Institute of Pharmacy, Aurangabad. DBATU, M.S. India.

Article Received on
02 November 2020,
Revised on 23 Nov. 2020,
Accepted on 13 Dec. 2020
DOI: 10.20959/wjpps20211-17998

***Corresponding Author**
Bharati R. Hole
Department of Quality Assurance Technique.
annabhaugaikwad@gmail.com,

ABSTRACT

Chromatography is defined as a set of techniques which is used for the separation of constituents in a mixture. This technique involves two phases' stationary and mobile phases. High performance liquid chromatography (HPLC) is an important qualitative and quantitative technique, generally used for the estimation of pharmaceutical and biological samples. In these chromatography techniques, HPLC is one of the chromatographic techniques, which is mostly used analytical technique. Analytical Chemistry is the branch of Science that uses advance technologies in determining the composition by analytical technique. These strategies are fundamental for various purposes, including testing for quality control discharge, testing of solidness tests, testing of reference materials and to give information to bolster determinations. It is the most versatile, safest, dependable and fastest chromatographic technique for the quality control of drug components. This review article aims to focus on the different aspects of HPLC technique its principle, types, instrumentation and applications.

KEYWORDS: High Performance Liquid Chromatography, Mobile phase, Stationary phase.

INTRODUCTION

High Performance Liquid Chromatography which is also known as High Pressure Liquid Chromatography. The journey of chromatography was started first by Mikhail Tweet. (Russian botanist) in 19031It is a popular analytical technique used for the separation, identification and quantification of each constituent of mixture. In these chromatography techniques, HPLC is one of the chromatographic techniques, which is mostly used analytical

WJPPS COPY PROOF

Back to Academic Year 2020-21

ISSN: 2455-2631 © April 2020 IJSDR | Volume 5, Issue 4

Development and Validation of RP-HPLC and UV-Spectrophotometric Absorptivity Method for Simultaneous Estimation of Cyclobenzaprine hydrochloride and Aceclofenac in Pharmaceutical dosage form

Annasaheb S Gaikwad*[1], Monika N Madibone[2], Jeevan R Rajguru[3], Prashant B Jadhav[4]

[1]Department of Quality Assurance Techniques, Pravara Rural College of Pharmacy, Sinnar, Dist- Nashik, M.S. India, 422103,
[2,4]Department of Pharmaceutics, Srinath College of Pharmacy, Bajajnagar, Dist-Aurangabad, M.S. India- 431136
[3]Department of Pharmaceutics, AAEMFS Delight Institute of Pharmacy, Koregaon Bhima, Dist- Pune, M.S. India, 422103,

***Abstract*: The objective of present research is to develop a simple, sensitive, linear, precise and accurate RP-HPLC and UV-Spectrophotometric method for simultaneous estimation of Cyclobenzaprine hydrochloride and Aceclofenac in bulk and tablet formulation as developed and validated. UV-Spectrophotometric method Calibration plot were linear R^2=0.9996 over the concentration range 1-5µg/ml for Cyclobenzaprine hydrochloride, R^2= 0.9995 for the Aceclofenac 13-65µg/ml. And Chromatographic conditions used are stationary phase Inertsil ODS column (250 × 4.6 mm×5µ) (5µm particle size) the mobile phase Methanol: 10mm KH_2PO_4 Buffer (Ph-3) (70:30) and flow rate was maintained 0.9ml/min, detection wavelength was 220nm. The retention times were 4.7min and 2.9 min for Cyclobenzaprine hydrochloride and Aceclofenac respectively. Calibration plot were linear R^2 = 0.9996 over the concentration range 3-15µg/ml for Cyclobenzaprine hydrochloride, R^2 = 0.9994 for the Aceclofenac 40-200µg/ml. No interference from any component of pharmaceutical dosage form was observed. The proposed method has been validated as per ICH guidelines, validation studies revealed that method id specific, rapid, reliable and reproducible. The developed method successfully employed for routine quality control analysis in the combined pharmaceutical dosage form.**

***Keywords*: Cyclobenzaprine hydrochloride and Aceclofenac, UV- Spectrophotometric, RP-HPLC Method.**

I. INTRODUCTION:-

Cyclobenzaprine hydrochloride is a once daily extended release skeletal muscle relaxant which relieves muscle spasm of local origin without interfering with muscle function. The exact mechanism of action is unknown. The IUPAC name is 3-(5H-Dibenzo [A, D] Cyclohepten-5-Ylidene) N, N-Dimethyl-1-Propanamine Absorption: Immediate-release (IR) mean oral bioavailability ranges from 33% to 55. Distribution: About 93% is plasma protein-bound. Metabolism: During first pass through GI tract and liver, drug and metabolites undergo enterohepatic recycling. Cyclobenzaprine is eliminated quite slowly, with an effective half-life of 18 hours (range 8-37 hours; n=18); plasma clearance is 0.7 L/min. Metabolites excreted in the urine are likely water-soluble glucuronide conjugates.



Cyclobenzaprine Hydrochloride

Aceclofenac is a non-steroidal agent with marked anti-inflammatory and analgesic properties. The mode of action of aceclofenac is largely based on the inhibition to prostaglandin synthesis. Aceclofenac is a potent inhibitor of the enzyme cyclo-oxygenase, which is involved in the production of prostaglandins. The IUPAC name is [(2, 6-dichlorophenyl) amino] phenylacetoxyacetic acid. After oral administration, aceclofenac is rapidly absorbed and the bioavailability is almost 100%. Peak plasma concentrations are reached approximately1.25 to 3 hours. Aceclofenac is highly protein-bound (> 99.7%). Aceclofenac penetrates into the synovial fluid where the concentrations reach approximately 60%of those in plasma. Aceclofenac is probably metabolized via CYP2C9 to the main metabolite 4-hydroxyaceclofenac. The mean plasma elimination half-life is 4-4.3 hours.Approximately two-thirds of the administered dose is excreted via the urine, analysis conjugated hydroxymetabolites.

Back to Academic Year 2020-21

Original Article

Synthesis, Characterization and Antidiabetic Evaluation of Sulfonamide in Corporated with 1,3,4-Oxadiazole Derivatives

Megha Tukaram Salve*, Shailaja Bhanudas Jadhav Sr

Department of Pharmaceutical Chemistry of Modern College of Pharmacy, Nigdi, Pune, Maharashtra, INDIA.

ABSTRACT

Aim/Background: There is an increasing prevalence of diabetes mellitus throughout the world, and new compounds are necessary to combat this. While the current available antidiabetic therapies are long-term complicated and side effects-prone, this has led to a demand for more affordable, more effective methods of tackling diabetes. Research is focused on finding alternative medicinal remedies with significant antidiabetic efficacy as well as low adverse effects. This study synthesized, characterized, and evaluated anti-diabetic properties of synthetic sulphonamide hybrid of 1,3,4-oxadiazole derivatives. **Materials and Methods:** An *in vivo* antihyperglycemic evaluation of the sulphonamide hybrid of 1,3,4-oxadiazole derivatives was conducted using wistar rat models of type II diabetes derived from a two-step synthesis. Our study examined the effects of synthesized compounds on a model induced by a high-fat diet combined with streptozotocin and nicotinamide injection. For assessment of diabetic effects, Vildagliptin (10 mg/kg/day) was used as the standard drug. On day 14th, 1,3,4-oxadiazole derivatives (50 mg/kg/day) significantly lowered the blood sugar of hyperglycemic rats. **Results:** Due HFD STZ with Nicotinamide blood glucose level of wistar rat was increased (295 ± 8.2). After 14th day administration of derivatives random blood glucose level under controlled. A-III (220± 7.5 B) and A-IV (222 ± 3.62 B) were lowered random blood glucose levels on wistar rat. As compared to diabetes control (295 ± 8.5), derivatives of 1,3,4-oxadiazole are considered promising lead compounds. Compounds A-III and A-IV were found to be the most effective in lowering blood glucose, indicating the potential of these compounds as antidiabetic agents. **Conclusion:** Hybrids developed in this study provide new classes of anti-diabetic agents, and further optimization can be performed using this information.

Key words: *In vivo* study, 1,3,4-Oxadiazole, Sulfonamide derivative, anti-diabetic agents.

Submission Date: 24-05-2021;
Revision Date: 29-07-2021;
Accepted Date: 01-11-2021

DOI: 10.5530/ijper.55.4.214

Correspondence:
Mrs. Megha Tukaram Salve Jr,
Department of Pharmaceutical Chemistry of Modern College of Pharmacy, Nigdi, Pune-411044, Maharashtra, INDIA.
Phone: 91-0241-2328977,
Email – meghasalve77@gmail.com

www.ijper.org

INTRODUCTION

Hyperglycemia caused by defects in insulin secretion, insulin action, or both, is recognized as a diabetic condition.[1] A diabetic condition (DM) is a chronic disease characterized by hyperglycemia and is categorized into types 1 and 2 which may differ from each other in pathogenesis.[2] A diabetic condition is defined as high blood sugar levels (above 130 mg/dl) that cause long-term damage, dysfunction, and even failure to various body systems including the eyes, kidneys, nerves, heart, and vessels that carry blood.[3]

Several compounds containing heterocyclic rings are important components of antidiabetic pharmaceutical products. Nitrogen, sulfur, and oxygen containing heterocyclic compounds have attracted the attention of medicines chemical due to their wide range of biological applications. Among the heterocyclic family, 1,3,4-oxadiazole derivatives have shown many promising applications in pharmaceuticals.[4,5] Blood glucose stimulates the release of insulin from beta cells.[3] In addition to insulin deficiency and insulin resistance, there is also a genetic defect in the cells of the ß axis of the

Back to Academic Year 2020-21

Int. J. Pharm. Investigation, 2021; 11(4) : 374-378

Original Article

Synthesis, Characterization, and Evaluation of *in vitro* Antidiabetic Activity of Novel Pyrrolidine Sulphonamide Derivative

Megha Tukaram Salve*, Shailaja Bhanudas Jadhav
Department of Pharmaceutical Chemistry, Modern College of Pharmacy, Nigdi, Pune, Maharashtra, INDIA.

ABSTRACT

Background: Diabetes is a long-term illness characterized by high blood sugar levels. It is estimated that by 2045, there will be nearly 693 million diabetic patients worldwide, with half of the population remaining undiagnosed. Metformin, insulin, sulfonylureas, and thiazolidinediones were related to several risk factors, including hypoglycemia, bone fracture, weight gain, cardiovascular, renal, and other complications. In the present study, we have explored the DPP-IV inhibitors as a new class of antidiabetic drugs. **Objectives:** The goal of DPP-IV inhibitors is to raise levels of incretins (GLP-1 and GIP), which block glucagon release while boosting insulin secretion, slowing stomach emptying, and reducing blood glucose levels. **Methods:** A series of derivatives substituted on Oxadiazole of sulfonamide pyrrolidine were produced by reacting 1,2,4-oxadiazol and sulfonyl chloride at room temperature in the presence of ethanol and stirring until the reaction was complete. The compounds were characterized using IR, ^{1}H NMR, C^{13} NMR, mass spectroscopy, elemental analysis, and screened for *in vitro* assay of DPP-IV inhibition. **Results and Discussion:** The IC_{50} was calculated for compounds B-I, B-V, B-VI, B-XI, and B-XIII that inhibited enzymes significantly, IC_{50} values ranging from 19.65 ± 2.60 nM to 11.32 ± 1.59 nM, respectively, and Vildagliptin (4.79±1.66 IC_{50} nM) was used as a standard. The most active derivative substituted Oxadiazole of pyrrolidine sulfonamide is B-XI (11.32 ± 1.59 IC_{50} nM) among all synthesized compounds. **Conclusion:** B-XI derivative has shown appreciable DPP-IVinhibitory action. The 1,2,4-oxadiazol-3-yl pyrrolidine-1-sulfonamide derivatives have shown anti-diabetic properties.

Key words: Pyrrolidine sulphonamide derivative, 1,2,4-oxadiazole, *In vitro* study, Antidiabetic, Dipeptidyl peptidase-4.

Correspondence

Ms. Salve Megha Tukaram, Department of Pharmaceutical Chemistry, Modern College of Pharmacy, Nigdi, Pune, Maharastra-411044, INDIA.

Email id: meghasalve77@gmail.com;

DOI: 10.5530/ijpi.2021.4.67

INTRODUCTION

Diabetes is a group of chronic metabolic disorders caused by high blood sugar levels over a prolonged period.[1] Diabetes has become a global epidemic problem It was predicted that 451 million diabetics existed in 2017, with this Figure expected to increase to 693 million by 2045.[2] Most patients with type 2 diabetes, even when using anti-diabetic drugs, struggle to maintain sufficient glycemic regulation and develop microvascular and macrovascular diabetic complications.[3,4] Current diabetes therapy has lots of side effects.[5] The most frequent adverse events were hypoglycemia, stomach issues, and weight gain. DPP-IV inhibitors' possible advantages include a complementary mode of action with other anti-diabetic drugs, a desirable adverse-effect history, and a weight-neutral effect.[6,7]

By increasing endogenous GLP-1 levels, a DPP-IV inhibitor activates insulin and suppresses glucagon synthesis causing intestinal hypoglycemia.[8] Incretins and glucagon-like peptide-1 breakdown are suppressed by a DPP-IV inhibitor (GLP-1). DPP-IV inhibitor is an alternative diabetic therapy.[7,9,10] Because of their various pharmaceutical properties, five-member heterocyclic compounds, especially 1,2,4-Oxadiazole, have shown activity against a variety of diseases like Alzheimer's disease, parasitic worms (helminths) and other internal parasites, management and treatment of edematous and another non-edematous disease, infectious diseases, diabetes, pain and cramp, cardiovascular disease, HIV disease, tuberculosis, antioxidant, cancer, seizure disorders, inflammation of a joint.[11] Our efforts to enhance the efficacy, selectivity of the novel 1,2,4- oxadiazole derivative of pyrrolidine sulphonamide as DPP-IV inhibitors.

The sulfonamide moiety ($-SO_2NH_2$) is an active pharmacophore that the clinical and medicinal importance of sulfonamide drugs and compounds in the new drug discovery.[12] They exhibit a wide range of pharmacological activities, such as antimicrobial, antimalarial, anti-HIV, insulin-releasing antidiabetic, high ceiling diuretic, antithyroid, antitumor analgesic, and anti-inflammatory.[13-16] Some laboratories reported entities with sulfonamide portion; compounds 1 (IC_{50}; 6.7 nM) and 2 (IC_{50}; 39 nM) as potent DPP-IV inhibitors.[17] Omarigliptin (MK-3102) is sulfonamide containing moiety, which produces its antihyperglycaemic action by inhibiting the DPP-4 enzyme. Its pharmacokinetic studies have shown that it is suitable for once-a-week dosing, which makes it unique among the other DPP-4 inhibitors.[18,19] In this research we have designed novel 1,2,4- oxadiazole derivatives with sulfonamide and pyrrolidine-2-carbonitrile scaffolds specifically *in vitro* DPP-IV inhibitor assay as antidiabetic activity. In the continuation of our research to develop small molecules as biologically active antihyperglycaemic compounds, new derivatives having a pyrrolidine-2-carbonitrile-sulfonamide backbone hybrid were designed and synthesized in search for new potent DPP-4 inhibitors as antidiabetic agents.[20]

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

Back to Academic Year 2020-21

WORLD JOURNAL OF PHARMACEUTICAL RESEARCH
SJIF Impact Factor 8.084
Volume 10, Issue 14, 353-361. Review Article ISSN 2277– 7105

QUALITY BY DESIGN APPROACHES TO ANALYTICAL METHOD DEVELOPMENT: A MINI REVIEW

Shingne Snehal S.[1]*, Bhujbal Akshay E.[2], Bulbule Laxman D.[3*], Proff. Santosh Waghmare[4] and Dr. H. V. kamble[5]**

[1,2]Student, Department of Pharmaceutical Chemistry.
[4]HOD, Department of Pharmaceutical Chemistry.
[5]Principle, Loknete Shri Dada Patil Pharate College of Pharmacy, Mandavgan Pharata Tal. Shirur Dist. Pune, Maharashtra (412211).
[3]Student, Department of Pharmaceutical Chemistry, Pravara Rural College of Pharmacy, Pravaranagar, Tal. Rahata Dist. Ahemednagar, Maharashtra (423603).

Article Received on
21 September 2021,
Revised on 11 Oct. 2021,
Accepted on 01 Nov. 2021,
DOI: 10.20959/wjpr202114-22237

***Corresponding Author**
Shingne Snehal S.
Student, Department of Pharmaceutical Chemistry.

ABSTRACT

(QBD) Quality by design is a systematic approach to and drug development. Predefined product specifications are done by, planning, developing, manufacturing respectively. Science risk-based management approaches are used for a predefined product. Analytical development method based on the same principle of QBD known as AQBD. The Traditional approach and analytical approach are specified in QBD. Analytical profile, risk Assessment, different critical quality attributes are used for various analytical techniques. Method validation and control strategy is put in place. Design space for the development of a product is required according to ICH Q8 guideline i.e. method operational design region. Analytical techniques are streamlined pathways for new drug development. Gives a new effective and unique approach.

KEYWORDS: Quality, Analytical QBD, CQA, MODR.

INTRODUCTION

ICH Q8 guidelines define QBD as, "A systematic approach to development that begins with predefined objectives and emphasizes product and process understanding and process control, based on sound science and quality risk management." QBD is nothing but the relation between predefined product specifications in order by planning, developing formulations,

Back to Academic Year 2020-21

WORLD JOURNAL OF PHARMACEUTICAL RESEARCH
SJIF Impact Factor 8.084
Volume 10, Issue 14, 462-469. Review Article ISSN 2277– 7105

GENOTOXICITY: MALIGNANCY, DAMAGE TO DNA; A VOLUMINOUS REVIEW

Bhujbal Akshay E.*[1], Shingne Snehal S.[2], Proff. Santosh Waghmare[3] and Dr. H. V. Kamble[4]

[1,2]Student, Department of Pharmaceutical Chemistry, Loknete Shree Dada Patil Pharate Colloge of Pharmacy, Mandavgan Pharata Tal. Shirur, Dist. Pune, Maharashtra (412211).
[3]HOD, Department of Pharmaceutical Chemistry, Loknete Shree Dada Patil Pharate Colloge of Pharmacy, Mandavgan Pharata Tal. Shirur, Dist. Pune, Maharashtra (412211).
[4]Principle, Loknete Shree Dada Patil Pharate Colloge of Pharmacy, Mandavgan Pharata Tal. Shirur, Dist. Pune, Maharashtra (412211).

Article Received on
26 Sept. 2021,
Revised on 15 October 2021,
Accepted on 05 Nov. 2021
DOI: 10.20959/wjpr202114-22302

***Corresponding Author**
Bhujbal Akshay E.
Student, Department of Pharmaceutical Chemistry, Loknete Shree Dada Patil Pharate Colloge Of Pharmacy, Mandavgan Pharata Tal. Shirur, Dist. Pune, Maharashtra (412211).

ABSTRACT

A chemical agent that causes DNA damage is genotoxin. Germline mutation, somatic mutation lead to malignant transformation. The in-vivo, in-vitro test is performed for genotoxicity. The various assay is helped to detect DNA damage and is also used to evaluate the safety of environmental chemicals and to know the mechanism of action of suspected carcinogens. Many mutagens undergo metabolic activation for binding with DNA and DNA adduct in a reactive species can be detected in cells and human tissues by using various sensitive techniques. The characterization and the detection of DNA adducts provide causes of human cancer.

KEYWORDS: Carcinogens, mutagenicity, DNA adduct, immunoassay, genotoxin.

INTRODUCTION

Genotoxicity is a Latin term used for a chemical or biological agent having the ability to damage the DNA or genetic code of a cell, resulting in genetic mutation or carcinogenicity. A destructive effect on chromosomal material affects the integrity of the cell. Genotoxicity and mutagenicity both are different terms as all mutagens are genotoxic, but not all mutagenic compounds are genotoxic. Chemical substances and radiations are genotoxins. Indirect or

Isolation and characterization of phytoconstituents from pe Back to Academic Year 2020-21 a cochinchinensis fruits

Journal of Pharmacognosy and Phytochemistry
Available online at www.phytojournal.com

E-ISSN: 2278-4136
P-ISSN: 2349-8234
JPP 2018; 7(6): 2489-2492
Received: 19-09-2018
Accepted: 21-10-2018

Mohan Agrawal
Assistant Professor, Department of Pharmacognosy, Sitabai Thite College of Pharmacy, Shirur, Pune Maharashtra, India

Dr. Anilkumar Aher
MVP's College of Pharmacy, Nashik, Maharashtra, India

Dr. Subodh Pal
MVP's College of Pharmacy, Nashik, Maharashtra, India

Dr. Deelip Derle
MVP's College of Pharmacy, Nashik, Maharashtra, India

Correspondence
Mohan Agrawal
Assistant Professor, Department of Pharmacognosy, Sitabai Thite College of Pharmacy, Shirur, Pune Maharashtra, India

Isolation and characterization of phytoconstituents from petroleum ether extract of *Momordica cochinchinensis* fruits

Mohan Agrawal, Dr. Anilkumar Aher, Dr. Subodh Pal and Dr. Deelip Derle

Abstract
The purpose of the study is to isolate and characterize the chemical constituents from fruits of *Momordica cochinchinensis*. The principal theme of the study is to develop applied chromatographic techniques for the separation, isolation and detection of the compounds. The petroleum ether extract of fruits of *Momordica cochinchinensis* was saponified and unsaponifiable matter was subjected to column chromatography and elution of column was carried out by Petroleum ether (100%) with increasing concentration of ethyl acetate for the separation of phytoconstituents. The isolated compounds were characterized and analyzed by physical characteristics, IR, NMR and Mass spectroscopy. Three known compounds lupeol, β-amyrin and β-sitosterol were determined for the first time from fruits of *Momordica cochinchinensis*. From the present study, it is concluded that the plant consists of phytosterols which can be isolated and characterized by chromatographical and spectroscopical method.

Keywords: *Momordica cochinchinensis,* phytosterols, lupeol, β-amyrin and β-sitosterol

Introduction
From thousands of years Natural products have been used by human societies. Natural sources have provided considerable value to the pharmaceutical industry over the past half century. Research studies leading to extraction, isolation and biological study of plant constituents have formed the major field of study. Various leads from plant sources were taken for discovering the new active therapeutic agents. Hence herbal medicine has played important role in managing the health conditions like diabetes, hypertension, inflammation, obesity, cancer, etc [1].
The plant *Momordica cochinchinensis* (Gac) is a Southeast Asian fruit found throughout the region from Southern China to Northeastern Australia, mostly Vietnam and throughout India. It grows on dioecious vines and usually collected from fence climbers or from wild plants [2]. It is reported that *Momordica cochinchinensis* Spreng, is one of the special fruits containing extraordinarily high levels of carotenoids, especially β -carotene (more than 16 mg/100 g) and lycopene (more than 50 mg/100 g), mainly in the red aril [3]. It is also reported to contain a protein that may inhibit the proliferation of cancer cells and also β-carotene with several phytonutrients, Vit-E, fatty acids, carbohydrates, flavonoidal glycosides [4].
Traditionally, Gac has been used as both food and medicine and promotes healthy vision by relief of dry eyes. It also possesses antioxidant, anti-microbial and antidiabetic properties. The seeds are considered to be good for cough and pains in the chest [5, 6].
But these studies are not enough for identifying and characterizing the bioactive compounds in the plant. The purpose of the study is to identify and characterize the bioactive principles from fruits of *Momordica cochinchinensis*.

Materials and Methods
Plant material: The fruits of *Momordica cochinchinensis* was collected from rural area of Kolkata. The herbarium of the *Momordica cochinchinensis* was authenticated by Botanical Survey of India, Pune Voucher specimen (MA 01) was deposited in library.

Preparation of Extracts: Fruits of *Momordica cochinchinensis* was extracted by Soxhlet extractor with Pet ether and macerated with ethanol and water successively.

Storage of Extracts: All the extracts were stored in tightly closed glass bottles in refrigerator at 2-8 ^{0}C.

Back to Academic Year 2020-21

ORIGINAL ARTICLE

Analgesic activity of *Momordica cochinchinensis* and *Momordica balsamina* fruit extracts

Mohan R. Agrawal[1]*, Anilkumar N. Aher[2], Subodh C. Pal[2], Deelip V. Derle[2]

[1]Department of Pharmacognosy, Sitabai Thite College of Pharmacy, Pune, Maharashtra, India, [2]Department of Pharmacognosy, MVP's College of Pharmacy, Nashik, Maharashtra, India

Abstract

Introduction: In the present study, fruit extracts of *Momordica cochinchinensis (Cucurbitaceae)* and *Momordica balsamina (Cucurbitaceae)* were investigated for analgesic activity by Eddy's hot plate and Tail immersion method. **Materials and Methods:** The extracts were prepared successively using powdered material with petroleum ether, ethanol, and water, and concentrated under vacuum and were evaluated for analgesic activity at three dose level (100, 200, and 400 mg/kg). **Results and Discussion:** In Eddy's hot plate method, oral administration of petroleum ether extracts of both the plants at the dose of 200 mg/kg ($P < 0.01$) and 400 mg/kg ($P < 0.001$) significantly reduced the thermal stimulation. Analgesic activity of petroleum ether extracts of both plants at the dose of 400 mg/kg after 90 min was comparable to standard drug pentazocine (10 mg/kg). In tail immersion method, petroleum ether extract at the dose of 100 mg/kg ($P < 0.05$), 200 mg/kg, and 400 mg/kg ($P < 0.01$) and alcoholic extract at the dose of 200 mg/kg and 400 mg/kg ($P < 0.05$) of both plant material has shown significant analgesic activity and was comparable to standard drug pentazocine (10 mg/kg) after 90 min. **Conclusion:** It is concluded that petroleum ether extracts of both plant material have central analgesic effects.

Key words: Analgesic activity, Eddy's hot plate, *Momordica balsamina*, *Momordica cochinchinensis*, phytosterols, tail immersion

INTRODUCTION

Momordica is a genus of about 60 species of annual or perennial climbers herbaceous or rarely small shrubs belonging to the family Cucurbitaceae, natives of tropical and subtropical Africa, Asia and Australia.[1,2]

Momordica cochinchinensis (Gac) is a Southeast Asian fruit found throughout the region from Southern China to Northeastern Australia, mostly Vietnam and throughout India. It grows on dioecious vines and is usually collected from fence climbers or wild plants. The vines can be commonly seen growing on lattices at the entrances to rural homes or in gardens. It bears fruits annually and is found in local markets. The fruit becomes a dark orange color on ripening, and is typically round or oblong, maturing to a size of about 13 cm in length and 10 cm in diameter. The exterior skin is covered in small spines, while dark red interior consists of clusters of fleshy pulp and seeds.[3] Gac fruit, *M. cochinchinensis* Spreng, is one of the special fruits containing extraordinarily high levels of carotenoids, especially β-carotene (>16 mg/100 g), and lycopene (>50 mg/100 g), mainly in the red aril.[4] Conventionally, Gac has been used as both food and medicine and promotes healthy vision by relief of dry eyes. It also possesses antioxidant, antimicrobial, and antidiabetic properties. The seeds are considered to be good for cough and pains in the chest.[5-7]

Momordica balsamina Family: Cucurbitaceae is climber with bright green leaves bears striking orange to red spindle-shaped ripe fruit. Shurb is fairly common and widespread in Malaya, Australia, West Asia, Africa, America, and India (Sind, Gujarat, and Deccan). Conventionally used as a purgative agent, purification of blood and dissipate

Address for correspondence:
Mohan R. Agrawal, Department of Pharmacognosy, Sitabai Thite College of Pharmacy, Pune, Maharashtra, India. Tel.: +91-2138-222680/9860947093.
E-mail: agrawalmohan19@gmail.com

Received: 15-05-2018
Revised: 10-08-2018
Accepted: 12-10-2018

Back to Academic Year 2020-21

Research Journal of Pharmacognosy and Phytochemistry. 12(3): July - September, 2020

ISSN 0975-2331 (Print)
0975-4385 (Online)
DOI: 10.5958/0975-4385.2020.00026.6

Vol. 12 |Issue-03|
July-September| 2020

Available online at
www.anvpublication.org
Research Journal of Pharmacognosy and Phytochemistry
Home page *www.rjpponline.org*

RESEARCH ARTICLE

Isolation, Characterization of Phytoconstituents from petroleum ether extract of *Momordica balsamina* fruits and its quantitative analysis by HPTLC

M.R. Agrawal[1]*, Dr. A.N. Aher[2], Dr. S.C. Pal[2]
[1]Sitabai Thite College of Pharmacy, Shirur, Pune (District), Maharashtra-412210, India.
[2]MVP's College of Pharmacy, Nashik, Maharashtra, India.
*Corresponding Author E-mail: **agrawalmohan19@gmail.com**

ABSTRACT:
The purpose of the study is to isolate and characterize the chemical constituents from fruits of *Momordica balsamina.* The principal theme of the study is to develop applied chromatographic techniques for the separation, isolation and detection of the compounds. The petroleum ether extract of fruits of *Momordica balsamina* was saponified and unsaponifiable matter was subjected to column chromatography andelution of column was carried out by Petroleum ether (100%) with increasing concentration of ethyl acetate for the separation of phytoconstituents. The isolated compounds were characterized and analyzed by physical characteristics, IR, NMR and Mass spectroscopy. Four known compounds lupeol, ursolic acid, β-amyrin and β-sitosterol were isolated from fruits of *Momordica balsamina*. The isolated compounds were quantitatively analyzed by HPTLC. The concentration of lupeol, ursolic acid and β-amyrin was found to be 1.15%, 0.6% and 2.8%. From the present study, it is concluded that the plant consists of phytosterols and triterpene.

KEYWORDS: *Momordica balsamina,* Phytosterols, lupeol, ursolic acid, β-amyrin and HPTLC.

INTRODUCTION:

Momordica is a genus of annual or perennial climbers is essentially a native of tropical regions on Asia distributed extensively in China, Japan, South East Asia, Polynesia besides tropical Africa and South America. Many species of *Momordica* genus have been found to grow wildly in India, Bangladesh, Srilanka, Myanmar, Malay etc[1].

Momordica balsamina also refers to as balsam fruit or apple belongs to the family of *Cucurbitaceae*. It is a plant commonly used by local poultry farmers in Plateau State, Nigeria for general well being of birds. The Balsam apple is a climber or trailer with annuals stems attaining 4-5 m length. The fruit is orange yellow, beaked, 21/2 inches in length bursting and exposing red brown seeds. The species is closely related to *Momordicacharantia* (Bitter mellon) which occurs in areas of greater rainfall and whose properties and actions includes antibacterial, anti-inflammatory, anti-oxidant, antiviral, immunostilmulant, hypoglycemic etc[2-5]. Some *Momordica* species are already established as medicine and their marketed preparations are available and used in the treatment of different diseases. The rationale behind the study is to isolate, characterize and quantitatively determine the constituents present in petroleum ether extract of *Momordica balsamina* fruits.

Received on 09.07.2020 Modified on 21.07.2020
Accepted on 28.07.2020 ©AandV Publications All right reserved
Res. J. Pharmacognosy and Phytochem. 2020; 12(3):156-161.
DOI: 10.5958/0975-4385.2020.00026.6

Back to Academic Year 2020-21

Academic Year 2021-2022

Sr. no	Title of the paper	Name of Staff	Department	Name of the journal	ISSN number	Document	Link
1	Excipients Compatibility Study Of Antilipidemic And Antihypertensive Drugs In Oral Solid Dosage Form	Dr. V. M. Satpute	Pharmaceutics	Neuroquantology	ISSN1303-5150	View Document	https://www.google.com/search?q=Excipients+Compatibility+Study+Of+Antilipidemic+And+Antihypertensive+Drugs+In+Oral+Solid+Dosage+Form+research+paper+by+dr.+v.+m.+satpute&rlz=1C1VDKB_en-GBIN1070IN1070&oq=Excipients+Compatibility+Study+Of+Antilipidemic+And+Antihypertensive+Drugs+In+Oral+Solid+Dosage+Form+research+paper+by+dr.+v.+m.+satpute&gs_lcrp=EgZjaHJvbWUyBggAEEUYOTIGCAEQRxgnMgYIAhAjGCfSAQkyMTQwNjJ2owajeoAgiwAgE&sourceid=chrome&ie=UTF-8
2	Formulation And Evaluation Of Selexipag And Ezetimibe Bi-Layer Tablet By Novel Approach	Dr. V. M. Satpute	Pharmaceutics	Neuroquantology	ISSN1303-5150	View Document	https://www.neliti.com/publications/429774/formulation-and-evaluation-of-selexipag-and-ezetimibe-bi-layer-tablet-by-novel-a
3	Evaluation of Some Natural Polymers as Pharmaceutical Excipient For Sustained Drug Delivery System	Mr. A. V. Dhobale	Pharmaceutics	International Journal of Research and Analytical Reviews (IJRAR)	www.ijrar.org E-ISSN 2348-1269	View Document	https://www.researchgate.net/publication/350782955_Evaluation_of_Some_Natural_Polymers_as_Pharmaceutical_Excipient_For_Sustained_Drug_Delivery_System

4	Green Chemistry	Mr. P. B. Cholke & Y. S. Temak	Pharmaceutics	International Journal of Creative Research Thoughts	ISSN: 2320-2882	View Document	https://www.google.com/search?q=Green+Chemistry+research+paper+by+pravin+cholake&sca_esv=9c9edb63dbab71a4&sca_upv=1&rlz=1C1VDKB_en-GBIN1070IN1070&biw=1366&bih=641&sxsrf=ACQVn09_J2GQj8sO9QI6WxcNZ8to4sgBjw%3A1713679595795&ei=66wkZvuJMIzT1e8Pnaug2A4&ved=0ahUKEwi7q8P80dKFAxWMafUHHZ0VCOsQ4dUDCBA&uact=5&oq=Green+Chemistry+research+paper+by+pravin+cholake&gs_lp=Egxnd3Mtd2l6LXNlcnAiMEdyZWVuIENoZW1pc3RyeSByZXNlYXJjaCBwYXBlciBieSBwcmF2aW4gY2hvbGFrZTIHECEYoAEYCjIHECEYoAEYCkjBYFAAWJVccAB4AJABAZgB7AOgAd40qgEKMC42LjE3LjQuMbgBA8gBAPgBAvgBAZgCGKAC2C7CAggQABiABBiiBMICBRAhGKABwgIEECEYCsICBRAhGJ8FwgIGEAAYFhgewgILEAAYgAQYhgMYigXCAgQQIxgnmAMAkgcKMC41LjE0LjQuMaAH5YQB&sclient=gws-wiz-serp
5	Gastro Retentive Drug Delivery System	Dr. H. V. Kamble	Pharmacology	World Journal of Pharmaceutical Research	ISSN 2277 – 7105	View Document	https://wjpr.net/abstract_show/18498
6	Gastro Retentive Drug Delivery System	Dr. S. A. Waghamare	Pharmaceutical Chemistry	World Journal of Pharmaceutical Research	ISSN 2277 – 7105	View Document	https://wjpr.net/abstract_show/18498

7	A Review on Method Development, Validation, Optimization and Applications of HPLC	Dr. H. V. Kamble	Pharmacology	International Journal for Research in Applied Science & Engineering Technology	ISSN: 2321-9653	View Document	https://www.ijraset.com/research-paper/method-development-validation-optimization-and-applications-of-hplc
8	A Review on Method Development, Validation, Optimization and Applications of HPLC	Dr. S. A. Waghamare	Pharmaceutical Chemistry	International Journal for Research in Applied Science & Engineering Technology	ISSN: 2321-9653	View Document	https://www.ijraset.com/research-paper/method-development-validation-optimization-and-applications-of-hplc
9	A Review: Drug-Excipient Interactions Study	Dr. H. V. Kamble	Pharmacology	World Journal of Pharmaceutical Research	ISSN 2277 – 7105	View Document	https://ijnrd.org/viewpaperforall.php?paper=IJNRD2302162
10	A Review: Drug-Excipient Interactions Study	Dr. S. A. Waghamare	Pharmaceutical Chemistry	World Journal of Pharmaceutical Research	ISSN 2277 – 7105	View Document	https://ijnrd.org/viewpaperforall.php?paper=IJNRD2302162
11	A Review On: Preservatives Used in Pharmaceuticals and Impacts on Health	Dr. S. A. Waghamare	Pharmaceutical Chemistry	Iconic Research And Engineering Journals	| ISSN: 2456-8880	View Document	https://scholar.google.co.in/scholar?q=A+Review+On:+Preservatives+Used+in+Pharmaceuticals+and+Impacts+on+Health+by+-dr.+santosh+a.+waghmare&hl=en&as_sdt=0&as_vis=1&oi=scholart
12	A Review: Drug-Excipient	Dr. V. M. Satpute	Pharmaceutical Chemistry	World Journal of Pharmace	ISSN 2277	View Document	https://ijnrd.org/viewpaperforall.php?paper=IJNRD2302162

	Interactions Study			utical Research	– 7105		
13	An Overview on Artificial Blood	Ms. S. B. Kshirsagar	Pharmaceutical Chemistry	International Journal In Pharmaceutical Sciences	Vol 1, Issue 1, 42-48	View Document	https://scholar.google.co.in/scholar?q=An+Overview+on+Artificial+Blood+by+s.+b.+kshirsagar&hl=en&as_sdt=0&as_vis=1&oi=scholart
14	A Review On: Preservatives Used in Pharmaceuticals and Impacts on Health	**Dr. H. V. Kamble**	Pharmacology	Iconic Research And Engineering Journals	| ISSN: 2456-8880	View Document	https://scholar.google.co.in/scholar?q=A+Review+On:+Preservatives+Used+in+Pharmaceuticals+and+Impacts+on+Health+by+-dr.+santosh+a.+waghmare&hl=en&as_sdt=0&as_vis=1&oi=scholart

Back to Academic Year 2021-22

Excipients Compatibility Study Of Antilipidemic And Antihypertensive Drugs In Oral Solid Dosage Form



EXCIPIENTS COMPATIBILITY STUDY OF ANTILIPIDEMIC AND ANTIHYPERTENSIVE DRUGS IN ORAL SOLID DOSAGE FORM

Satpute Vivek Mahavir
Ph.D. Scholar, Bhagwant University Sikar Road, Ajmer (RAJASTHAN)
Correspondence author email: satputevivek9@gmail.com
Dr. Punit R. Rachh
Bhagwant University Sikar Road, Ajmer (RAJASTHAN)

Abstract

Excipients compatibilitystudy of antilipidemic and antihypertensive drugs like Selexipag and Ezetimibe respectively was carried out with selected generally regarded as safeexcipients to prepare oral solid dosage formi.e.bilayer tablet. Isothermal stress testing was performed in binary a mixture which was subjected to 50°C for 4 weeks. Isothermal stressed samples were evaluated with RP-HPLC method and FTIR analysis. A trial was conducted with single disintegrants and diluent with direct compression technique and in-vitro dissolution was carried out. To improve the release profile of Immediate Release layer of Selexipag, multiple disintegrants were used namely croscarmellose sodium (CCS), sodium starch glycolate (SSG) and Crospovidone (CP). Similarly, extended release layer of Ezetimibe was prepared by wet granulation technique. Intra and extra granulation was done with mixtures of disintegrants. Extended release layer of Ezetimibe was formulated using polymers and gums like HPMC, MCC, PVP, Talc, Gaur gum, and Xanthan gum, Magnesium stearate. The results show that Selexipag and Ezetimibe were compatible with the all the polymers used in the study. The polymers used in the present study and the formulations developed using the compatible excipients were found to be stable.

Keywords: Ezetimibe, Selexipag, Compatibility, Isothermal stress testing, Bilayer tablet.

DOI Number: 10.14704/nq.2022.20.9.NQ44744 **Neuro Quantology 2022; 20(9):6349-6356**

Introduction

The safety, efficacy, quality and stability of a formulation are the cornerstones of any new drug development process. In order to consistently maintain these attributes in a finished dosage form, it is important to have a comprehensive understanding of the Physico-chemical characteristics of the active pharmaceutical ingredient (API), as well as all other components (e.g. excipients, manufacturing aids, packaging materials) of the drug product[1].The preformulation stage involves characterization of several aspects of the API including solubility, dissolution, permeability, polymorph/salt screening, stability, ionization properties, particle size distribution, API-excipient compatibilities etc[2].

Due to the intimate contact of the API with one or more excipients in a formulation, there exists a likelihood of physical and/or chemical interactions between them. Any such interactions may result in a negative impact on the physical, stability or performance attributes of the drug product. The choice of excipients is of crucial importance to avoid these negative effects, and to facilitate the development of a robust and an effective formulation. Thus, for a rational selection of excipients, screening of excipient-API compatibility is recognized as an important aspect of formulation[3].

Selexipag is used in adults to treat pulmonary arterial hypertension (PAH, high blood pressure in the vessels that carry blood to the lungs) to slow down the worsening of symptoms and reduce the chance of being hospitalized for PAH.

Back to Academic Year 2021-22

Gastro Retentive Drug Delivery System

WORLD JOURNAL OF PHARMACEUTICAL RESEARCH
SJIF Impact Factor 8.084
Volume 11, Issue 2, 254-272. Review Article ISSN 2277– 7105

A REVIEW: GASTRO RETENTIVE DRUG DELIVERY SYSTEM

***Suresh Choudhary, Santosh Waghmare and Hemant Kamble**

Loknete Shri Dadapatil Pharate College of Pharamcy, Mandavgoan Pharat Tal-shirur, Dist-Pune, Maharashtra.

Article Received on
06 December 2021,
Revised on 26 Dec. 2021,
Accepted on 16 Jan. 2022,
DOI: 10.20959/wjpr20222-22766

***Corresponding Author**
Suresh Choudhary
Loknete Shri Dadapatil Pharate College of Pharamcy, Mandavgoan Pharat Tal-shirur, Dist-Pune, Maharashtra.

ABSTRACT

The goal of this review on gastroretentive drug delivery systems (GRDDS) was to compile the current literature with a focus on a few gastroretentive approaches that have recently emerged as important methodologies in the field of site-specific orally administered sustained/controlled release drug delivery. To overcome physiological challenges such as short gastric residence times (GRT) and unpredictable gastric emptying times, technological efforts have been made in the study and development of rate-controlled oral drug delivery systems (GET). GRDDS is a method for extending the GRT and targeting site-specific drug release in the upper gastrointestinal tract (GIT) for a local or systemic effect. Because of the fast gastric transition from the stomach, conventional oral dose forms have low bioavailability issues, especially for medications that are less soluble at an alkaline pH of the intestine. Additionally, medications that have a local action in the stomach are quickly emptied and do not spend enough time in the stomach. To lower the frequency of dose administration, several efforts have been made to extend the retention duration of drug delivery systems. GRDFs not only extend dosing intervals, but they also improve patient compliance beyond what is currently possible with controlled release dosage forms. The benefits, drawbacks, and characterisation of gastroretentive drug delivery devices are discussed in this article. This study contains patents and commercially marketed gastroretentive products.

KEYWORDS: Gastroretentive drug delivery system; approach; floating system; Evaluation.

INTRODUCTION

The oral delivery of drugs is the most favoured route of administration because of ease of administration. The bioavailability of drugs in oral dose forms is affected by a variety of

Back to Academic Year 2021-22

International Journal for Research in Applied Science & Engineering Technology (IJRASET)
ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538
Volume 10 Issue V May 2022- Available at www.ijraset.com

A Review on Method Development, Validation, Optimization and Applications of HPLC

Sumit P. Isane[1], Santosh A. Waghmare[2], Hemant V. Kamble[3]
[1]Student, Department of Pharmaceutics, [2]Professor, Department of Pharmacy, [3]Principle, Loknete Shri Dadapatil Pharate College of Pharmacy, Mandavgan Pharata, Shirur, Pune, Maharashtra, India

Abstract: ***High-Performance Liquid Chromatography (HPLC) is a technique used to separate components from the mixture. The technology consists of two phases, the stationary phase, and the mobile phase. Component separation is based on differences in partition coefficients in two phases.***
The present review article focuses on the principle, types, instrumentation, process validation, application, advantages, disadvantages of HPLC. HPLC methods development and validation play important roles in the discovery, development, manufacture of pharmaceutical drugs, and various other studies related to humans and animals. An analytical procedure is developed to test a defining characteristic of the drug substance or drug product against established acceptance criteria for that characteristic. Validation of HPLC method as per ICH Guidelines covers all the performance characteristics of validation, like Accuracy, precision, specificity, linearity, range, the limit of detection, the limit of quantification, robustness, and system suitability testing.
Keywords: HPLC, Types, Method validation, Method development, Instrumentation, and Applications.

I. INTRODUCTION

High-Performance Liquid Chromatography (HPLC) involves separations in which the components to be separated are distributed between two immiscible phases.
One of these phases is the mobile phase, and the other is a stationary phase[1,2]. High-Performance Liquid Chromatography is now one of the most powerful tools in analytical chemistry. It can separate, identify, and quantify the compounds that are present in any sample that can be dissolved in a liquid. High-performance liquid chromatography (HPLC) is the most accurate analytical method widely used for the quantitative as well as qualitative analysis of drug products[3]. HPLC is an advanced technique of column liquid chromatography. The solvent usually flows through the column with the help of gravity but in the HPLC technique, the solvent will be forced under high pressure up to 400 atmospheres so that sample can be separated into different constituents with the help of difference in relative affinities[4-9]. n HPLC, pumps will be used to pass pressurized liquid solvent including the sample mixture which is allowed to enter into a column filled with a solid adsorbent material. The interaction of each sample component will be varied and this causes the difference in flow rates of each component and finally leads to the separation of components of the column.
Chromatography can be depicted as a mass exchange process including adsorption. HPLC depends on pumps to pass a pressurized fluid and an example blend through a section loaded with adsorbent, prompting the partition of the specimen segments. The dynamic segment of the section, the adsorbent, is regularly a granular material made of solid particles (e.g. silica, polymers, etc.) 2 µm to 50 µm in size.
The segments of the example mixture/blend are isolated from each other because of their distinctive degrees of connection with the retentive particles. The pressurized fluid is commonly a blend of solvents (e.g. water, acetonitrile/methanol) and is known as the 'mobile phase'. Its organization and temperature play an important part in the partition procedure by affecting the connections occurring between sample segments and adsorbent[10-17].
HPLC is recognized from traditional ("low weight") liquid chromatography because operational pressures are fundamentally higher (50 bar to 350 bar), while normal liquid chromatography regularly depends on the power of gravity to pass the portable stage through the segment. Because of the small sample amount isolated in scientific HPLC, column section measurements are 2.1 mm to 4.6 mm distance across, and 30 mm to 250 mm length. Additionally, HPLC segments are made with smaller sorbent particles (2 µm to 50 µm in normal molecule size). This gives HPLC high determining or resolving power (the capacity to recognize components) while isolating mixtures, which makes it a prominent chromatographic method.

© IJRASET: All Rights are Reserved | SJ Impact Factor 7.538 | ISRA Journal Impact Factor 7.894 |

Back to Academic Year 2021-22

A Review: Drug- Excipient Interactions Study

WORLD JOURNAL OF PHARMACEUTICAL RESEARCH
SJIF Impact Factor 8.084
Volume 10, Issue 12, 214-224. Review Article ISSN 2277– 7105

A REVIEW: DRUG- EXCIPIENT INTERACTIONS STUDY

Manish Wagh*, Santosh Waghmare and Hemant Kamble

Loknete Shri Dadasaheb Pharate Patil College of Pharmacy, Mandavgoan Pharata, Tal-Shirur, Dist.-Pune. Maharashtra.

Article Received on
03 August 2021,
Revised on 23 August 2021,
Accepted on 13 Sept. 2021
DOI: 10.20959/wjpr202112-21601

***Corresponding Author**
Manish Wagh
Loknete Shri Dadasaheb Pharate Patil College of Pharmacy, Mandavgoan Pharata, Tal-Shirur
Dist.-Pune. Maharashtra.

ABSTRACT

Excipients are included in dosage forms to aid manufacture, administration or absorption. Although considered pharmacologically inert, excipients can initiate, propagate or participate in chemical or physical interactions with drug compounds, which may compromise the effectiveness of a medication. Exicipients are not exquisitely pure. Even for the most commonly used excipients, it is necessary to understand the context of their manufacture in order to identify potential active pharmaceutical ingredients interactions with trace components. Chemical interactions can lead to degradation of the active ingredient, thereby reducing the amount available for therapeutic effect. Physical interactions can affect rate of dissolution, uniformity of dose or ease of administration.

KEYWORDS: Excipient, Drug, Interaction, Physical, Chemical.

INTRODUCTION

Pharmaceutical dosage form is a combination of active pharmaceutical ingredients (API) and excipients. Excipients are included in dosage forms to aid manufacture, administration or absorption (Crowley and Martini). The ideal excipients must be able to fulfill the important functions i.e. dose, stability and release of API from the formulation. Although considered pharmacologically inert, excipients can initiate, propagate or participate in chemical or physical interactions with drug compounds, which may compromise the effectiveness of a medication. Exicipients are not exquisitely pure. In common with virtually all materials of minerals, synthetic, semi-synthetic or natural origin manufacture involves using starting materials, reagents and solvents. Residues invariably remain after isolation. Often, it is the multi-component nature of the excipient that drives many of the interactions with APIs. Even

Back to Academic Year 2021-22

Formulation And Evaluation Of Selexipag And Ezetimibe Bi-Layer Tablet By Novel Approach



FORMULATION AND EVALUATION OF SELEXIPAG AND EZETIMIBE BI-LAYER TABLET BY NOVEL APPROACH

Satpute Vivek Mahavir
Ph.D. Scholar, Bhagwant University Sikar Road, Ajmer (RAJASTHAN)
Correspondence author email: satputevivek9@gmail.com
Dr. Punit R. Rachh
Bhagwant University Sikar Road, Ajmer (RAJASTHAN)

Abstract

Hypertension and hypercholesterolemia are two common diseases that often coexist. The combined presence of hypertension and hypercholesterolemia increases considerably the risk for cardiovascular complications.Selexipag is used to treat symptoms of pulmonary arterial hypertension and Ezetimibe is a lipid-lowering compound that inhibits intestinal cholesterol and phytosterols absorption.The bilayer tablets of antilipidemic and antihypertensive can be employed for the treatment of hypertension and hypercholesterolemia together through oral administration of single tablet. The rationale for this fixed-dose combination is to co-administer two drugs acting by different mechanisms of action together, reduce dosing frequency, and increase patient compliance.In the present investigation we combine Selexipag and Ezetimibe. Immediate release layer of Selexipag was formulated using various superdisintegrants like Crospovidone, Croscarmellose sodium, and sodium starch glycolate, Microcrystalline Cellulose, Magnesium Stearate and Extended release layer of Ezetimibe was formulated using polymers and gums like Hydroxy Propyl methyl cellulose,Poly vinyl Pyrrolidone, Talc, Gaur gum, and Xanthan gum, Magnesium stearate. The disintegration and dissolution study of both layers showed that inclusion of surfactant to the tablet formulation (Immediate Release) and dissolution medium (Extended Release) enhanced the release of drugs from both layers. Dissolution studies of optimized IR layer (SL6) and ER layer (EZ5) showed good dissolution. The above results reveal that the optimized immediate release layer Selexipag (SL6) and extended-release layer of Ezetimibe (EZ5) might be suitable for the treatment of hypertension and hypercholesterolemia by sequential release of the two drugs in a bilayer tablet.

Keywords: Ezetimibe, Selexipag, Sustained release, immediate release, Bilayer tablet.

DOI Number: 10.14704/nq.2022.20.9.NQ44745 **Neuro Quantology 2022; 20(9):6357-6368**

Introduction

Early trials on the treatment of hypertension based upon high doses of diuretics or beta blockers demonstrated a significant reduction in stroke but a less prominent decline in the incidence of coronary heart disease. This relative lack of coronary benefit may have been related to the adverse effect of these antihypertensive agents on plasma lipids since even small elevations in serum cholesterol may significantly increase coronary risk, particularly in patients with underlying hypertension. Drug-induced changes in lipid levels may be particularly important in hypertensive since up to 40 percent of untreated patients with primary hypertension (formerly called "essential" hypertension) already have lipid abnormalities, such as a high low-density lipoprotein-cholesterol and a low high-density lipoprotein-cholesterol.[1]

Hence, hypertension frequently coexists with hyperlipidemia and both are considered to be major risk factors for developing cardiac disease ultimately resulting in adverse cardiac events. This clustering of risk factors is potentially due to a common mechanism. Further, patient compliance with the management of hypertension is generally better than patient compliance with hyperlipidemia. It would, therefore, be advantageous for patients with the intent of providing commercially viable multi-

Back to Academic Year 2021-22

Evaluation of Some Natural Polymers as Pharmaceutical Excipient For Sustained Drug Delivery System

ResearchGate

See discussions, stats, and author profiles for this publication at: https://www.researchgate.net/publication/350782955

"Evaluation of Some Natural Polymers as Pharmaceutical Excipient For Sustained Drug Delivery System"

Article *in* International Journal of Humanities and Social Science · January 2021

CITATIONS
3

READS
660

2 authors, including:

Avinash V Dhobale
LSDP college of pharmacy mandavgan pharata
28 PUBLICATIONS **42** CITATIONS

SEE PROFILE

All content following this page was uploaded by Avinash V Dhobale on 10 April 2021.

The user has requested enhancement of the downloaded file.

Back to Academic Year 2021-22

Green Chemistry

IJCRT.ORG ISSN : 2320-2882

INTERNATIONAL JOURNAL OF CREATIVE RESEARCH THOUGHTS (IJCRT)

An International Open Access, Peer-reviewed, Refereed Journal

GREEN CHEMISTRY

[1]YOGITA S.TEMAK, [2]PRAVIN BALASAHEB CHOLKE, [3]Nasir Taboli
[1]ASSISTANT PROFESOR, [2]ASSOCIATED PROFESOR, [3]Student
[1]DBTU,
[2]DBTU,
[3]DBTU

1.Abstract

The object of Green Chemistry is the reduction of chemical pollutants flowing to the environment. The Chemistry and the Environmental Division of EuChem has assumed Green Chemistry as one of its areas of interests, but one question to solve is where Green Chemistry should be placed within the context of Chemistry and Environment. The concept of Green Chemistry, as primarily conceived by Paul Anastas and John Warner, is commonly presented through the twelve principles of Green Chemistry. However, these Twelve principles through fruits of a great intuition and common sense, do not a clear connection between aims, concepts, and related research areas of Green Chemistry. The Twelve unsolved questions are the object of the present article.

2. Introduction

New chemistry is required to improve the economics of chemical manufacturing and to enhance the environmental protection. The green chemistry concept presents an attractive
technology to chemists, researchers, and industrialists for innovative chemistry research and applications .Primarily, green chemistry is characterized as reduction of the environmental damage accompanied by the production of materials and respective minimization and proper disposal of wastes generated during different chemical processes.

According to another definition, green chemistry is a new technique devoted to the synthesis, processing, and application of chemical materials in such manner as to minimize hazards to humankind and the environment.

Numerous new terms have been introduced associated with the concept of “green chemistry,” such as eco-efficiency, sustainable chemistry, atom efficient or atom economy, process intensification and integration inherent safety, product life cycle analysis, ionic liquids, alternate feed stocks, and “Renewable Energy Source.”

Hence, there is an essential need to improve the synthetic and engineering chemistry either by environmental friendly starting materials or by properly designing novel synthesis routes
that reduce the use and generation of toxic substances by using modern energy sources.

Back to Academic Year 2021-22

A Review On: Preservatives Used in Pharmaceuticals and Impacts on Health

© JAN 2022 | IRE Journals | Volume 5 Issue 7 | ISSN: 2456-8880

A Review On: Preservatives Used in Pharmaceuticals and Impacts on Health

SHUBHAM SHITOLE[1], SHWETA SHINDE[2], SANTOSH WAGHMARE[3], HEMANT KAMBLE[4]
[1] *Dept of Pharmaceutics, Loknete Shri Dadapatil Pharate Patil College of Pharmacy, Mandavgan Pharata, Shirur, Pune, Maharashtra.*
[2] *Dept of Pharmaceutical Chemistry, Loknete Shri Dadapatil Pharate Patil College of Pharmacy, Mandavgan Pharata, Shirur, Pune, Maharashtra.*
[3] *Dept of Pharmacology, Loknete Shri Dadapatil Pharate Patil College of Pharmacy, Mandavgan Pharata, Shirur, Pune, Maharashtra.*
[4] *Loknete Shri Dadapatil Pharate Patil College of Pharmacy, Mandavgan Pharata, Shirur, Pune, Maharashtra.*

Abstract- ***For several decades pharmacist have been aware of the need to protect their products against microbial contamination but it is only during the last one or perhaps two decades the serious thought of has been applied to the science of preservation. Preservatives are commonly used as additives in pharmaceutical products, food and cosmetics. Some of the liquid preparations are susceptible to microbial contamination because of the nature of ingredients present in it. Such preparation is protected by preservatives which avoids degradation and alteration of the product. A preservative is a natural or synthetic chemical added to various products which helps to prevent microbial decomposition. Present article deals with the study of ideal properties, classification, mechanism of action, pharmaceutical applications and its impact on health of various preservatives used in pharmaceuticals.***

I. INTRODUCTION

A preservative is a natural or synthetic chemical that is added to products such as foods, pharmaceuticals, paints, biological samples, wood, etc. to prevent decomposition by microbial growth or by undesirable chemical changes. Preservatives are substances that are commonly added to various foods and pharmaceutical products in order to prolong their shelf life. The addition of preservatives to such products, especially to those that have higher water content, is essential for avoiding alteration and degradation by microorganisms during storage. Preservatives are put in foods to inhibit growth of bacteria, yeasts, or molds that can cause disease. Chemical preservation cannot totally keep products from spoiling, but they slow the spoiling process caused by microorganisms. Frozen and canned foods often do not contain any preservatives. Processed Foods are foods that are put through a process to kill harmful bacteria that may form in the food. These processes are supposed to be helpful to the products, but they can also add harmful substances.

When a natural food is processed it may be crushed, heated and have chemicals added to it that may kill all the nutritional value of the food. Then often additives are added to the product to put back some vitamins and nutrients that were lost when the food product was processed. Preservative is added to keep foods fresh and to keep them from spoiling.[1] The first preservatives to be used were vinegar, salt and sugar. Now many of the preservatives are man-made chemicals. Many of the synthetic food and cosmetic additives are considered to be safe, but some of them were found to be carcinogenic and toxic and it is better to limit their use. In general, all synthetic chemical additives and preservatives may be avoided, as many of them have not been properly tested. It happens fairly often that a synthetic chemical additive that has been judged to be safe and has been used for year is later found to be toxic and, in some cases, causes children and adult to die.

All of the chemical food coloring can be potentially harmful, so it is better to avoid all food coloring made from synthetic chemicals. It is important to understand

Back to Academic Year 2021-22

INTERNATIONAL JOURNAL IN PHARMACEUTICAL SCIENCES

Journal Homepage: http://ijpsjournal.com/

OPEN ACCESS

Review Article

An Overview on Artificial Blood

Rajnandini Ahire*, Mukesh Ratnaparkhi, Shital Kshirsagar, Devang Mahale

Department of Pharmaceutics, Marathwada Mitra Mandal's College of Pharmacy, Thergaon (Kalewadi), Pune411033, Maharashtra, India

ARTICLE INFO

Received: 18 Sept. 2021
Revised: 24 Sept. 2021
Accepted: 15 Oct. 2021
Published: 8 Nov. 2021
Keywords:
artificial blood, hemoglobin, hemoglobin-based oxygen carriers, oxygen carrier, Perfluorocarbons, red blood cells
DOI:
10.5281/zenodo.5655215

ABSTRACT

The 21st century is a challenge for people. Population growth, aging population, generation of new infectious agents, and natural disasters are some of the factors that threaten the status quo of blood transfusions. Artificial blood is designed solely to carry oxygen and carbon dioxide into the body. Depending on the type of blood replacer, it can be produced with a variety of techniques using chemical separation, synthetic production, or recombinant biochemical techniques. The technologies focused on the two major alternatives to blood are perfluorocarbons and hemoglobin-based oxygen carriers (HBOCs). With the availability of blood products, the United States alone is expected to generate more than $ 7.6 billion annually.

HISTORY

The concept of artificial blood sounds simple, but it's not. In 1616, William Harvey first described blood circulation, and scientists began to wonder if blood could be taken and replaced with other liquids such as wine and milk. 6 Milk was one of the first ingredients in 1854, and patients used milk to treat cholera in Asia. The patient died of bleeding due to a serious injury and needed a blood exchange. [1] Since World War II, research on blood proposals has been taken seriously in order to overcome large-scale situations such as private sector events. The composition of blood and the function of each substitution are clear except for the mobility of oxygen. [3] They wanted to change a person's blood for a variety of beneficial effects, including healing of illness and personality. In 1667, the first successful human blood transfusion took place.

***Corresponding Author**: Rajnandini Ahire
Address: *Department of Pharmaceutics, Marathwada Mitra Mandal's College of Pharmacy, Thergaon (Kalewadi), Pune-411033, Maharashtra, India*
Email ✉ : _rajnandiniahire99@gmail.com

Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Back to Academic Year 2021-22

Sr. no	Title of the paper	Name of Staff	Department	Name of the journal	ISSN number	Document	Link
1	Face Care Cosmetic-A Review On Herbal Face Toner	Dr. V. M. Satpute	Pharmaceutics	World J Pharm Pharm Sci	ISSN 2277 – 7105	View Document	https://www.wjpr.net/abstract_show/21922
2	Chloroquine and Hydroxychloroquine in Coronavirus Disease-19: The Real Savior or a False-positive Treatment	Dr. V. M. Satpute	Pharmaceutics	JETIR	ISSN 2349-5162	View Document	https://www.google.com/search?q=Chloroquine+and+Hydroxychloroquine+in+Coronavirus+Disease-19%3A+The+Real+Savior+or+a+False-positive+Treatment+research+paper+by+dr.v.m.satpute%3AJETIR&sca_esv=d143a126214438bd&sca_upv=1&rlz=1C1VDKB_en-GBIN1070IN1070&sxsrf=ACQVn08BcAMGn4TbxWJyDzkQHRBmakjxuw%3A1713681337038&ei=ubMkZsz7AY6Yvr0P5MGs0A8&ved=0ahUKEwiMvOi62NKFAxUOjK8BHeQgC_oQ4dUDCBA&uact=5&oq=Chloroquine+and+Hydroxychloroquine+in+Coronavirus+Disease-19%3A+The+Real+Savior+or+a+False-positive+Treatment+research+paper+by+dr.v.m.satpute%3AJETIR&gs_lp=Egxnd3Mtd2l6LXNlcnAikgFDaGxvcm9xdWluZSBhbmQgSHlkcm94eWNobG9yb3F1aW5lIGluIENvcm9uYXZpcnVzIERpc2Vhc2UtMTk6IFRoZSBSZWFsIFNhdmlvciBvciBhIEZhbHNlLXBvc2l0aXZlIFRyZWF0bWVudCByZXNlYXJjaCBwYXBlciBieSBkci52Lm0uc2F0cHV0ZTpKRVRJUkgAUABYAHAAeACQAQCYAQCgAQCqAQC4AQPIAQD4AQGYAgCgAgCYAwCSBwCgBwA&sclient=gws-wiz-serp

3	A Review: Etiology, Pathogenesis And Treatment Of Peptic Ulcer	Dr. H. V. Kamble	Pharmacology	World Journal of Pharmaceutical Research	ISSN 2277 – 7105	View Document	https://scholar.google.co.in/scholar?q=A+Review:+Etiology,+Pathogenesis+And+Treatment+Of+Peptic+Ulcer+by+H.V.+Kamble&hl=en&as_sdt=0&as_vis=1&oi=scholart
4	Formulation and Evaluation of Sustained Release Ibuprofen Tablet	Dr. H. V. Kamble	Pharmacology	International Journal of Scientific Research and Engineering Development	Volume 6 Issue 2	View Document	https://scholar.google.co.in/scholar?q=Formulation+and+Evaluation+of+Sustained+Release+Ibuprofen+Tablet++by+H.V.+Kamble&hl=en&as_sdt=0&as_vis=1&oi=scholart
5	Formulation and Evaluation of Sustained Release Ibuprofen Tablet	Dr. S. A. Waghmare	Pharmaceutical Chemistry	International Journal of Scientific Research and Engineering Development	Volume 6 Issue 2	View Document	https://scholar.google.co.in/scholar?q=Formulation+and+Evaluation+of+Sustained+Release+Ibuprofen+Tablet++by+H.V.+Kamble&hl=en&as_sdt=0&as_vis=1&oi=scholart
6	A Comprehensive Review On Clinical Trials	Dr. H. V. Kamble	Pharmacology	World Journal of Pharmaceutical Research World Journal of Pharmaceutical	ISSN 2277 – 7105	View Document	https://scholar.google.co.in/scholar?q=A+Comprehensive+Review+On+Clinical+Trials+by+H.V.+Kamble&hl=en&as_sdt=0&as_vis=1&oi=scholart

				Research			
7	A Comprehensive Review On Clinical Trials	Dr. S. A. Waghmare	Pharmaceutical Chemistry	World Journal of Pharmaceutical Research World Journal of Pharmaceutical Research	ISSN 2277 – 7105	View Document	https://scholar.google.co.in/scholar?q=A+Comprehensive+Review+On+Clinical+Trials+by+H.V.+Kamble&hl=en&as_sdt=0&as_vis=1&oi=scholart
8	Solid Lipid Nanoparticle (Sln)	Dr. S. A. Waghmare	Pharmaceutical Chemistry	International Research Journal of Modernization in Engineering Technology and Science	ISSN: 2582-5208	View Document	https://www.researchgate.net/publication/283600259_Solid_lipid_nanoparticles_A_promising_drug_delivery_system
9	Novel Maos of Novel Cinnoline Derivatives	Ms. Bhalerao Monali Ramchandra	Pharmaceutical Chemistry	Eur.Chem.Bull	ISSN 2063-5346	View Document	https://www.google.com/search?q=Development+and+validation+of+chromatographic+method+for+analysis+of+some+polycyclic+hydrocarbons+on+soybean+oil+by+ms.+monali+r.+bhalerao&rlz=1C1VDKB_en-GBIN1070IN1070&oq=Development+and+validation+of+chromatographic+method+for+analysis+of+some+polycyclic+hydrocarbons+on+soybean+oil+by+ms.+monali+r.+bhalerao&gs_lcrp=EgZjaHJvbWUyBggAEEUYOTIHCAEQIRiPAjIHCAIQIR

							iPAtIBCTIwMDUzajBqN6gCCLACAQ&sourceid=chrome&ie=UTF-8
10	Development and validation of chromatographic method for analysis of some polycyclic hydrocarbons on soybean oil	Ms. Bhalerao Monali Ramchandra	Pharmaceutical Chemistry	Eur.Chem.Bull	ISSN 2063-5346	View Document	https://www.google.com/search?q=Development+and+validation+of+chromatographic+method+for+analysis+of+some+polycyclic+hydrocarbons+on+soybean+oil+by+ms.+monali+r.+bhalerao&rlz=1C1VDKB_en-GBIN1070IN1070&oq=Development+and+validation+of+chromatographic+method+for+analysis+of+some+polycyclic+hydrocarbons+on+soybean+oil+by+ms.+monali+r.+bhalerao&gs_lcrp=EgZjaHJvbWUyBggAEEUYOTIHCAEQIRiPAjIHCAIQIRiPAtIBCTIwMDUzajBqN6gCCLACAQ&sourceid=chrome&ie=UTF-8
11	A Review on Beta Vulgaris (Beet Root)	Mr. A. S. Gaikwad	Pharmaceutical Chemistry	PENSEE	ISSN NO: 0031-4773	View Document	https://www.researchgate.net/publication/374813414_A_Review_on_Beta_Vulgaris_Beet_Roo
12	Microsponge Drug Delivery System - A Review	Mr. A. S. Gaikwad	Pharmaceutical Chemistry	PENSEE	ISSN NO: 0031-4773	View Document	https://www.researchgate.net/publication/374813521_Microsponge_drug_delivery_Scopus
13	Development And Validation Of RP HPLC Method For Simultaneous Estimation Of Zaltoprofen And	Mr. A. S. Gaikwad	Pharmaceutical Chemistry	International journal Of Current Research	ISSN: 0975-833	View Document	https://www.researchgate.net/publication/374813419_DEVELOPMENT_AND_VALIDATION_OF_RP_ZALTOPROFEN_AND_PARACETAMOL

	Paracetamol In Bulk And Tablet Formulation						
14	Synthesis of Cyclopyrrolidine Clubbed with Oxadiazole Bases and Evaluation of their Anti-Diabetic Activity through in vivo Model	Ms. M. T. Salve		Indian Journal of Pharmaceutical Education and Research	Vol 57, Issue 3(Suppl), Ju	View Document	https://www.ijper.org/article/2084
15	Novel Drug Delivery System : Review Article	Ms. S. S. Khillare	Pharmaceutics	WJPR	ISSN: 2277-7105	View Document	https://www.wjpr.net/abstract_show/22172
16	A Review on Nanotechnology in Cancer Treatment	Mr. M. M.Garje	Pharmaceutical Chemistry	IJSRT	Vol 7: Isue 9	View Document	https://www.google.com/search?q=A+Review+on+Nanotechnology+in+Cancer+Treatment+by+mr.m.m.+garje%3AIJCRT&sca_esv=d143a126214438bd&sca_upv=1&rlz=1C1VDKB_en-GBIN1070IN1070&biw=1366&bih=641&sxsrf=ACQVn08bbuYmF_F1tdObiUbhEUGOJQqEzg%3A1713682995582&ei=M7okZsmQI73fseMPxqWQeA&ved=0ahUKEwiJ9tXR3tKFAxW9b2wGHcYSBA8Q4dUDCBA&uact=5&oq=A+Review+on+Nanotechnology+in+Cancer+Treatment+by+mr.m.m.+garje%3AIJCRT&gs_lp=Egxnd3Mtd2l6LXNlcnAiRUEgU

							mV2aWV3IG9uIE5hbm90ZWNobm9sb2d5IGluIENhbmNlciBUcmVhdG1lbnQgYnkgbXIubS5tLiBnYXJqZTpJSkNSVEgAUABYAHAAeACQAQCYAQCgAQCqAQC4AQPIAQD4AQGYAgCgAgCYAwCSBwCgBwA&sclient=gws-wiz-serp
17	Face Care Cosmetic-A Review On Herbal Face Toner	Dr. H. V. Kamble	Pharmacology	World J Pharm Pharm Sci	ISSN 2277 – 7105	View Document	https://www.wjpr.net/abstract_show/21922
18	Face Care Cosmetic-A Review On Herbal Face Toner	**Mr. S. A. Waghmare**	Pharmaceutical Chemistry	World J Pharm Pharm Sci	ISSN 2277 – 7105	View Document	https://www.wjpr.net/abstract_show/21922
19	Novel Drug Delivery System : Review Article	Dr. H. V. Kamble	Pharmacology	WJPR	ISSN: 2277-7105	View Document	https://www.wjpr.net/abstract_show/22172
20	Novel Drug Delivery System : Review Article	**Mr. S. A. Waghmare**	Pharmaceutical Chemistry	WJPR	ISSN: 2277-7105	View Document	https://www.wjpr.net/abstract_show/22172

Back to Academic Year 2022-23

WORLD JOURNAL OF PHARMACEUTICAL RESEARCH
SJIF Impact Factor 8.084
Volume 12, Issue 5, 59-70. Review Article ISSN 2277– 7105

FACE CARE COSMETIC-A REVIEW ON HERBAL FACE TONER

Suraj Agaldare*[1], Dr. Vivek Satpute[2], Mr. Santosh A. Waghmare[3] and Dr. Hemant V. Kamble[4]

[1]Student Department of Pharmaceutics, Loknete, Shri Dadapatil Pharate College of Pharmacy, Mandavgan Pharata, Shirur, Pune, Maharastra, India.
[2,3]Professor, Department of Pharmaceutics, Loknete, Shri Dadapatil Pharate College of Pharmacy, Mandavgan Pharata, Shirur, Pune, Maharastra, India.
[4]Principle, Department of Pharmaceutics, Loknete, Shri Dadapatil Pharate College of Pharmacy, Mandavgan Pharata, Shirur, Pune, Maharastra, India.

Article Received on
24 February 2023,
Revised on 16 March 2023,
Accepted on 05 April 2023
DOI: 10.20959/wjpr20235-27485

***Corresponding Author**
Suraj Agaldare
Student Department of Pharmaceutics, Loknete, Shri Dadapatil Pharate College of Pharmacy, Mandavgan Pharata, Shirur, Pune, Maharastra, India.

ABSTRACT

Natural remedies have lesser side effects, secure and also aceeptable than chemical ones. In the world market, formulations with natural ingredients have more accesible. For delivering the drug imediately to the site of action, which gives prolonged action is the benefits of topical drug delivery system. Skin is the main path of delivery of drug in TDDS. The ingredients are easily available which are being used. They are not only easily available but also has nutritional value from topical point of view and more economical. To formulate and evaluate the formulation is the motive of this project. It is natural and safe herbal preparation which gives calming, soothing and astringent effect on the face. The natural ingredients like aloe vera, cucumber also the pippermint, lemmongrass and rose water used in the formulation. It having ability to reduce the facial irritation as well as to enhance beauty. We can use it in our daily busy schedule. Face toner is estimated for its physicochemical properties, surface tension, pH and stability. Most popular advantages of herbal cosmetics are, they are non toxic in nature and they having tendancy to reduce allergic reactions. The main reason behind this study, we found good properties of the face toner.

KEYWORDS: Toner, Aloe vera, Cucumber, Pipper mint, Lemmon grass.

Back to Academic Year 2022-23

© 2023 JETIR April 2023, Volume 10, Issue 4 www.jetir.org (ISSN-2349-5162)

JETIR.ORG ISSN: 2349-5162 | ESTD Year : 2014 | Monthly Issue

JOURNAL OF EMERGING TECHNOLOGIES AND INNOVATIVE RESEARCH (JETIR)

An International Scholarly Open Access, Peer-reviewed, Refereed Journal

Chloroquine and Hydroxychloroquine in Coronavirus Disease-19: The Real Savior or a False-positive Testament

Shubham A. Patil* Student, Department of Pharmaceutics, Loknete Shri Dadapatil Pharate College of Pharmacy, Mandavgan Pharata, Shirur, Pune, Maharashtra, India *Corresponding Author.

Prasad A. Mhaske ,Student, Department of Pharmaceutics, Loknete Shri Dadapatil Pharate College of Pharmacy, Mandavgan Pharata, Shirur, Pune, Maharashtra, India.

Dr. Vivek M. Satpute Professor, Department of Pharmaceutics, Loknete Shri Dadapatil Pharate College of Pharmacy, Mandavgan Pharata, Shirur, Pune, Maharashtra, India.

Mr Santosh A. Waghmare Professor, Department of Pharmaceutics, Loknete Shri Dadapatil Pharate College of Pharmacy, Mandavgan Pharata, Shirur, Pune, Maharashtra, India.

Dr. Hemant V. Kamble Principal, Department of Pharmaceutics, Loknete Shri Dadapatil Pharate College of Pharmacy, Mandavgan Pharata, Shirur, Pune, Maharashtra, India.

Abstract :-

Objective: A novel coronavirus disease (COVID-19) has spread all around the world. The progression from initial signs to a diagnosis of acute respiratory failure is usually related to spontaneous cytokine production. There is a growing need to classify appropriate medicines for treatment care. The inhibitory effect of chloroquine (CQ) is potential. However, CQ can lead to serious side effects. Various studies recommend hydroxychloroquine (HCO) have similar antiviral effect as of CQ and maybe a better therapeutic solution. Therefore, we aim to explore the mechanism by which HCQ can inhibit replication of coronavirus. Materials and Methods: A retrospective study was carried out using online databases from 2003 to 2020, Results: The obtained results showed that HCQ can inhibit viral replication and entry inside the cell through raising lysosomal pH and hinding to specific receptors on cells, thereby, preventing viral entry. Conclusion: HCQ has a better safety profile than CQ and also modulates cytokine syndrome. However, further studies are needed to explore this mechanism.

Key words: Chloroquine, coronavirus disease-19, hydroxychloroquine, severe acute respiratory syndrome .

Back to Academic Year 2022-23

WORLD JOURNAL OF PHARMACEUTICAL RESEARCH
SJIF Impact Factor 8.084
Volume 12, Issue 19, 36-47. Review Article ISSN 2277– 7105

A REVIEW: ETIOLOGY, PATHOGENESIS AND TREATMENT OF PEPTIC ULCER

Ms. Sonal S. Ithape*, Dr. Hemant V. Kamble and Mr. Santosh A. Waghmare

India.

Article Received on
13 Sept. 2023,
Revised on 03 October 2023,
Accepted on 23 October 2023
DOI: 10.20959/wjpr202319-29270

***Corresponding Author**
Ms. Sonal S. Ithape
India.

ABSTRACT

Peptic ulcer disease, which includes both stomach and duodenal ulcers, accounts for a significant portion of people seeking surgical advice around the world. A peptic ulcer is a lesion that develops on the stomach or duodenal lining. "Gastric ulcers" and "duodenal ulcers" are the two most prevalent peptic ulcer kinds. Peptic ulcers are caused by an imbalance between aggressive factors like hydrochloric acid (HCL), pepsin, refluxed bile, leukotrienes (LTs), reactive oxygen species (ROS), and defensive factors like the mucus-bicarbonate barrier, prostaglandins (PGs), mucosal blood flow, cell renewal and migration, nonenzymatic and enzymatic antioxidants. The most common causes of peptic ulcer disease are H. pylori infection and the use of nonsteroidal anti-inflammatory medications (NSAIDs). In addition, a variety of variables are implicated in the pathogenesis of gastric ulcer, including bacterial infection (Helicobacter pylori), certain drugs (NSAID), chemicals (Hcl/ethanol), and stomach cancer, with minor factors including stress, smoking, spicy food, and nutritional deficiencies.

KEYWORDS: Peptic Ulcer, Acid Secretion, NSAID, Helicobactor Pylori, Proton Pump Inhibitor.

INTRODUCTION

Peptic ulcer occurs in that part of the gastrointestinal tract (G.I.T.) which is exposed to gastric acid and pepsin, i.e. the stomach and duodenum. The etiology of peptic ulcer is not clearly known. It results probably due to an imbalance between the aggressive (acid, pepsin, bile and H. pylori) and the defensive (gastric mucus and bicarbonate secretion, prostaglandins, nitric oxide, high mucosal blood flow, innate resistance of the mucosal cells) factors. A variety of psychosomatic, humoral and vascular derangements have been implicated and the causative

Back to Academic Year 2022-23

Formulation and Evaluation of Sustained Release Ibuprofen Tablet

International Journal of Scientific Research and Engineering Development-– Volume 6 Issue 2, Mar-Apr 2023

Available at www.ijsred.com

RESEARCH ARTICLE OPEN ACCESS

Formulation and Evaluation of Sustained Release Ibuprofen Tablet

Sahil A Dalavi*, Mr Santosh A Waghmare**, Dr Hemant V Kamble**

**(Student, Department of Pharmaceutical Chemistry, Loknete Shri Dadapatil Pharate Collage of Pharmacy , A/p- MandavganPharata,Tal-Shirur , Dist- Pune*

Email:sahildalavi4412@gmail.com

*** (Professor, Loknete Shri Dadapatil Pharate Collage of Pharmacy , A/p- MandavganPharata,Tal-Shirur , Dist- Pune*

*** (Principle,Loknete Shri Dadapatil Pharate Collage of Pharmacy , A/p- MandavganPharata,Tal-Shirur , Dist- Pune*

----------**********----------

Abstract:

In this study of Ibuprofen it is an (NSAIDs) non steroidal anti- inflammatory drug and used as analgesic & anti-inflammatory drug. It can be also used in the treatment of rheumatoid arthritis, osteoarthritis, and primary dysmenorrheal. Ibuprofen is absorbed rapidly, it has low aqueous solubility so it also lowers the dissolution profile of drug.The main aim of proposed work was to develop Ibuprofen tablets, sustained release dosage form, for the treatment inflammation and pain in the body. Ibuprofen is used to reduce fever and treat pain or inflammation caused by many conditions such as headache, toothache, back pain, arthritis, menstrual cramps, or minor injury. Sustained release formulation is the drug delivery system that designed to achieve a prolonged therapeutic effect by continuously releasing medication over an extended period of time after administration of single dose. The tablets were prepared by direct compression method using Hydroxy Propylemethyl Cellulose HPMC(HV),HPMC(LV) and talc.Tablets blends were evaluated for bulk density, tapped density,Carr's Index compressibility index and angle of repose shows satisfactory results.The results of all these tests were found to be satisfactory. The in vitro dissolution study was carried out for 12 hours using paddle method in phosphate buffer (pH7.4) as dissolution media. Formulation F4 shows – of drug release at the end of 12 hours.

Keywords: Ibuprofen, Sustained Release, Dissolution Rate

Keywords — Clinical studies, target discovery and validation, lead optimization, new medication.

----------**********----------

INTRODUCTION

1. **Oral Drug Delivery System**

Oral route of drug delivery is the most preferred route of administration All other routes of drug delivery because of ease of administration,patient compliance and flexible design of dosage form. An oral drug delivery system providing a uniform drug delivery can only partly satisfy therapeutic and biopharmaceutical needs, as it doesn't take in to account the site specific absorption rates within the gastrointestinal tract (GIT). Therefore there is a need of developing drug delivery system that release the drug at the right time, at the specific site and with the desired rate.[1,2]

 ©IJSRED:All Rights are Reserved

Back to Academic Year 2022-23

WORLD JOURNAL OF PHARMACEUTICAL RESEARCH
SJIF Impact Factor 8.084
Volume 12, Issue 11, 1368-1379. Review Article ISSN 2277– 7105

A COMPREHENSIVE REVIEW ON CLINICAL TRIALS

Dhananjay Narsale*, Hemant Kamble and Santosh Waghmare

Loknete Shree Dadapatil Pharate College of Pharmacy, at Post Mandavgan Pharata, Ta: Shirur, Dist: Pune. Pincode: 412211.

Article Received on
19 May 2023,
Revised on 09 June 2023,
Accepted on 29 June 2023
DOI: 10.20959/wjpr202311-28892

***Corresponding Author**
Dhananjay Narsale
Loknete Shree Dadapatil Pharate College of Pharmacy, at Post Mandavgan Pharata, Ta: Shirur, Dist: Pune. Pincode: 412211.

ABSTRACT

A clinical trial is a type of research that looks at specific health issues with human volunteers. The quickest and safest way to find a treatment that works for people and a way to improve health is through carefully conducted clinical trials. Under controlled conditions, investigational trials determine whether experimental treatments or new ways to use known therapies are safe and effective. Health issues are addressed in large populations or natural populations through observational trials. A crucial and highly specialized type of biological assay, clinical trials aim to measure therapeutic efficacy. Clinical pharmacologists conduct research on human volunteers for phase I pharmacokinetics, safety, and gross effects. If the drug passes the test, it moves on to phase II testing, where clinical pharmacologists study the pharmacokinetics, safety, and therapeutic efficacy of selected patients. If the drug passes, clinical investigators now study hundreds of selected patients in phase III, primarily for safety and therapeutic efficacy. The drug is now approved and can be sold if this is passed. Physicians from a variety of hospitals and clinics still provide feedback on the drug's efficacy, ADR, and marketing after phase IV.

KEYWORDS:- Clinical trials, Phases of clinical trials, ICH guidelines, IND, NDA.

INTRODUCTION

A clinical trial is a systematic process that is intended to find out the safety and efficacy of a drug/device in treating/preventing/diagnosing a disease or a medical condition.[1] For any new drug to enter in clinical trial, it must pass preclinical studies. Preclinical studies involve in vitro (i.e. test-tube or Laboratory) studies and trials on animal populations. Wide range of

Back to Academic Year 2022-23

Solid Lipid Nanoparticle (Sln)

INTERNATIONAL RESEARCH JOURNAL OF PHARMACY
www.irjponline.com ISSN 2230 – 8407
Review Article

SOLID LIPID NANOPARTICLES: A PROMISING DRUG DELIVERY SYSTEM

Waghmare A. S.*, Grampurohit N.D., Gadhave M.V., Gaikwad D.D., Jadhav S.L
Department of Pharmaceutics, Vishal Institute of Pharmaceutical Education & Research, Ale, Pune, Maharashtra, India

Article Received on: 17/02/12 Revised on: 29/03/12 Approved for publication: 05/04/12

*Email: amolwaghmare5@gmail.com

ABSTRACT
Solid lipid nanoparticles are at the forefront of the rapidly developing field of nanotechnology with several potential applications in drug delivery, research and clinical medicine, as well as in other varied sciences. Solid lipid nanoparticles (SLNs) technology represents a promising new approach to lipophilic drug delivery. Nanoparticles are solid colloidal particles ranging in size from 1 to 1000 nm and composed of macromolecular material. The biodegradable and bioacceptable nature of SLNs makes them less toxic as compared to polymeric nanoparticles. SLNs can also be used to improve the bioavalability of drugs. In this present review this new approach is discussed in terms of their advantages, characterization, pharmacokinetic studies, in-vivo studies, in-vitro studies, and special features.

KEYWORDS: Nanotechnology, Solid lipid nanoparticle, Colloidal carriers, Bioavailability enhancement.

INTRODUCTION

A high potential for drug delivery has been attributed to particulate drug carriers, especially small particles such as microparticles and colloidal system in nanometer range.[1] Nano particulate drug delivery system may offer plenty of advantages over conventional dosage forms which include improved, reduced toxicity, enhancedbio distribution and improved patient compliance[2]. Colloidal particles ranging in size between 10 and 1000 nm are known as nanoparticles. They are manufactured from synthetic/natural polymers and ideally suited to optimize drug delivery and reduce toxicity. Over the years, they have emerged as a variable substitute to liposomes as drug carriers. The successful implementation of nanoparticles for drug delivery depends on their ability to penetrate through several anatomical barriers, sustained release of their contents and their stability in the nanometer size. However, the scarcity of safe polymers with regulatory approval and their high cost have limited the wide spread application of nanoparticles to clinical medicine [3].To overcome these limitations of polymeric nanoparticles, lipids have been put forward as an alternative carrier, particularly for lipophilic pharmaceuticals. These lipid nanoparticles are known as solid lipid nanoparticles (SLNs), which are attracting wide attention of formulators world-wide.[4] SLNs are colloidal carriers developed in the last decade as an alternative
system to the existing traditional carriers (emulsions, liposomes and polymeric nanoparticles). They are a new generation of submicron-sized lipid emulsions where the liquid lipid (oil) has been substituted by a solid lipid. SLN offer unique properties such as small size, large surface area, high drug loading and the interaction of phases at the interfaces, and are attractive for their potential to improve performance of pharmaceuticals, neutraceuticals and other materials.[5].

Nanotechnology

Nanotechnology is the science of matter and material that deal with the particle size in nanometers. These are small colloidal particles that are made of nonbiodegradable & biodegradable polymers and their diameter is around 200nm[6]

Nanoparticles

Nanoparticles are solid polymeric, submicronic colloidal system range between 5-300nm consisting of macromolecular substances that vary in size 10nm to 100nm. The drug of interest is dissolved, entrapped, adsorbed, attached or encapsulated into the nanoparticle matrix.[7] In recent years, significant effort has been devoted to develop nanotechnology for drug delivery, since it offers a suitable means of delivering small molecular weight drugs, as well as macromolecules such as proteins, peptides or genes to cells and tissue. Nanoparticles hold promise for peroral drug delivery, which represents so far the most common and convenient route of administration.

The advantages of using nano particles for nanoparticles loaded with drugs, because of their small size can penetrate through small capillaries and are taken up by cells and allow the drug release at right rate and dose at specific sites in the body for a certain time to release the accurate delivery, which enhances the therapeutic effect and reduces the toxicity and side effects. The use of biodegradable materials for nanoparticles preparation allows sustained release within the target site over a period of days or even weeks. The advantages of nanoparticles as drug delivery system are that they are biodegradable, non-toxic, and capable of being stored for longer periods. In addition, their potential uptake as well as their stability in the GIT that indicates that nanoparticles are expected to be the promising carriers for the transport of drugs. These attributes make nanoparticles more suitable for the purpose of sustained release and improvement of bioavailability[8]

SOLID LIQUID NANOPARTICLES

Solid lipid nanoparticles (SLNs) are considered to be the most effective lipid based colloidal carriers (fig 1), introduced in early nineties. This is the one of the most popular approaches to improve the oral bioavailability of the poorly water soluble drugs. SLNs are in the range of submicron size (50-1000 nm) and are composed of physiologically tolerated lipid components which are in solid state at room temperature[9,10] They have many advantages such as good biocompatibility, low toxicity andlipophillic drugs are better delivered by solid lipid nanoparticles and the

Back to Academic Year 2022-23

ISSN 2063-5346

NOVEL MAOS OF NOVEL CINNOLINE DERIVATIVES

Atul Baravkar[1*], Dipali Kadam[2], Sheetal Gaikwad[3], Nitin Shinde[4], Sujata Veer[5], Amit Lunkad[6], Vitthal Chopade[7], Vishnu Neharkar[8], Makarand Puri[9], Rajanikant Kakade[10], Nilesh Jadhav[11], Shyam Panga[11], Vijay M. Kale[12], Yojana Kunjir[12], Sachin Vijapure[13], Reshma Devkate[14], Priti Kolpe[14], Sonali Pawar[15], Baliram Sarvade[16], Sachin Anbhule[17], Raju Kawade[18], Shaikh Sana M Jafar Shaikh[18], Rahul Mohan[18], Monali Bhalerao[19], Meghana Muley[20], Gaffar Sayyad[21]

Article History: **Received:** 01.02.2023 **Revised:** 07.03.2023 **Accepted:** 10.04.2023

Abstract

An efficient and green novel microwave assisted organic synthesis (MAOS) method has been developed for dinitro cinnoline derivatives with better yields. The framework of these derivatives was constructed from dinitrophenyl arylethylidene hydrazines. Tetrabutylammonium bromide (TBAB) was used as a phase transfer catalyst (PTC), potassium carbonate as an inexpensive and efficient catalyst and water as solvent due to its polarity which helps to increase the temperature substantially. This methodology features a simple, environmentally friendly approach, employing water as a green solvent and using a one-pot reaction. The use of microwave increases the rate of reaction and it was observed that dinitro arylcinnolines can be synthesised in 8-12 min of microwave irradiation compared to conventional thermal heating protocol which requires more than 2 h. Spectral data confirms the identity of synthesized derivatives and satisfactory yields are obtained by this process.

[1]Shardabai Pawar Institute of Pharmaceutical Sciences and Research, Baramati, Pune, India. 413115
[2]K. K. Wagh College of Pharmacy, Nashik, India. 422006
[3]Samarth Institute of Pharmacy, Belhe, Pune, India. 412410
[4]Shardabai Pawar Mahila Mahavidyalaya, Baramati, Pune, India. 413115
[5]Saikrupa Institute of Pharmacy, Ghargaon, Ahmednagar, India. 413728
[6]Sitabai Thite College of Pharmacy, Shirur, Pune, India. 412210
[7]Modern College of Pharmacy, Pune, India. 411044
[8]Rasiklal M. Dhariwal Institute of Pharmaceutical Education and Research, Pune, India. 411019
[9]School of Pharmacy, Vishwakarma University, Pune, India. 411048
[10]Siddhi's Institute of Pharmacy, Nandagon, India. 421401
[11]Dr. N. J. Paulbudhe College of Pharmacy, Ahmednagar, India. 414003
[12]Mahadev Kanchan College of Pharmaceutical Education and Research, Uruli Kanchan, Pune, India. 412202
[13]Sarsam College of Pharmacy, Palashiwadi, Pune, India. 413110
[14]Institute of Pharmaceutical Sciences and Research for Girls, Bhigwan, Pune, India. 413130
[15]Vishal Institute of Pharmaceutical Education and Research, Ale, Junnar, India. 412211
[16]Padmini College of Pharmacy, Dighanchi, Sangali, India. 415315
[17]Parikrama GOI College of Pharmacy, Kashti, Ahmednagar, India. 414701
[18]Nandkumar Shinde College of Pharmacy, Vaijapur, Aurangabad, India. 413701
[19]LSDP College of Pharmacy, Mandavgan Pharata, Shirur, India. 412211
[20]Delight College of Pharmacy, Koregaon Bhima, Pune, India. 412216
[21]SAJVPM's College of Pharmaceutical Science and Research Center, Beed, India. 431122
Corresponding author: atul200678@gmail.com

DOI: 10.31838/ecb/2023.12.s1.118

Back to Academic Year 2022-23

Development and validation of chromatographic method for analysis of some polycyclic hydrocarbons on soybean oil



ISSN 2063-5346

DEVELOPMENT AND VALIDATION OF CHROMATOGRAPHIC METHOD FOR ANALYSIS OF SOME POLYCYCLIC HYDROCARBONS IN SOYBEAN OIL

Atul Baravkar[1*], Deepali Kadam[2], Monali Bhalerao[3]

Article History: Received: 01.02.2023 **Revised:** 07.03.2023 **Accepted:** 10.04.2023

Abstract

Some polycyclic hydrocarbons (PHs) present in almost all edible oils are carcinogenic, teratogenic, neurotoxic and mutagenic at certain levels. There are total 16 PHs which are present in various edible oils. Soybean oil is one of most widely and routinely used for cooking food by all over population. Hence the detection of PHs levels in various soybean oil brands available in market is necessary. Many analytical methods are available for the same but these methods are time consuming and secondly quantity of organic solvents required is also higher. For example HPLC can be used for analysis of PHs in edible oils but it takes more time for analysis. Hence a novel method is developed using supercritical fluid chromatography for analysis of some PHs in soybean oil which reduces time by 5 times as compared to normal HPLC method and also requires less amount of organic solvents for analysis because of the lower viscosity and higher diffusivity in the mobile phases of SFC. The method is validated using various validation parameters and statistically proved using ANOVA.

KEY WORDS- Polycyclic hydrocarbons, Soybean oil, HPLC, SFC, Cancer.

[1]Shardabai Pawar Institute of Pharmaceutical Sciences and Research, Baramati, Pune, India. 413115

[2]K. K. Wagh College of Pharmacy, Nashik, India. 422006

[3]LSDP College of Pharmacy, Mandavgan Pharata, Shirur, Pune, India. 412211

Corresponding author- atul200678@gmail.com

DOI: 10.31838/ecb/2023.12.s1.087

Back to Academic Year 2022-23

A Review on Beta Vulgaris (Beet Root)

A Review on Beta Vulgaris (Beet Root)

Bharati R Hole*[1], Annasaheb S Gaikwad[2], Ankita A Giramkar[3], Dr. Deepak G Umalkar[4]

Department of Quality Assurance Technique,
[1, 3, 4,] Saikrupa Institute of Pharmacy, Ghargaon, DBATU, M.S. India.
[2] Mangaldeep Institute of Pharmacy, Aurangabad. DBATU, M.S. India.

Correspondence address –
Bharati Raju Hole
Saikrupa Institute of Pharmacy, Ghargaon.
Email address- bharatihole1997@gmail.com

Back to Academic Year 2022-23

PENSEE ISSN NO: 0031-4773

Microsponge Drug Delivery System - A Review

Ankita A Giramkar *, Annasaheb S Gaikwad, Bharati R Hole, Vikram J Pansare

Department of Quality Assurance Technique,
[1,3,4,] Saikrupa Institute of Pharmacy, Ghargaon, DBATU, M.S. India.
[2] Mangaldeep Institute of Pharmacy, Aurangabad. DBATU, M.S. India.

Correspondence Address –
Ankita A Giramkar
Saikrupa Instituteof Pharmacy, Ghargaon.
Email address- giramkarankita95@gmail.com

Back to Academic Year 2022-23

Development And Validation Of RP HPLC Method For Simultaneous Estimation Of Zaltoprofen And Paracetamol In Bulk And Tablet Formulation

ISSN: 0975-833X

Availableonlineathttp://www.journalcra.com

International Journal of Current Research
Vol. 11, Issue, 05, pp.3572-3576, May, 2019

DOI: https://doi.org/10.24941/ijcr.35203.05.2019

INTERNATIONAL JOURNAL OF CURRENT RESEARCH

RESEARCH ARTICLE

DEVELOPMENT AND VALIDATION OF RP-HPLC METHOD FOR SIMULTANEOUS ESTIMATION OF ZALTOPROFEN AND PARACETAMOL IN BULK AND TABLET FORMULATION

***Annasaheb S Gaikwad, Yogesh R Thombare, Nisha S Mhaske and Mahendra B Datir**

Department of Quality Assurance Technique, Pravara Rural Education Society's College of Pharmacy, Sinnar , Dist- Nashik, M.S. India, Sp, Pune University, Pune, M.S. India

ARTICLEINFO

Article History:
Received 19th February, 2019
Received in revised form
25th March, 2019
Accepted 15th April, 2019
Published online 30th May, 2019

Key Words:
Zaltoprofen, Paracetamol,
RP-HPLC Method,
ICH Guideline.

****Corresponding author:***

ABSTRACT

A simple, sensitive, linear, precise and accurate RP-HPLC method for simultaneous estimation of Zaltoprofen and Paracetamol in bulk and tablet formulation as developed and validated. Chromatographic conditions used are stationary phase Grace C^{18} column (250mm × 4.6mm, 5µ particle size. The mobile phase Methanol: Phosphate buffer (PH 3.0) in the ratio 75:25 v/v and flow rate was maintained 0.8ml/min, detection wavelength was 241nm. The retention times were 3.101min and 5.838min for Zaltoprofen and Paracetamol respectively. Calibration plot were linear R^2=0.9994 over the concentration range 10-18µg/ml for Zaltoprofen, R^2 = 0.9994 for the Paracetamol 40-72µg/ml. No interference from any component of pharmaceutical dosage form was observed. The proposed method has been validated as per ICH guidelines, validation studies revealed that method id specific, rapid, reliable and reproducible. The developed method successfully employed for routine quality control analysis in the combined pharmaceutical dosage form.

Copyright © 2019, Annasaheb S Gaikwad et al. *This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.*

Citation: Annasaheb S Gaikwad, Yogesh R Thombare, Nisha S Mhaske and Mahendra B Datir. **2019.** "Development and Validation of RP-HPLC Method for Simultaneous Estimation of Zaltoprofen and Paracetamol in Bulk and Tablet Formulation", *International Journal of Current Research*, 11, (05), 3572-3576.

INTRODUCTION

Zaltoprofen 2-(10, 11-dihydro-10-oxdibenzo (b, f) thiepin-2-yl propionic acid is a potent NSAID. Zaltoprofen is a preferential COX-2 inhibitor and selectively inhibits prostaglandin E_2 (PGE_2) production at inflammatory sites. Paracetamol is 4-hydroxy phenyl acetamide. The central analgesic action of paracetamol is like aspirin, i.e. it raise pain threshold, but has weak peripheral anti-inflammatory action. It is poor inhibitor of PG synthesis in peripheral tissues, but more active on COX inhibitor in the brain.

The ability of paracetamol to inhibit COX-3 could also account for its analgesic, antipyretic action. The combined paracetamol treatment may increase the effect and decrease the dose dependent side effect of NSAIDs and combination of Zaltoprofen with Paracetamol will be potent analgesic and anti-inflammatory drug for future in the pain management. ZAL is marketed in the combination with PCM under the trade name ZOTT ® P by Aeon Formulations Pvt. Ltd. Literature reveals that there are many methods for the individual determination of ZAL and PCM; but few methods are cited for determination of combined dosage form so, it was proposed to develop an economical, rapid and simple RP-HPLC method for simultaneous estimation of these drugs in combined dosage form.

MATERIAL OF METHODS

Chemicals: HPLC grade Methanol, HPLC grade Water, Potassium dihydrogen phosphate AR grade. All other chemicals were of analytical grade.

Instrumentation:HPLC3000 series instrument, P-3000-M Reciprocating pump 40M pal. RP-HPLC Binary gradient system with grace C_{18}column (250mm × 4.6 mm id, particle size 5µ) equipped with UV 3000 –M series detector use, Wenser high precision balance (PGB 100).

Chromatographic Conditions: The mobile phase ratio was optimized in isocratic mode for analysis of Zaltoprofen and Paracetamol. Different ratio was studied such as, 70:30, 80:20, 75:25. Of Methanol: Phosphate Buffer for Zaltoprofen and Paracetamol. The final mobile consisted 75:25 and mobile phase was clarified by filtration through nylon filter paper with pore size 0.45µm and degassed through sonicator then pumped at flow rate 08ml/min, in gradient mode on grace C_{18} column. The peak response was monitored at 241nm wave length. The sample solution was injected (20µl) HPLC system and data was acquired LC solution workstation software.

Preparation of Standard Solution: Weigh accurately 10mg of Zaltoprofen and Paracetamol was transferred into 10ml volumetric flask it was dissolved with methanol from this

Back to Academic Year 2022-23

Synthesis of Cyclopyrrolidine Clubbed with Oxadiazole Bases and Evaluation of their Anti-Diabetic Activity through in vivo Model

Ind. J. Pharm. Edu. Res., 2023; 57(3s):s749-s757.
https://www.ijper.org

Original Article

Synthesis of Cyclopyrrolidine Clubbed with Oxadiazole Bases and Evaluation of their Anti-Diabetic Activity through *in vivo* Model

Salve Megha Tukaram*, Jadhav Shailaja

Department of Pharmaceutical Chemistry, Modern College of Pharmacy, Nigdi, Pune, Maharastra, INDIA.

ABSTRACT

Background and Aim: Based on inhibitors of DPP-IV, there is a fantastic method for creating anti-diabetic medications. These inhibitors regulate diabetic patients' blood sugar levels to prevent difficulties with their health. In the current work, we created a brand-new series of compounds Cyclopyrrolidine clubbed with oxadiazole bases. **Materials and Methods:** Cyclopyrrolidine clubbed with oxadiazole bases (B-1 to B-16) were synthesized and characterized through IR, NMR, mass spectrometry, and elemental analysis. Docking studies were performed to assess interactions and binding modes of synthesized hits at the binding site of receptor DPP-4 (PDB 3W2T). Using vildagliptin as a standard drug, six of the synthesized compounds were tested for their antidiabetic activity in diabetic rats induced with HFD-STZ-Nicotinamide. **Results:** The results showed that compound B-XIV (220*4.56B) resulted in the greatest reduction in blood glucose level from all synthesized compounds compared to that of vildagliptin (215*7.52B) in HFD-STZ-Nicotinamide. Other compounds showed moderate to good antihyperglycemic activity. **Conclusion:** From he presents work it can be concluded that synthesized compounds possess good DPP-IV inhibitory activity. Compounds containing electron-withdrawing groups (chlorine, nitro, methoxy) were displayed a good anti-diabetic effect than electron-donating groups (methyl, hydroxyl). Oxadiazole derivatives could be used for further development to obtain more promising drug candidates.

Keywords: Furadiazole, Hyperglycaemic, HFD-STZ-Nicotinamide.

Correspondence:
Dr. Megha Tukaram Salve Jr,
Department of Pharmaceutical Chemistry, Modern College of Pharmacy, Nigdi, Pune-411044, Maharashtra, INDIA.
Email: meghasalve77@gmail.com

Received: 02-10-2022;
Revised: 30-03-2023;
Accepted: 08-06-2023.

INTRODUCTION

Since the early 1970s, 1,2,4-oxadiazole heterocycles have been extensively investigated, yielding a variety of bioactive compounds, including anti-cancers,[1] anti-inflammatory,[2] anticonvulsant,[3] antiviral,[4] antibacterial,[5] antifungal,[6] anti-depressant,[7] anti-angiogenic,[8] analgesic,[9] anti-insomnia, anti-oedema,[10] anti-parasitic,[11] anti-Alzheimer[12] and anti-diabetic.[13] In addition, they showed inhibition of dipeptidyl peptidase IV,[14] α-Glucosidase,[15] α-Amylase,[16] Protein tyrosine phosphatase 1B.[17] A number of 1,2,4-oxadiazoles were screened as GLP19 (Glucagon-Like Peptide 1),[18] PPAR (peroxisome proliferator-activated receptor) alpha/gamma agonists,[19] GPR40 (G-Protein Receptor 40)[20] agonists as a antidiabetic activity.

Recently, GLP-1 has been identified as a potential target for treatment of T2DM. Gut hormone GLP-1 increases insulin secretion after eating. GLP-1 has been shown to improve glycemic control in patients with Type 2 diabetes.[21] As GLP-1 cleaves its N-terminal, DPP-IV controls its activity [7-36]-amide to form GLP-1[9-36]-amide, an inert compound. This method can be used to increase GLP-1 in the blood by inhibiting DPP-IV.[22] As a result, the investigation of DPP-IV inhibitors as possible treatments has been devoted a significant amount of time and effort as shown in Figure 1.

DOI: 10.5530/ijper.57.3s.85

Copyright Information :
Copyright Author (s) 2023 Distributed under Creative Commons CC-BY 4.0

Publishing Partner : EManuscript Tech. [www.emanuscript.in]

MATERIALS AND METHODS

ChemSketch software was used to simulate the 2D structures of the compounds, and AutoDock version 1.5.6 was used to dock the hits. A protein (PDB NO: 3W2T) was used as DPP-IV. Chemicals of analytical grade came from Nice Chemicals Pvt. Ltd., (India), Fizmerck (India), M Lychem chemical (India), and Poona chemical (India). We used the chemicals and solvents in their original forms, without purification.

Using Thin-Layer Chromatography (TLC), a reaction was monitored to determine if the reactants had been consumed and if a new product had been formed. Using the Labronics LT-115 digital melting point apparatus, we measured the melting

Back to Academic Year 2022-23

Novel Drug Delivery System: Review Article



WORLD JOURNAL OF PHARMACEUTICAL RESEARCH
SJIF Impact Factor 8.084
Volume 12, Issue 9, XXX-XXX. Review Article ISSN 2277– 7105

NOVEL DRUG DELIVERY SYSTEM: A REVIEW ARTICLE

Samta Khillare[1]*, Rushikesh Ghule[2], S. A. Waghmare[3] and Hemant Kamble[4]

[1]Department of Pharmaceutics, Loknete Shri Dadapatil Pharate Collage of Pharmacy Mandavgan Pharate Tal – Shirur District – Pune, Maharashtra, India.

[2]Department of Chemistry, Loknete Shri Dadapatil Pharate Collage of Pharmacy Mandavgan Pharate Tal – Shirur District – Pune, Maharashtra, India.

[3]CEO, Loknete Shri Dadapatil Pharate College of Pharmacy, Mandavgan Pharate Tal – Shirur District – Pune, Maharashtra, India.

[4]Principle, Loknete Shri Dadapatil Pharate College of Pharmacy, Mandavgan Pharate, Tal - Shirur, District- Pune, Maharashtra, India.

Article Received on
14 April 2023,
Revised on 04 May 2023,
Accepted on 24 May 2023
DOI: 10.20959/wjpr20239-27036

***Corresponding Author**
Samta Khillare
Department of Pharmaceutics, Loknete Shri Dadapatil Pharate Collage of Pharmacy Mandavgan Pharate Tal – Shirur District – Pune, Maharashtra, India.
samtakhillare1234@gmail.com

ABSTRACT

In recent years, a more logical approach to the creation of an ideal drug delivery system has been made possible by our growing understanding of the pharmacokinetic and pharmacodynamic behaviour of the medication. It is already apparent that multidisciplinary efforts will contribute significantly to future success in drug delivery research. The use of enhanced drug delivery systems, which are both safer and more effective, represents a step forward in the treatment of many human diseases as well as attractive marketing prospects for pharmaceutical companies. When a drug delivery system is designed properly, the target is met or the desired rate of drug administration is reached or demanded by the physiological needs of the body. Traditional pharmaceutical dosage forms are unable to regulate how quickly drugs are delivered to the target place. The distribution of the drug in non-target tissue and body fluids necessitates therapeutic doses that may be greatly greater than those needed in target cells, and the higher doses frequently result in severe adverse reactions during treatment. Therefore, novel drug delivery systems (NDDS) are those carriers that keep the drug concentration in the therapeutic range for a longer period of time and may also deliver the necessary dosage to the specific site of action as well.

A Review on Nanotechnology in Cancer Treatment

Back to Academic Year 2022-23

RT - Volume 7 Issue 9 – SEPTEMBER 2021

A Review on Nanotechnology in Cancer Treatment

Mr.Garje Mayur[1], Miss.Prerana Aurange[2], Prof.Santosh Waghmare[3], Dr. H. V Kamble[4]
[1, 2, 3, 4, 5]Loknete Shri Dadapatil Pharate College Of Pharmacy,Mandavgan Pharata

tract- *Cancer is a disease in which some of the body's s grow uncontrollably and spread to other parts of the ly. Cancer treatment is one of the major challenges of dern medicine. Cancer targeting treatment has been atly improved by new tools an approaches based on notechnology. Nanotechnology can be applied to target ugs to the surface or to the interior of specific cells. it can so be used in diagnosis and prognosis of diseases. In ldition , Nano shells and nanoparticles are the most romising applications for various cancer treatments.*

Keywords- cancer, nanotechnology, nanoparticles , cancer reatment

I. INTRODUCTION

Cancer is a disease in which some of the body's cells grow uncontrollably and spread to other parts of the body. Cancer is a group of diseases involving abnormal cell growth with the potential to invade or spread to other parts of the body. There are many types of cancer treatment. The types of treatment that you receive will depend on the type of cancer. Cancer therapy currently limited to surgery, radiation and chemotherapyall three methods risk damage to normal cells tissue selectively to cancerous cells and neoplasms and enhance the therapeutic efficacy of radiation based and other current treatment modalities.

Nanotechnology is science, engineering, and technology conducted at the nanoscale, which is about 1 to 100 manometers. Nano refers to the 10-9 power, or one billionth. For comparison, a human hair is about 100,000 nanometres thick. There are many interesting Nano devices being developed that have a potential to improve cancer detection, diagnosis, and treatment of cancer nanoparticles is defined as a partical of matter is between 1-100 nm

Nanoparticles can be defined as particles less than 100 nm in diameter that exhibit new or enhanced size - dependent properties compared with larger particles of the same material

Metal nanoparticles (silver and gold) are widely used in

1) cell imaging
2) DNA hybridization
3) proteins interaction and
4) photo thermal therapy

Advantages of nanoparticles :

1) Improved bioavailability by enhancing aqueous solubility
2) increasing resistance time in the body (increasing half life for clearance)
3) Stable dosage form of drugs
4) Faster dissolution
5) Decrease toxicity
6) Smaller dosage form

Nanotechnology in drug delivery system and cancer treatment

1. Dentrimers :

Dentrimersare highly branched, star-shaped macromolecules with manometer-scale dimensions. Dentrimers are defined by three components: a central core and an exterior surface with functional surface group The varied combination of these components yields products of different shapes and sizes with shielded interior cores .dentrimers have been explored for the encapsulation of hydrophobic compounds and for the delivery of anticancer drugs. There are three methods for using dendrimers in drug delivery. First, the drug is covalently attached to the periphery of the dendrimer to form dendrimer prodrugs. Second the drug is coordinated to the outer functional groups via ionic interactions .Third the dendrimer acts as a unimolecular micelle by encapsulating a pharmaceutical through the formation of a dendrimer-drug supramolecular assembly.



Back to Academic Year 2022-23

Number of books and chapters in edited volumes/books published and papers published in national/ international conference proceedings per teacher during last five years.

Summary

2018-2019	2019-2020	2020-2021	2021-2022	2022-2023
44	25	25	18	22

PRINCIPAL
SWGP'S LOKNETE SHRI DADAPATIL PHARATE
COLLEGE OF PHARMACY
Mandavgan Pharata, Tal. Shirur, Dist. Pune 412211

Number of books and chapters in edited volumes/books published and papers published in national/ international conference proceedings per teacher during last five years.

a. Book Publications

b. Papers Publications

Index

Book Publications

Sr.No.	Content	Document
1	Academic Year 2018-19	View Document
2	Academic Year 2019-20	View Document
3	Academic Year 2020-21	View Document
4	Academic Year 2021-22	View Document
5	Academic Year 2022-23	View Document

SWGP'S LOKNETE SHRI DADAPATIL PHARATE COLLEGE OF PHARMACY MANDAVGAN PHARATA, PUNE 412211

PRINCIPAL
SWGP'S LOKNETE SHRI DADAPATIL PHARATE
COLLEGE OF PHARMACY
Mandavgan Pharata, Tal. Shirur, Dist. Pune 412211

Number of books and chapters in edited volumes/books published and papers published in national/ international conference proceedings per teacher during last five years

Book Publications Academic Year 2018-2019

Sr. No	Name of The Teacher	Title of The Book/Chapter Published	Year of Publication	ISBN/ISSN Number of The Proceedings	Name of The Publisher	Content	Link
1	Dr. Vivek M. Saptupe	Practical Book of Pharmaceutical Inorganic Chemistry I	2018	ISBN: 9788193927632	Global Education	View Document	https://www.flipkart.com/pharmaceutical-inorganic-chemistry/p/itma2656044483804
2	Mr. Avinash V. Dhobale	Community Pharmacy and Management	2018	ISBN: 9789393834744	Pritam Publication	View Document	https://www.flipkart.com/community-pharmacy-management/p/itm4e2f16fde8ab1

SHRI LOKNETE SHRI DADAPATIL PHARATE COLLEGE OF PHARMACY
MANDAVGAN PHARATA, PUNE 412211

PRINCIPAL
SWGP'S LOKNETE SHRI DADAPATIL PHARATE
COLLEGE OF PHARMACY
Mandavgan Pharata, Tal. Shirur, Dist. Pune 412211

Back to Book Publications Index

Dr. V. M. Satpute



PRINCIPAL
SWGP'S LOKNETE SHRI DADAPATIL PHARATE
COLLEGE OF PHARMACY
Mandavgan Pharata, Tal. Shirur, Dist. Pune 412211

Mr. A. V. Dhobale

Back to Book Publications Index



About the Author

Mrs. Sneha Akash Mane
B. Pharm, M. Pharm

Mrs.Sneha Akash Mane Presently workingin Vidarbha Institute Of Pharmacy, Washim. Working as Assistant Professor in the Deparment of Pharmacy.Shehas Completed her Diploma and Graduation from Chatrapati Shahu Maharaj Shikshan Sanstha Maregaon.Tq Wani District Yavatmal and from Balpande College Of Pharmacy Mane wada Besa ,Nagpur.She is having 8 year teaching experience.She worked as Quality Control Incharge in Oswal Pharmaceutical for 1.5years.She attended one day seminar of Modern Teaching Technique and also she is lifetime member of APTI.

Dr. K. V. Otari

Dr. K. V. Otari is currently working as Professor and Principal at Navsahyadri Institute of Pharmacy,Pune.He completad M.Pharm in Pharmacology from Rajiv Gandhi University of Health Sciences, Bangalore, India and Ph.D.in Pharmacology from Savitribai PhulePune University,Pune,India.

He has over 16 year experience in the field of pharmacy education at graduate level & post-graduate level,10 years of research experience and 6 year so f industrial experience as a Director.He has a bout 41 peer-reviewed national and international publications, 03books/Chapter & one patent to his credit.He is approved PhD Guide by DBATU and Career Point University ,Kota and guided over 35 scholars for their PG and PhD research.He is recipient of grants for research and international travel.He is serving as are viewer for Scientific Journal sanda Member of Editorial Board of various Scientific Journals.

Mr. Avinash V. Dhobale Patil

Mr. Avinash V. Dhobale Patil is presently working as Assistant Professor Pharmaceutics Department at LSDP College of Pharmacy Mandavgan Pharata,Shirur Pune.(Savitribai Phule University Pune).He has completed his Graduation and Post Graduation in Pharmaceutics from Sudhakarrao Naik Institute of Pharmacy,Pusad,and Yavatmal. From SGBAU.He is pursing PhD in Pharmaceutical Science.(Pharmaceutics) He has published several research and review paper sin National and International journal so freputed.He has EightBook so in his credit. He has Lifetime member of APP (Membership ,MH State Branch) He has organized Two Conferences one is National Conference in year 2017-18, and also International conference organized at L.S.D.P College Pharmacy, Pune in 2019 and he achieved Best Researcher Award in Academic year 2018-19 by Association of Pharmacy Professionals APP,Maharashtra State Branch. He has more than 05Years Teaching Experience.

Mr.Hanumant Sopan Patil

Mr.Hanumant Sopan Patil Presently workingin Pune District Education Association's, Seth Govind Raghunath Sable College of Pharmacy,Saswad,Dit.PuneasaHOD/Lecturer In the Department of Pharmacy.He has Completed his M.Pharm.in Pharmaceutics,fromK.T.Patil College of Pharmacy,Osmanabad,Dr.BabasahebAmbedkar Marathwada University, Aurangabad. He is having 26 year teaching Experience.He has awarded as a best teacher by late Eknath Kaka Jagtap pratisthan,Saswad,in2020.He has Published Practical manual for Second Year Diploma in Pharmacy for Brilliant Publication,Pune.He has Published 3International papers.

Pritam Publication
5, Vardhaman Chamber, University Road, Jalgaon. Mob.: 98502 22351.
email - pritamjalgaon@rediffmail.com, website-www.pritampublication.com

COMMUNITY PHARMACY AND MANAGEMENT

COMMUNITY PHARMACY AND MANAGEMENT

- Mrs. Sneha Akash Mane
- Mr. Hanumant Sopan Patil
- Dr. K. V Otari
- Mr. Avinash V. Dhobale Patil

PRINCIPAL
SWGP'S LOKNETE SHRI DADAPATIL PHARATE
COLLEGE OF PHARMACY
Mandavgan Pharata, Tal. Shirur, Dist. Pune 412211

Back to Book Publications Index

Book Publications Academic Year 2019-2020

Sr. No	Name Of The Teacher	Title Of The Book/ Chapter Published	Year Of Publication	ISBN/ ISSN Number Of The Proceedings	Name Of The Publisher	Contents	Link
1	Mr. P. B. Cholake	A Practical Book Of Medicinal Chemistry-I	2019	ISBN: 978-93-24457-15-7 PT-1948	Success Publication	View Document	https://www.researchgate.net/publication/330038031_A_PRACTICAL_BOOK_OF_MEDICINAL_CHEMISTRY-I_FOR_THIRD_YEAR_BPHARMACY
2	Mr. R. M. Wani	Physical Pharmaceutics II	2020	-	PV Publication	View Document	https://pvbooks.in/?s=Physical+Pharmaceutics+II&product_cat=0&book_author=0&post_type=product
3	Mr. A. V. Dhobale						
4	Mr. S. N. Kharde						
5	Mr. R. M. Wani	Pharmacognosy & Phytochemistry I	2020	ISBN : 973-98-89739-17-6	Success Publication	View Document	https://www.researchgate.net/publication/338832332_Practical_Handbook_of_Pharmacognosy_and_Phytochemistry-I
6	Mr. A. V. Dhobale						
7	Mr. S. N. Kharde						
8	Mr. R. M. Wani	Practical Book of Pharmaceutical Organic Chemistry II	2020	ISBN:97815-43343-94	Medical Publishers	View Document	https://www.researchgate.net/publication/340248637_Practical_Book_of_Pharmaceutical_Organic_Chemistry_II
9	Mr. A. V. Dhobale						

PRINCIPAL
SWGP'S LOKNETE SHRI DADAPATIL PHARATE
COLLEGE OF PHARMACY
Mandavgan Pharata, Tal. Shirur, Dist. Pune 412211

Mr. P. B. Cholake

Back to Book Publications Index

A
PRACTICAL BOOK
OF

MEDICINAL CHEMISTRY- I

FOR
THIRD YEAR B.PHARMACY

(Semester V)

Prof (Dr.) Mrunal K.Shirsat
Principal, Shri Wagheshwar Gramvikas Pratishthan Sanchalit,
Loknete Shri Dadapatil Pharate College of Pharmacy,
Madavgoan Pharata, Shirur, Pune.

Prof (Dr.) Indrajeet Singhvi
Dean ,Faculty of Pharmacy, Pacific University, Udaipur
Pacific Hills, Pratap Nagar Udaipur-313003

Prof. P. B. Cholke
Assistant Professor& HOD of Pharmaceutics Department,
Shri Wagheshwar Gramvikas Pratishthan Sanchalit,
Loknete Shri Dadapatil Pharate College of Pharmacy,
Madavgoan Pharata, Shirur, Pune.

Prof. Y. S. Temak
Assistant Professor& HOD of Pharmaceutical Chemistry Department,
Shri Wagheshwar Gramvikas Pratishthan Sanchalit,
Loknete Shri Dadapatil Pharate College of Pharmacy,
Madavgoan Pharata, Shirur, Pune.

SUCCESS PUBLICATIONS

SWGP'S LOKNETE SHRI DADAPATIL PHARATE COLLEGE OF PHARMACY
MANDAVGAN PHARATA, PUNE 412211

PRINCIPAL
SWGP'S LOKNETE SHRI DADAPATIL PHARATE
COLLEGE OF PHARMACY
Mandavgan Pharata, Tal. Shirur, Dist. Pune 412211

Back to Book Publications Index

Mr. R. M. Wani, Mr. A. V. Dhobale & Mr. S. N. Kharde



PRINCIPAL
SWGP'S LOKNETE SHRI DADAPATIL PHARATE COLLEGE OF PHARMACY
Mandavgan Pharata, Tal. Shirur, Dist. Pune 412211

Back to Book Publications Index

Mr. R. M. Wani, Mr. A. V. Dhobale & Mr. S. N.Kharde





Back to Book Publications Index

Mr. R. M. Wani & Mr. A. V. Dhobale





SWGP'S LOKNETE SHRI DADAPATIL PHARATE COLLEGE OF PHARMACY MANDAVGAN PHARATA, PUNE 412211

PRINCIPAL
SWGP'S LOKNETE SHRI DADAPATIL PHARATE
COLLEGE OF PHARMACY
Mandavgan Pharata, Tal. Shirur, Dist. Pune 412211

Back to Book Publications Index

Book Publications Academic Year 2020-2021

Sr. No	Name of The Teacher	Title of The Book / Chapter Published	Year of Publication	ISBN/ISSN Number of The Proceedings	Name of The Publisher	Contents	Link
1	Mr. Avinash V. Dhobale	Pharmacognosy	2021		Brilliant Publication	View Documents	https://pvbooks.in/product/concise-course-in-pharmacognosy-and-phtochem-ii-sem-v/
2	Mr. P. B. Cholke	Textbook of Medicinal Chemistry - I	2021	ISBN : 9789393834133	Pritam Publication	View Documents	https://www.amazon.in/Textbook-Medicinal-Chemistry-I/dp/939383413X
3	Ms. Y. S. Temak						
4	Ms. Monali R. Bhalerao	Biochemistry	2021	ISBN : 978-93-5735-000-6	Pritam Publication	View Documents	https://www.amazon.in/Textbook-Biochemistry-Clinical-Pathology-Bhalerao/dp/9393834768

PRINCIPAL
SWGP'S LOKNETE SHRI DADAPATIL PHARATE
COLLEGE OF PHARMACY
Mandavgan Pharata, Tal. Shirur, Dist. Pune 412211

Back to Book Publications Index

Mr. Avinash V. Dhobale



PRINCIPAL
SWOP'S LOKNETE SHRI DADAPATIL PHARATE
COLLEGE OF PHARMACY
Mandavgan Pharata, Tal. Shirur, Dist. Pune 412211

Back to Book Publications Index

Dr. Y. S. Temak & Dr. P. B. Cholake



A TEXTBOOK OF
MEDICINAL CHEMISTRY-I

FOR
SECOND YEAR B. PHARMACY
(Semester IV)
(As per syllabus of PCI, New Delhi SPPU, Pune & DBATU, Lonere)

Dr. Sandip S. Kshirsagar
Dr. Rupali B. Dhamdhere
Dr. Yogita S. Temak
Dr. Pravin B. Cholke



PRINCIPAL
SWGP'S LOKNETE SHRI DADAPATIL PHARATE
COLLEGE OF PHARMACY
Mandavgan Pharata, Tal. Shirur, Dist. Pune 412211

Back to Book Publications Index

Ms. M. R. Bhalerao



PRINCIPAL
SWGP'S LOKNETE SHRI DADAPATIL PHARATE
COLLEGE OF PHARMACY
Mandavgan Pharata, Tal. Shirur, Dist. Pune 412211

Back to Book Publications Index

Book Publications Academic Year 2021-2022

Sr. No	Name of The Teacher	Title of The Book / Chapter Published	Year of Publication	ISBN/ISSN Number of The Proceedings	Name of The Publisher	Contents	Link
1	Ms. Monali R. Bhalerao	Industrial Pharmacy I	2022	ISBN : 978-93-5735-000-6	Pritam Publication	View Document	https://www.pritampublications.com/collection/Pharmacy/B-Pharm-3rd-Semester-Textbooks
2	Mr. Santosh A. Waghmare	Pharmaceutical Organic Chemistry - III	2022	ISBN-10 : 9392159099 ISBN-13: 9789392159091	Pritam Publication	View Document	https://www.amazon.in/Textbook-Pharmaceutical-Organic-Chemistry-III/dp/9392159099
3	Ms. A. A. Najan	Textbook of Novel Drug Delivery System	2022	ISBN : 978-93-5735-063-1	Pritam Publication	View Document	https://www.flipkart.com/textbook-novel-drug-delivery-system/p/itm1da8b02c9e731?pid=9789357350631&lid=LSTBOK9789357350631 75VO6D&marketplace=FLIPKART

4	Dr. M. R. Agrawal	Pharmacognosy And Phytochemistry I	2022	ISBN-10 : 9391567762 ISBN-13 : 978-9391567767	Technical Publications	View Document	https://www.amazon.in/Pharmacognosy-Phytochemistry-B-PHARMACY-SYLLABUS/dp/9391567762



ack to Book Publications Index

Ms. Monali R. Bhalerao

About the Author

Mr. Nilesh Mahadeo Ghuge
M. Pharm (Pharmaceutics)

Mr. Nilesh Mahadeo Ghuge Presently working in Dr. S. K. Toshniwal Educational & Research Trust's VIDARBHA INSTITUTE OF PHARMACY Gut. No 114. Anjankhed Borala Phata Hingoli Road Washim-444505 as Assistant Professor in the Department of Pharmaceutics. He has completed his Graduation (B.Pharm) from S.G.S.P.S Institute of Pharmacy Kaulkhed Akola and Post-Graduation (M.Pharm) from P. Wadhwani College of Pharmacy, Yavatmal Moha Phata Dhamangaon Road Yavatmal,(MH). He is having 11 years of Teaching Experience with Published many Research & Review Articles in Pharmaceutical Field.

Dr. Nandu Kayande
M.Pharm (Pharmacology)

Dr. Nandu Kayande, M.Pharm (Pharmacology) Ph.D. Principal Thakur Shivkumar Singh Memorial Pharmacy College, Burhanpur, MP. He has completed his Graduation from Shri Gurudatta Shikhan Prasarak Sanstha's Institute of Pharmacy, Kaulkhed, Akola From SGBAU Amaravati & Postgraduation From SRM College of Pharmacy, SRM University, Chennai. He has completed his Ph.D. in Pharmaceutical science from OPJS University, Churu Rajasthan. He has more than 14 years of experience in Teaching. He has one book on his credit. He has life time membership of APTI. He has published 35 Research & Review articles in reputed national and international journals. He as attended & presented 10 research paper in national & International conferences. He has guided 36 UG students for their minor projects and co-guided 4 students for Ph.D. dissertation.

Ms. M. R. Bhalerao

Ms. M. R. Bhalerao is currently working as Professor at L. S. D. P. College of Pharmacy, Mandavgan Pharata, Pune. She completed M. Pharm in Pharmaceutical Chemistry from Sinhgad Institute of Pharmacy, Lonavala, Pune.
She is having 7 years experience in the field of pharmacy education at graduate level and 1 year of industrial experience. Her 5 papers are published in India. She is **Life time member of APTI,** New Delhi.
She is received **Bharat Gaurav Award** for year 2022 from New Delhi.

Ms. Priyanka D. Borude
M.Pharm (Pharmaceutics), Ph.D appeared.

Ms. Priyanka Borude is presently working as lecturer at Pune District Education Association's, Shankarrao Ursal College of Pharmacy (Diploma), Kharadi; Dist.Pune.She has teaching and Industrial experience of 12 years. She has attended many National, International conferences and seminars. She has published several national and international research and review papers in reputed international journals. She has completed post graduation from MCE Society's Allana college of pharmacy,Camp,Pune and UG from AISSPMS College of Pharmacy, Pune.and persuing Ph.D from JJTU ,Rajasthan. She has also delivered a guest lectures on Personality development in various reputed Institutes.

Mr. Sunil Mahadeo Bhoyar
M Pharm

Mr. Sunil Mahadeo Bhoyar Presently working in MahilaUtkarshaPratisthan's Institute of Pharmacy, Degaon Risod Dist-Washim as Principal in the Department of Pharmacy. He has completed his Graduation (B.Pharm) and Post-Graduation (M.Pharm) from Sudhakarrao Naik Institute of Pharmacy Pusad Tq Pusad District Yavatmal, He is having 09 years of Teaching Experience with Published many Research & Review Articles in Pharmaceutical Field.

Pritam Publications
5, Vardhaman Chamber, University Road, Jalgaon. Mob.: 98502 22351.
email - pritamjalgaon@rediffmail.com, website-www.pritampublication.com

Textbook of Industrial Pharmacy



SWGP'S LOKNETE SHRI DADAPATIL PHARATE COLLEGE OF PHARMACY
MANDAVGAN PHARATA, PUNE 412211

PRINCIPAL
SWGP'S LOKNETE SHRI DADAPATIL PHARATE
COLLEGE OF PHARMACY
Mandavgan Pharata, Tal. Shirur, Dist. Pune 412211

Back to Book Publications Index

Prof. S. A. Waghmare

A Textbook of

Pharmaceutical Organic Chemistry - III

As per PCI Syllabus
for B. Pharmacy Semester - IV



Ms. Priyanka R. Varade
Mrs. Jyoti N. Kadam
Mrs. Bhagyashri J. Warude
Prof. Santosh A. Waghmare

SWGP'S LOKNETE SHRI DADAPATIL PHARATE COLLEGE OF PHARMACY
MANDAVGAN PHARATA, PUNE 412211

PRINCIPAL
SWGP'S LOKNETE SHRI DADAPATIL PHARATE
COLLEGE OF PHARMACY
Mandavgan Pharata, Tal. Shirur, Dist. Pune 412211

Back to Book Publications Index

Ms. A. A. Najan

A Textbook of

Novel Drug Delivery System

(Semester VII)

Dr. Rahul S. Solunke
Mr. Vinod D. Usnale
Ms. Anita A. Najan
Ms. Pooja P. Dahale
Mr. Vijaykumar R. Chakote

PRINCIPAL
SWGP'S LOKNETE SHRI DADAPATIL PHARATE
COLLEGE OF PHARMACY
Mandavgan Pharata, Tal. Shirur, Dist. Pune 412211

Back to Book Publications Index

Dr. M. R. Agrawal



PRINCIPAL
SWGP'S LOKNETE SHRI DADAPATIL PHARATE
COLLEGE OF PHARMACY
Mandavgan Pharata, Tal. Shirur, Dist. Pune 412211

Back to Book Publications Index

Book Publications Academic Year 2022-2023

Sr. No	Name of The Teacher	Title of The Book/Chapter Published	Year of Publication	ISBN / ISSN Number of The Proceedings	Name of The Publisher	Contents	Link
1	Ms. Monali R. Bhalerao	Pharmaceutical Management	2023	ISBN : 978-93-5735-000-6	Pritam Publication	View Document	https://www.amazon.in/Textbook-Marketing-Management-Bhalerao-Priyanka/dp/9357350004
2	Mr. A. V. Dhobale	Human Anatomy & Physiology	2023	-	Brilliant Publication	View Document	https://www.flipkart.com/human-anatomy-physiology/p/itm6cc825e220eea

SWGP'S LOKNETE SHRI DADAPATIL PHARATE COLLEGE OF PHARMACY MANDAVGAN PHARATA, PUNE 412211

PRINCIPAL
SWGP'S LOKNETE SHRI DADAPATIL PHARATE
COLLEGE OF PHARMACY
Mandavgan Pharata, Tal. Shirur, Dist. Pune 412211

Back to Book Publications Index

Ms. Bhalerao Monali R.



SWGP'S LOKNETE SHRI DADAPATIL PHARATE COLLEGE OF PHARMACY
MANDAVGAN PHARATA, PUNE 412211

PRINCIPAL
SWGP'S LOKNETE SHRI DADAPATIL PHARATE COLLEGE OF PHARMACY
Mandavgan Pharata, Tal. Shirur, Dist. Pune 412211

Back to Book Publications Index

Dr. A. V. Dhobale



PRINCIPAL
SWGP'S LOKNETE SHRI DADAPATIL PHARATE
COLLEGE OF PHARMACY
Mandavgan Pharata, Tal. Shirur, Dist. Pune 412211

Back to Book Publications Index

Papers Publications

Index

Sr. No.	Content	Document
1	Academic Year 2018-2019	View Document
2	Academic Year 2019-2020	View Document
3	Academic Year 2020-2021	View Document
4	Academic Year 2021-2022	View Document
5	Academic Year 2022-2023	View Document

PRINCIPAL
SWGP'S LOKNETE SHRI DADAPATIL PHARATE
COLLEGE OF PHARMACY
Mandavgan Pharata, Tal. Shirur, Dist. Pune 412211

Academic Year 2018-2019

Sr. no	Title of the paper	Name of Staff	Department	Name of the journal	ISSN number	Link
1	Evaluation Of Some Natural Polymers As Pharmaceutical Excipient For Sustained Drug Delivery System	Mr. A. V. Dhobale	Pharmaceutics	International Journal Of Humanities And Social Science	E-ISSN 2348 – 1269	https://www.researchgate.net/publication/350782955_Evaluation_of_Some_Natural_Polymers_as_Pharmaceutical_Excipient_For_Sustained_Drug_Delivery_System
2	Recent Advances In Mucoadhesive Buccal Drug Delivery System And Its Marketed Scope And Opportunities	Mr. A. V. Dhobale	Pharmaceutics	International Journal Of Advanced Pharmaceutical Sciences	ISSN: 2456-8147	https://www.researchgate.net/publication/327111123_RECENT_ADVANCES_IN_MUCOADHESIVE_BUCCAL_DRUG_DELIVERY_SYSTEM_AND_ITS_MARKETED_SCOPE_AND_OPPORTUNITIES
3	Evaluation Of Antiulcer Activity Of Aconitum Heterophyllum On Experimental Animal	Mr. A. V. Dhobale	Pharmaceutics	World J Pharm Pharm Sci	ISSN 2278 – 4357	https://www.researchgate.net/publication/326837658_EVALUATION_OF_ANTIULCER_ACTIVITY_OF_ACONITUM_HETEROPHYLLUM_ON_EXPERIMENTAL_ANIMAL
4	Design And In-Vitro Characterization Of Gastroretentive Floating Tablets Containing Gliclazide	Mr. A. V. Dhobale	Pharmaceutics	World J Pharm Pharm Sci	ISSN 2278 – 4357	https://www.researchgate.net/publication/322307709_DESIGN_AND_IN-VITRO_CHARACTERIZATION_OF_GASTRORETENTIVE_FLOATING_TABLETS_CONTAINING_GLICLAZIDE

5	Mouth Dissolving Strips	Mr. A. V. Dhobale	Pharmaceutics	International Journal Of Advanced Pharmaceutical Sciences	ISSN: 2456-8147	https://www.researchgate.net/profile/Avinash-Dhobale-2
6	Formulation And Evaluation Of Fast Disintegrating Tablet Of Carvedilol.	Mr. A. V. Dhobale	Pharmaceutics	International Journal Of Pharmacy Pharmaceutical Sciences	ISSN 2349 7203	https://www.researchgate.net/publication/327690399_Formulation_and_Evaluation_of_Fast_Disintegrating_Tablet_of_Carvedilol
7	Evaluation Of Antiulcer Activity Of Aconitum Heterophyllum On Experimental Animal	Mr. A. V. Dhobale	Pharmaceutics	World Journal Of Pharmacy And Pharmaceutical Sciences	ISSN 2278 4357	https://www.researchgate.net/publication/326837658_EVALUATION_OF_ANTIULCER_ACTIVITY_OF_ACONITUM_HETEROPHYLLUM_ON_EXPERIMENTAL_ANIMAL
8	Formulation And Evaluation Of Atorvastatin Calcium Oral Dispersible Table	Mr. A. V. Dhobale	Pharmaceutics	International Journal Of Current Advanced Research	ISSN: O: 2319-6475, ISSN: P: 2319-6505,	https://www.researchgate.net/publication/326831851_FORMULATION_AND_EVALUATION_OF_ATORVASTATIN_CALCIUM_ORAL_DISPERSIBLE_TABLET
9	Formulation And Evaluation Of Carbamazepine Nanoemulsion For Brain Targeted Drug Delivery Via Intranasal Route	Mr. A. V. Dhobale	Pharmaceutics	Indo American Journal Of Pharmaceutical Research	ISSN NO: 2231-6876	https://www.researchgate.net/publication/326803830_FORMULATION_AND_EVALUATION_OF_CARBAMAZEPINE_NANOEMULSION_FOR_BRAIN_TARGETED_DRUG_DELIVERY_VIA_INTRANASAL_ROUTE
10	Recent Advances In Pilot Plant Scale Up Techniques	Mr. A. V. Dhobale	Pharmaceutics	Indo American Journal Of Pharmace	ISSN NO: 2231-6876	https://www.researchgate.net/publication/326304751_RECENT_ADVANCES_IN_PILOT_PLANT_SCALE_UP_TECHNIQUES-A_Review

				utical Research,		Back to Paper Publication
11	Design, Development And Characterization Oflosartan Potassium Sustained Release Matrix Tablet	Mr. A. V. Dhobale	Pharmaceutics	World Journal Of Pharmacy And Pharmaceutical Scinces	ISSN 2278 – 4357	https://www.researchgate.net/publication/322308014_DESIGN_DEVELOPMENT_AND_CHARACTERIZATION_OF_LOSARTAN_POTASSIUM_SUSTAINED_RELEASE_MATRIX_TABLET
12	Design And In-Vitro Characterization Of Gastroretentive Floating Tablets Containing Gliclazide	Mr. A. V. Dhobale	Pharmaceutics	World Journal Of Pharmacy And Pharmaceutical Sciences	ISSN 2278 – 4357	https://www.researchgate.net/publication/322307709_DESIGN_AND_IN-VITRO_CHARACTERIZATION_OF_GASTRORETENTIVE_FLOATING_TABLETS_CONTAINING_GLICLAZIDE
14	Formulation And Evaluation Of Rosuvastatin Oral Dispersible Tablet Under A Creative Commons	Dr. V. M. Satpute	Pharmaceutics	International Journal Of Advanced Pharmaceutical Sciences	ISSN: 2456-8147	https://www.researchgate.net/publication/340262020_FORMULATION_AND_EVALUATION_OF_ROSUVASTATIN_ORAL_DISPERSIBLE_TABLET_under_a_Creative_Commons_Attribution_40
15	A Project On Human Papillomavirus-A Review Under A Creative Commons Attribution	Dr. V. M. Satpute	Pharmaceutics Pharmaceutics	International Journal Of Advanced Pharmaceutical Sciences	SSN:2456-8147	https://www.researchgate.net/publication/326571009_A_PROJECT_ON_HUMAN_PAPILLOMAVIRUS_-_A_REVIEW
16	Formulation And Evaluation Of Fast Disintegrating Tabletof Carvedilol	Dr. V. M. Satpute	Pharmaceutics	Human Journals IJPPR	ISSN 2349 7203	https://www.researchgate.net/publication/327690399_Formulation_and_Evaluation_of_Fast_Disintegrating_Tablet_of_Carvedilol
17	Isolation And Characterization			Journal Of Pharmaco	E-ISSN:	https://rjpponline.org/AbstractView.aspx?PID=2020-12-3-6

	Of Phytoconstituents From Petroleum Ether Extract Of Momordica Cochinchinensis Fruits	Mr. M. R. Agrawal	Pharmacognosy	gnosy And Phytochemistry	2278-4136 P-ISSN: 2349-8234	
18	Anti-Inflammatory Activity Of Momordica Cochinchinensis And Momordica Balsamina Fruit Extracts	Mr. M. R. Agrawal	Pharmacognosy	International Journal Of Green	DOI: 10.24214/IJGHC/HC/7/4/44655.	https://www.academia.edu/42305275/Analgesic_activity_of_Momordica_cochinchinensis_and_Momordica_balsamina_fruit_extracts
19	Analgesic Activity Of Momordica Cochinchinensis And Momordica Balsamina Fruit Extracts	Mr. M. R. Agrawal	Pharmacognosy	International Journal Of Gree Pharmacy	Vol 12(4)	https://www.academia.edu/42305275/Analgesic_activity_of_Momordica_cochinchinensis_and_Momordica_balsamina_fruit_extracts
20	Ameliorative Effects of Artemisia Pallens In A Murine Model of Ovalbumin-Induced Allergic Asthma Via Modulation of Biochemical perturbations	Ms. A. A. Mukherjee	Pharmacognosy	Biomedicine & Pharmacotherapy	Https://Doi.Org/10.1016/J.Fct.2018.08.028	https://pubmed.ncbi.nlm.nih.gov/28810518/#:~:text=Conclusion%3A%20Artemisia%20pallens%20simultaneously%20orchestrate,OVA%2Dinduced%20AHR%20in%20rats.
21	Evaluation f Health-Related Quality Of Life In Hemolytic Uraemic Syndrome Patients Treated With Eculizumab: A	Ms. A. A. Mukherjee	Pharmacognosy	Taylor & Francis	ISSN: 0886-022X (Print) 1525-6049	https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6014301/

	Systematic Evaluation On Basis Of EMPRO					
22	Curcumin Ameliorates Vitamin A Deficiency-Induced Urolithiasis In Neonatal Rats Via Inhibition Of KIM-1/NGAL, Nrf2, And Inos Pathways	Ms. A. Kandhare	Pharmacology	Latin American Journal Of Pharmacy	ISSN 0326 2383 ISSN 2362-3853	https://www.researchgate.net/publication/331045453_Curcumin_ameliorates_vitamin_A_deficiency-induced_urolithiasis_in_neonatal_rats_via_inhibition_of_KIM-1NGAL_Nrf2_and_iNOs_pathways
23	Neuroprotective Effect Of Naringin, A Flavone Glycoside In Quinolinic Acidinduced Neurotoxicity: Possible Role Of PPAR-Γ, Bax/Bcl-2, And Caspase-3	Ms. A. Kandhare	Pharmacology	Food And Chemical Toxicology 121 (2018) 95–108	Https://Doi.Org/10.1016/J.Fct.2018.08.028	https://pubmed.ncbi.nlm.nih.gov/30130594/
24	Low Molecular Weight Galactomannans-Based Standardized Fenugreek Seed Extract Ameliorates High-Fat Diet-Induced Obesity In Mice Via Modulation Of Fasn, IL-6, Leptin, And TRIP-Br_2	Ms. A. Kandhare	Pharmacology	RSC Advances	Rsc.Li/Rsc-Advances	https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9086199/

25	Anti-Epileptic Effect Of Morin Against Experimental Pentylenetetrazol-Induced Seizures Via Modulating Brain Monoamines And Oxidative Stress	Ms. A. Kandhare	Pharmacology	Asian Pacific Journal Of Tropical Biomedicine 2018; 8(7): 352-359	ISSN 2221-1619	https://www.researchgate.net/publication/326472470_Anti-epileptic_effect_of_morin_against_experimental_pentylenetetrazol-induced_seizures_via_modulating_brain_monoamines_and_oxidative_stress
26	Add-On Therapy Of Herbal Formulation Rich In Standardized Fenugreek Seed Extract In Type 2 Diabetes Mellitus Patients With Insulin Therapy: An Efficacy And Safety Study	Ms. A. Kandhare	Pharmacology	Asian Pacific Journal Of Tropical Biomedicine 2018; 8(7): 352-359	ISSN 2221-1619	https://www.researchgate.net/publication/327910450_Add-on_therapy_of_herbal_formulation_rich_in_standardized_fenugreek_seed_extract_in_type_2_diabetes_mellitus_patients_with_insulin_therapy_An_efficacy_and_safety_study
27	Efficacy And Safety Of Herbal Formulation Rich In Standardized Fenugreek Seed Extract As Add-On Supplementation In Patients With Type 2 Diabetes Mellitus On Sulfonylurea Therapy: A 12-Week, Randomized, Double-Blind, Placebo-Controlled, Multi-Center Study	Ms. A. Kandhare	Pharmacology	Pharmacognosy Magazine	ISSN : 0973-1296	https://www.researchgate.net/publication/327521981_Efficacy_and_safety_of_herbal_formulation_rich_in_standardized_fenugreek_seed_extract_as_add-on_supplementation_in_patients_with_type_2_diabetes_mellitus_on_sulfonylurea_therapy_A_12-week_randomized_d

Back to Paper Publication

28	Evaluation Of Health-Related Quality Of Life In Hemolytic Uraemic Syndrome Patients Treated With Eculizumab: A Systematic Evaluation On Basis Of EMPRO	Ms. A. Kandhare	Pharmacology	Taylor & Francis	ISSN: 1525-6049	https://pubmed.ncbi.nlm.nih.gov/29363392/
29	Hesperidin, A Plant Flavonoid Accelerated The Cutaneous Wound Healing In Streptozotocin-Induced Diabetic Rats: Role Of Tgf-B/Smads And Ang-1/Tie2	Ms. A. Kandhare	Pharmacology	EXCLI Journal	ISSN 1611-2156	https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5962903/
30	Anti-Allergic Potential Of Fisetin In A Murine Model Of Ova-Induced Allergic Rhinitis Via Inhibition Of Gata-3 And Th2 Cytokines Zhao L	Ms. A. Kandhare	Pharmacology	Biomedica	Vol. 34, Issue 2, Apr. – Jun., 2018	https://www.google.com/search?q=Anti-Allergic+Potential+Of+Fisetin+In+A+Murine+Model+Of+Ova-Induced+Allergic+Rhinitis+Via+Inhibition+Of+Gata-3+And+Th2+Cytokines+Zhao+L+research+paper+by+amit+kandhare&rlz=1C1VDKB_en-GBIN1070IN1070&oq=Anti-Allergic+Potential+Of+Fisetin+In+A+Murine+Model+Of+Ova-Induced+Allergic+Rhinitis+Via+Inhibition+Of+Gata-3+And+Th2+Cytokines+Zhao+L+research+paper+by+amit+kandhare&gs_lcrp=EgZjaHJvbWUqBggAEEUYOzIGCAAQRRg7MgcIARAhGI8C0gEJMTAwOTNqMGo3qAIAsAIA&sourceid=chrome&ie=UTF-8#ip=1

31	Elucidation Of The Molecular Mechanism Of Tempol In Pentylenetetrazol-Induced Epilepsy In Mice: Role Of Gamma - Amino butyric Acid, Tumor Necrosis Factor-Alpha, Interleukin-1β And C-Fos	Ms. A. Kandhare	Pharmacology	Pharmacognosy Magazine	ISSN : 0973-1296	https://ouci.dntb.gov.ua/en/works/4k3GEqn4/
32	Fisetin, A Plant Flavonoid Ameliorates Doxorubicin-Induced Cardiotoxicity In Experimental Rats: The Decisive Role OfCaspase-3, COX-II, Ctn-I, Inos And TNF	Ms. A. Kandhare	Pharmacology	Molecular Biology Reports	ISSN 0301-4851	https://ouci.dntb.gov.ua/en/works/7P2pwv0l/
33	Pioglitazone And Risk Of Bladder Cancer In Type 2 Diabetes Mellitus Patients: A Systematic Literature Review And Meta-Analysis Of Observational Studies Using Real-World Data	Ms. A. Kandhare	Pharmacology	Clinical Epidemiology And Global Health 6 (2018) 61–68	Http://Dx.Doi.Org/10.1016/J.Cegh.2017.08.002	https://www.sciencedirect.com/science/article/pii/S2213398417300544
34	Development And Validation Of UV-Visible Spectroscopy	Mr. Pravin	Pharmaceutical Chemistry	International Journal Of Research	ISSN: 2455-698	https://www.pharmacyjournal.in/archives/2018/vol3/issue4/3-4-16

	Method For Simultaneous Estimation Of Saxagliptin Hydrochloride And Metformin Hydrochloride In Tablet Dosage Form	Cholke & Ms. Yogita Temak		In Pharmacy And Pharmaceutical Sciences		
35	Formulation And Evaluation Of Mouth Dissolving Tablet Of Bendroflumethiazide	Ms. S. A. Nagare	Pharmaceutical Chemistry	International Journal Of Current Research	ISSN: 0975-833	https://www.researchgate.net/publication/326849498_FORMULATION_AND_EVALUATION_OF_MOUTH_DISSOLVING_TABLET_OF_BENDROFLUMETHIAZIDE
36	"Formulation And Evaluation Of Carbamazepine Nanoemulsion For Brain Targeted Drug Delivery Via Intranasal Route"	Ms. S. A. Nagare	Pharmaceutical Chemistry	Indo American Journal Of Pharmaceutical Research	ISSN NO: 2231-6876	https://www.researchgate.net/publication/326803830_FORMULATION_AND_EVALUATION_OF_CARBAMAZEPINE_NANOEMULSION_FOR_BRAIN_TARGETED_DRUG_DELIVERY_VIA_INTRANASAL_ROUTE
37	Formulation And Evaluation Of Mouth Dissolving Tablet Of Bendroflumethiazide	Ms. A. A. Najan	Pharmaceutical Chemistry	International Journal Of Current Research	ISSN: 0975-833	https://scholar.google.co.in/scholar?q=Formulation+And+Evalu...Ta...e+research+paper+by+a.a.najan&hl=en&as_sdt=0&as_vis=1&oi=scholart
38	Curcumin Ameliorates Vitamin A Deficiency-Induced Urolithiasis In Neonatal Rats Via Inhibition Of	Ms. A. A. Mukherjee	Pharmacology	Latin American Journal Of Pharmacy	ISSN 0326 2383 ISSN 2362-3853	https://www.researchgate.net/publication/331045453_Curcumin_ameliorates_vitamin_A_deficiency-induced_urolithiasis_in_neonatal_rats_via_inhibition_of_KIM-1NGAL_Nrf2_and_iNOs_pathways

Back to Paper Publication

	KIM-1/NGAL, Nrf2, And Inos Pathways					
39	Evaluation Of Health-Related Quality Of Life In Hemolytic Uraemic Syndrome Patients Treated With Eculizumab: A Systematic Evaluation On Basis Of EMPRO	Ms. A. A. Mukherjee	Pharmacology	Taylor & Francis	ISSN: 1525-6049	https://pubmed.ncbi.nlm.nih.gov/29363392/
40	Hesperidin, A Plant Flavonoid Accelerated The Cutaneous Wound Healing In Streptozotocin-Induced Diabetic Rats: Role Of Tgf-B/Smads And Ang-1/Tie2	Ms. A. A. Mukherjee	Pharmacology	EXCLI Journal	ISSN 1611-2156	https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5962903/
41	Morin ameliorates ovalbumin-induced allergic rhinitis via inhibition of STAT6/SOCS1 and GATA3/T-bet signaling pathway in BALB/c mice	Mr. Amit Kandhare	Pharmacology	Journal of Functional Foods	55(3):391-401	https://www.researchgate.net/publication/332123811_Morin_ameliorates_ovalbumin-induced_allergic_rhinitis_via_inhibition_of_STAT6SOCS1_and_GATA3T-bet_signaling_pathway_in_BALB_c_mice
42	Morin ameliorates ovalbumin-induced allergic rhinitis via inhibition of STAT6/SOCS1 and GATA3/T-bet signaling	Ms. A. A. Mukherjee	Pharmacology	Journal of Functional Foods	55(3):391-401	https://www.researchgate.net/publication/332123811_Morin_ameliorates_ovalbumin-induced_allergic_rhinitis_via_inhibition_of_STAT6SOCS1_and_GATA3T-

	pathway in BALB/c mice					bet signaling pathway in BALB c mice

Back to Paper Publication

Academic Year 2019-2020

Sr. no	Title of the paper	Name of Staff	Department	Name of the journal	ISSN number	Link
1	Ferulic acid ameliorates doxorubicin-induced cardiac toxicity in rats	Dr. A. D. Kandhare	Pharmacology	Naunyn-Schmiedeberg's Arch Pharmacol	https://doi.org/10.1007/s00210-019-01623-4	https://www.researchgate.net/publication/330967015_Ferulic_acid_ameliorates_doxorubicin-induced_cardiac_toxicity_in_rats
2	Diarylheptanoid, a constituent			Journal of Ethnopha	https://doi.	https://www.researchgate.net/publication/328300349_Diarylh

	isolated from methanol extract of Alpinia officinarum attenuates TNF-α level in Freund's complete adjuvant-induced arthritis in rats	Dr. A. D. Kandhare	Pharmacology	rmacology	org/10.1016/j.jep.2018.10.019	eptanoid a constituent isolated from methanol extract of Alpinia officinarum attenuates TNF-a level in Freund's complete adjuvant-induced arthritis in rats
3	Efficacy of Antioxidant Supplements on Prevention and Amelioration of Cisplatin-Induced Nephrotoxicity: A Systematic Review and Meta-analysis of Randomized Controlled Trials	Dr. A. D. Kandhare	Pharmacology	Jundishapur J Nat Pharm	Jundishapur J Nat Pharm Prod	https://www.researchgate.net/publication/334747602_Efficacy_of_Antioxidant_Supplements_on_Prevention_and_Amelioration_of_Cisplatin-Induced_Nephrotoxicity_A_Systematic_Review_and_Meta-analysis_of_Randomized_Controlled_Trials
4	A systematic literature review of fenugreek seed toxicity by using ToxRTool: evidence from preclinical and clinical studies	Dr. A. D. Kandhare	Pharmacology	Elsevier Ltd	https://doi.org/10.1016/j.heliyon.2019.e01536	https://www.researchgate.net/publication/332619750_A_systematic_literature_review_of_fenugreek_seed_toxicity_by_using_ToxRTool_evidence_from_preclinical_and_clinical_studies
5	Therapeutic Potential of Morin in Ovalbumin-induced Allergic Asthma Via Modulation of SUMF2/IL-13 and BLT2/NF-kB	Dr. A. D. Kandhare	Pharmacology	Current Molecular Pharmacology	Current Molecular Pharmacology, 2019, Vol. 12,	https://www.researchgate.net/publication/330122445_Therapeutic_Potential_of_Morin_in_Ovalbumin-induced_Allergic_Asthma_Via_Modulation_of_SUMF2IL-13_and_BLT2NF-kB_Signaling_Pathway

	Signaling Pathway					
6	Hesperidin Ameliorates Bleomycin-Induced Experimental Pulmonary Fibrosis Via Inhibition Of Tgf-B1/Smad3/Ampk And Iκbα/Nf-Kb Pathways	Dr. A. D. Kandhare	Pharmacology	EXCLI Journal	ISSN 1611-2156	https://www.researchgate.net/publication/336589331_Hesperidin_ameliorates_bleomycin-induced_experimental_pulmonary_fibrosis_via_inhibition_of_TGF-beta1Smad3AMPK_and_IkappaBalphaNF-kappaB_pathways
7	Acute and sub-chronic oral toxicity studies of hesperidin isolated from orange peel extract in Sprague Dawley rats	Dr. A. D. Kandhare	Pharmacology	Regulatory Toxicology and Pharmacology	www.elsevier.com/locate/yrtph	https://www.researchgate.net/publication/332404120_Acute_and_sub-chronic_oral_toxicity_studies_of_hesperidin_isolated_from_orange_peel_extract_in_Sprague_Dawley_rats
8	Efficacy of Antioxidant Supplements on Prevention and Amelioration of Cisplatin-Induced Nephrotoxicity: A Systematic Review and Meta-analysis of Randomized Controlled Trials	Ms. A. A. Mukharjee	Pharmacology	Jundishapur J Nat Pharm	Jundishapur J Nat Pharm Prod	https://www.researchgate.net/publication/334747602_Efficacy_of_Antioxidant_Supplements_on_Prevention_and_Amelioration_of_Cisplatin-Induced_Nephrotoxicity_A_Systematic_Review_and_Meta-analysis_of_Randomized_Controlled_Trials
9	Therapeutic Potential of Morin in Ovalbumin-induced Allergic Asthma Via Modulation of SUMF2/IL-13			Current Molecular Pharmacology	Current Molecular Pharmacology, 2019,	https://www.researchgate.net/publication/330122445_Therapeutic_Potential_of_Morin_in_Ovalbumin-induced_Allergic_Asthma_Via_Modulation_of_SUMF2IL-

	and BLT2/NF-kB Signaling Pathway	Ms. A. A. Mukharjee	Pharmacology		Vol. 12,	13 and BLT2NF-kB Signaling Pathway
10	Acute and sub-chronic oral toxicity studies of hesperidin isolated from orange peel extract in Sprague Dawley rats	Ms. A. A. Mukharjee	Pharmacology	Regulatory Toxicology and Pharmacology	www.elsevier.com/locate/yrtph	https://www.researchgate.net/publication/332404120_Acute_and_sub-chronic_oral_toxicity_studies_of_hesperidin_isolated_from_orange_peel_extract_in_Sprague_Dawley_rats
11	Development and Validation of RP-HPLC and UV Spectrophotometric Absorptivity Method for Simultaneous Estimation of Cyclobenzaprine hydrochloride and Aceclofenac in Pharmaceutical dosage form	Mr. A. S. Gaikwad	Pharmaceutical Chemistry	International Journal of Scientific Development and Research	ISSN: 2455-2631	https://scholar.google.com/citations?view_op=view_citation&hl=en&user=ORlyYt4AAAAJ&citation_for_view=ORlyYt4AAAAJ:kNdYIx-mwKoC
12	A Review - Corona Virus (Covid-19)	Mr. A. S. Gaikwad	Pharmaceutical Chemistry	International Journal for Research Trends and Innovation	ISSN: 2456-3315	https://scholar.google.com/citations?view_op=view_citation&hl=en&user=ORlyYt4AAAAJ&citation_for_view=ORlyYt4AAAAJ:YOwf2qJgpHMC
13	Targeted Drug Delivery – From Magic Bullet to Nanomedicine	Mr. A. S. Gaikwad	Pharmaceutical Chemistry	Journal of Xidian University	ISSN No:1001-240	https://scholar.google.com/citations?view_op=view_citation&hl=en&user=ORlyYt4AAAAJ&citation_for_view=ORlyYt4AAAAJ:kNdYIx-mwKoC

14	Cancer Chemotherapy with Peptides & Peptidomimetics drug & Peptide based Vaccine	Ms. S. A. Nagare	Pharmaceutical Chemistry	International Research Journal of Pharmacy and Medical Sciences	ISSN2581-3277	https://www.researchgate.net/publication/330957007_Cancer_Chemotherapy_with_Peptides_and_Peptidomimetics_Drug_and_Peptide_Based-Vaccines
15	A study to assess the knowledge regarding hand hygiene practice among secondary school going children in selected schools of Pune city	Ms. S. B. Kshirsagar	Pharmaceutical Chemistry	International Journal of Applied Research 2020; 6(12): 27-29	ISSN : 2394-7500	https://www.researchgate.net/publication/362517268_A_study_to_assess_the_knowledge_regarding_hand_hygiene_practice_among_secondary_school_going_children_in_selected_schools_of_Pune_city

Back to Paper Publication

Academic Year 2020 - 2021

Sr. no	**Title of the paper**	**Name of Staff**	**Department**	**Name of the journal**	**ISSN number**	**Link**
1	Genotoxicity: Malignancy, Damage To DNA	**Dr. H. V. Kamble**	Pharmacology	World Journal of Pharmaceutical Research	ISSN 2277 – 7105	https://wjpr.net/abstract_show/18131
2	Herbal Medicines and Screening Models for Hepatoprotective Agents	Dr. H. V. Kamble	Pharmacology	World Journal of Pharmaceutical Research	ISSN 2277 – 7105	https://www.google.com/search?q=Herbal+Medicines+and+Screening+Models+for+Hepatoprotective+Agents+research+paper+by+pratiksha+kothari&rlz=1C1VDKB_en-GBIN1070IN1070&oq=Herbal+Medicines+and+Screening+Models+for+Hepatoprotective+Agen

						ts+research+paper+by+pratiksha+kothari&gs_lcrp=EgZjaHJvbWUyBggAEEUYOTIGCAEQRRhAMgYIAhBFGDvSAQkyMDkxNWowajeoAgCwAgA&sourceid=chrome&ie=UTF-8
3	Drug- Excipient Interactions Study	Dr. H. V. Kamble	Pharmacology	World Journal of Pharmacy and Pharmaceutical Sciences	ISSN 2277 – 7105	https://www.ijnrd.org/viewpaperforall?paper=IJNRD2302162
4	Quality By Design Approaches To Analytical Method Development: A Review	Mr. S. A. Waghmare	Pharmaceutical Chemistry	World Journal of Pharmaceutical Research World Journal of Pharmaceutical Research	ISSN 2277 – 7105	https://wjpr.net/public/abstract_show/18121
5	Genotoxicity: Malignancy, Damage To DNA	Mr. S. A. Waghmare	Pharmaceutical Chemistry	World Journal of Pharmaceutical Research	ISSN 2277 – 7105	https://wjpr.net/abstract_show/18131
6	Herbal Medicines and Screening Models for Hepatoprotective Agents	Dr. S. A. Waghmare	Pharmacology	World Journal of Pharmaceutical Research	ISSN 2277 – 7105	https://www.google.com/search?q=Herbal+Medicines+and+Screening+Models+for+Hepatoprotective+Agents+research+paper+by+pratiksha+kothari&rlz=1C1VDKB_en-GBIN1070IN1070&oq=Herbal+Medicines+and+Screening+Models+for+Hepatoprotective+Agents+research+paper+by+pratiksha+kothari&gs_lcrp=EgZjaHJvbWUyBggAEEUYOTIGCAEQRRhAMgYIAhBFGDvSAQkyMDkxNWow

						ajeoAgCwAgA&sourceid=chrome&ie=UTF-8
7	Elucidation of Molecular Mechanism Involved in Nephroprotective Potential of Naringin in Ethylene Glycol-Induced Urolithiasis in Experimental Uninephrectomized Hypertensive Rats	Dr. A. D. Kandhare	Pharmacology	Latin American Journal of Pharmacy - 39 (5): 991-9 (2020)	ISSN 2362-3853	https://www.google.com/search?q=Elucidation+of+Molecular+Mechanism+Involved+in+Nephroprotective+Potential+of+Naringin+in+Ethylene+Glycol-Induced+Urolithiasis+in+Experimental+Uninephrectomized+Hypertensive+Rats+a.d.kandhare&rlz=1C1VDKB_en-GBIN1070IN1070&oq=Elucidation+of+Molecular+Mechanism+Involved+in+Nephroprotective+Potential+of+Naringin+in+Ethylene+Glycol-Induced+Urolithiasis+in+Experimental+Uninephrectomized+Hypertensive+Rats+a.d.kandhare&gs_lcrp=EgZjaHJvbWUyBggAEEUYOdIBCTQ1ODIxajBqNKgCALACAA&sourceid=chrome&ie=UTF-8
8	Herbal Medicines and Screening Models for Hepatoprotective Agents	Ms. P. P. Kothari	Pharmacology	World Journal of Pharmaceutical Research	ISSN 2277 – 7105	https://www.google.com/search?q=Herbal+Medicines+and+Screening+Models+for+Hepatoprotective+Agents+research+paper+by+pratiksha+kothari&rlz=1C1VDKB_en-GBIN1070IN1070&oq=Herbal+Medicines+and+Screening+Models+for+Hepatoprotective+Agents+research+paper+by+pratiksha+kothari&gs_lcrp=EgZjaHJvbWUyBggAEEUYOTIGCAEQRRhAMgYIAhBFGDvSAQkyMDkxNWowajeoAgCwAgA&sourceid=chrome&ie=UTF-8

9	Development and Validation of RP-HPLC and UV-Spectrophotometric Method for Simultaneous Estimation of Teneligliptin Hydrobromide Hydrate and Metformin Hydrochloride in Pharmaceutical dosage form	Mr. A. S. Gaikwad	Pharmacology	International Journal of Creative Research Thoughts	ISSN: 2320-2882	https://www.researchgate.net/publication/374813445_Development_and_Validation_of_RP-HPLC_and_UV-Spectrophotometric_Method_for_Simultaneous_Estimation_of_Teneligliptin_Hydrobromide_Hydrate_and_Metformin_Hydrochloride_in_Pharmaceutical_dosage_form
10	High Performance Liquid Chromatography -A Review	Mr. A. S. Gaikwad	Pharmacology	World Journal Of Pharmacy And Pharmaceutical Sciences	ISSN 2278 – 4357	
11	Development and Validation of RP-HPLC and UV Spectrophotometric Absorptivity Method for Simultaneous Estimation of Cyclobenzaprine hydrochloride and Aceclofenac in Pharmaceutical dosage form	Mr. A. S. Gaikwad	Pharmacology	International Journal of Scientific Development and Research	ISSN: 2455-2631	https://www.researchgate.net/publication/374813338_Development_and_Validation_of_RP-HPLC_and_UV-Spectrophotometric_Absorptivity_Method_for_Simultaneous_Estimation_of_Cyclobenzaprine_hydrochloride_and_Aceclofenac_in_Pharmaceutical_dosage_form
12	Synthesis, Characterization and Antidiabetic Evaluation of Sulfonamide in Corporated with 1,3,4-Oxadiazole Derivatives	Ms. M. T. Salve	Pharmacology	Indian Journal of Pharmaceutical Education and Research	Vol 55 Issue 4 1145	https://www.google.com/search?q=Synthesis%2C+Characterization+and+Antidiabetic+Evaluation+of+Sulfonamide+in+Corporated+with%0D%0A1%2C3%2C4-Oxadiazole+Derivatives%0D%0A+research+paper+by+m.t.salve

						&sca_esv=9c9edb63dbab71a4&sca_upv=1&rlz=1C1VDKB_en-GBIN1070IN1070&sxsrf=ACQVn0_YILL1hUj1r65PNTu3cPUJlg_93A%3A1713677958243&ei=hqYkZr-zDq3XseMP_pmd0A4&ved=0ahUKEwi_jtfvy9KFAxWta2wGHf5MB-oQ4dUDCBA&uact=5&oq=Synthesis%2C+Characterization+and+Antidiabetic+Evaluation+of+Sulfonamide+in+Corporated+with%0D%0A1%2C3%2C4-Oxadiazole+Derivatives%0D%0A+research+paper+by+m.t.salve&gs_lp=Egxnd3Mtd2l6LXNlcnAikwFTeW50aGVzaXMsIENoYXJhY3Rlcml6YXRpb24gYW5kIEFudGlkaWFiZXRpYyBFdmFsdWF0aW9uIG9mIFN1bGZvbmFtaWRlIGluIENvcnBvcmF0ZWQgd2l0aCAoxLDMsNC1PeGFkaWF6b2xlIERlcml2YXRpdmVzCiByZXNlYXJjaCBwYXBlciBieSBtLnQuQuc2FsdmVIAFAAWABwAHgAkAEAmAEAoAEAqgEAuAEDyAEA-AEC-AEBmAIAoAIAmAMAkgcAoAcA&sclient=gws-wiz-serp
13	Synthesis, Characterization, and Evaluation of in vitro Antidiabetic Activity of Novel Pyrrolidine Sulphonamide Derivative	Ms. M. T. Salve	Pharmacology	International Journal of Pharmaceutical Investigation	Vol 11, Issue 4	https://www.google.com/search?q=Synthesis%2C+Characterization%2C+and+Evaluation+of+in+vitro%0D%0AAntidiabetic+Activity+of+Novel+Pyrrolidine+Sulphonamide%0D%0ADerivative%0D%0A+research+paper+by+m.t.salve&sca_esv=9c9edb63dbab71a4&sca_upv=1&rlz=1C1VDKB_en-GBIN1070IN1070&sxsrf=ACQVn09IiPSXqpa0597xagN66VXIm4RFtw%3A1713678080795&ei=AKckZsqRMLaNseMPkbGW6AQ&ved=0ahUKEwjKkY-qzNKFAxW2RmwGHZGYBU0Q4

						dUDCBA&uact=5&oq=Synthesis%2C+Characterization%2C+and+Evaluation+of+in+vitro%0D%0AAntidiabetic+Activity+of+Novel+Pyrrolidine+Sulphonamide%0D%0ADerivative%0D%0A+research+paper+by+m.t.salve&gs_lp=Egxnd3Mtd2l6LXNlcnAilwFTeW50aGVzaXMsIENoYXJhY3Rlcml6YXRpb24sIGFuZCBFdmFsdWF0aW9uIG9mIGluIHZpdHJvCkFudGlkaWFiZXRpYyBBY3Rpdml0eSBvZiBOb3ZlbCBQeXJyb2xpZGluZSBTdWxwaG9uYW1pZGUKRGVyaXZhdGl2ZQogcmVzZWFyY2ggcGFwZXIgYnkgbS50LnNhbHZlMggQABiABBiiBEjWGFAAWMgQcAB4AJABAJgBpwGgAacBqgEDMC4xuAEDyAEA-AEBwAECmAIBoAKuAZgDAJIHAzAuMaAH2wI&sclient=gws-wiz-serp
14	Quality By Design Approaches To Analytical Method Development: A Review	Ms. S. S. Shingne	Pharmaceutical Chemistry	World Journal of Pharmaceutical Research World Journal of Pharmaceutical Research	ISSN 2277 – 7105	https://wjpr.net/public/abstract show/18121
15	Genotoxicity: Malignancy, Damage To DNA	Ms. S. S. Shingne	Pharmaceutical Chemistry	World Journal of Pharmaceutical Research	ISSN 2277 – 7105	https://wjpr.net/abstract_show/18131
16	Genotoxicity: Malignancy, Damage To DNA	Mr. A. E Bhujbal	Pharmaceutical Chemistry	World Journal of Pharmaceutical Research	ISSN 2277 – 7105	https://wjpr.net/abstract_show/18131

17	Quality By Design Approaches To Analytical Method Development: A Review	Mr. A. E Bhujbal	Pharmaceutical Chemistry	World Journal of Pharmaceutical Research World Journal of Pharmaceutical Research	ISSN 2277 – 7105	https://wjpr.net/public/abstract_show/18121
18	Isolation and characterization of phytoconstituents from petroleum ether extract of Momordica cochinchinensis fruits	Dr. M. R. Agrawal	Pharmacognosy	Journal of Pharmacognosy and Phytochemistry	E-ISSN: 2278-4136 P-ISSN: 2349-8234	https://scholar.google.co.in/scholar?q=Isolation+and+characterization+of+phytoconstituents+from+petroleum+ether+extract+of+Momordica+cochinchinensis+fruits+research+paper+by+m.+r.+agrawal&hl=en&as_sdt=0&as_vis=1&oi=scholart
19	Analgesic activity of Momordica cochinchinensis and Momordica balsamina fruit extracts	Dr. M. R. Agrawal	Pharmacognosy	International Journal of Green and Herbal Chemistry	Vol.7	https://scholar.google.co.in/scholar?q=Analgesic+activity+of+Momordica%0D%0Acochinchinensis+and+Momordica+balsamina+fruit+extracts%0D%0A+research+paper+by+m.+r.+agrawal&hl=en&as_sdt=0&as_vis=1&oi=scholart
20	Isolation, Characterization of Phytoconstituents from petroleum ether extract of Momordica balsamina fruits and its quantitative analysis by HPTLC	Dr. M. R. Agrawal	Pharmacognosy	Research Journal of Pharmacognosy and Phytochemistry	ISSN 0975-4385	https://rjpponline.org/AbstractView.aspx?PID=2020-12-3-6

Back to Paper Publication

Academic Year 2021-2022

Sr. No	Title of the paper	Name of Staff	Department	Name of the journal	ISSN number	Link
1	Gastro Retentive Drug Delivery System	Dr. H. V. Kamble	Pharmacology	World Journal of Pharmaceutical Research	ISSN 2277 – 7105	https://wjpr.net/abstract_show/18498

Back to Paper Publication

2	A Review: Drug-Excipient Interactions Study	Dr. H. V. Kamble	Pharmacology	World Journal of Pharmaceutical Research	ISSN 2277 – 7105	https://ijnrd.org/viewpaperforall.php?paper=IJNRD2302162
3	A Review On: Preservatives Used in Pharmaceuticals and Impacts on Health	**Dr. H. V. Kamble**	Pharmacology	Iconic Research And Engineering Journals	| ISSN: 2456-8880	https://scholar.google.co.in/scholar?q=A+Review+On:+Preservatives+Used+in+Pharmaceuticals+and+Impacts+on+Health+by+dr.+santosh+a.+waghmare&hl=en&as_sdt=0&as_vis=1&oi=scholart
4	A Review on Method Development, Validation, Optimization and Applications of HPLC	Dr. H. V. Kamble	Pharmacology	International Journal for Research in Applied Science & Engineering Technology	ISSN: 2321-9653	https://www.ijraset.com/research-paper/method-development-validation-optimization-and-applications-of-hplc
5	Excipients Compatibility Study Of Antilipidemic And Antihypertensive Drugs In Oral Solid Dosage Form	Dr. V. M. Satpute	Pharmaceutics	Neuroquantology	ISSN1303-5150	https://www.google.com/search?q=Excipients+Compatibility+Study+Of+Antilipidemic+And+Antihypertensive+Drugs+In+Oral+Solid+Dosage+Form+research+paper+by+dr.+v.+m.+satpute&rlz=1C1VDKB_en-GBIN1070IN1070&oq=Excipients+Compatibility+Study+Of+Antilipidemic+And+Antihypertensive+Drugs+In+Oral+Solid+Dosage+Form+research+paper+by+dr.+v.+m.+satpute&gs_lcrp=EgZjaHJvbWUyBggAEEUYOTIGCAEQIxgnMgYIAhAjGCfSAQkyMTQwN2owajeoAgiwAgE&sourceid=chrome&ie=UTF-8

6	Formulation And Evaluation Of Selexipag And Ezetimibe Bi-Layer Tablet By Novel Approach	Dr. V. M. Satpute	Pharmaceutics	Neuroquantology	ISSN1303-5150	https://www.neliti.com/publications/429774/formulation-and-evaluation-of-selexipag-and-ezetimibe-bi-layer-tablet-by-novel-a
7	A Review: Drug-Excipient Interactions Study	Dr. V. M. Satpute	Pharmaceutical Chemistry	World Journal of Pharmaceutical Research	ISSN 2277 – 7105	https://ijnrd.org/viewpaperforall.php?paper=IJNRD2302162
8	Gastro Retentive Drug Delivery System	Dr. S. A. Waghamare	Pharmaceutical Chemistry	World Journal of Pharmaceutical Research	ISSN 2277 – 7105	https://wjpr.net/abstract_show/18498
9	A Review: Drug-Excipient Interactions Study	Dr. S. A. Waghamare	Pharmaceutical Chemistry	World Journal of Pharmaceutical Research	ISSN 2277 – 7105	https://ijnrd.org/viewpaperforall.php?paper=IJNRD2302162
10	A Review on Method Development, Validation, Optimization and Applications of HPLC	Dr. S. A. Waghamare	Pharmaceutical Chemistry	International Journal for Research in Applied Science & Engineering Technology	ISSN: 2321-9653	https://www.ijraset.com/research-paper/method-development-validation-optimization-and-applications-of-hplc
11	A Review On: Preservatives Used in Pharmaceuticals and Impacts on Health	Dr. S. A. Waghamare	Pharmaceutical Chemistry	Iconic Research And Engineering Journals	| ISSN: 2456-8880	https://scholar.google.co.in/scholar?q=A+Review+On:+Preservatives+Used+in+Pharmaceuticals+and+Impacts+on+Health+by+dr.+santosh+a.+waghmare&hl=en&as_sdt=0&as_vis=1&oi=scholart
12	Evaluation of Some Natural Polymers as	Mr. A. V. Dhobale	Pharmaceutics	International Journal of	www.ijrar.org	https://www.researchgate.net/publication/350782955_Evaluation_of_Some_Natural_Polymers_as_

	Pharmaceutical Excipient For Sustained Drug Delivery System			Research and Analytical Reviews (IJRAR)	E-ISSN 2348-1269	Pharmaceutical Excipient For Sustained Drug Delivery System
13	Green Chemistry	Mr. P. B. Cholke	Pharmaceutics	International Journal of Creative Research T	ISSN: 2320-2882	https://www.google.com/search?q=Green+Chemistry+research+paper+by+pravin+cholake&sca_esdbab71a4&sca_upv=DKB_en-GBIN1070IN1070&biw=1366&bih=641&sxsrf=ACQVn09_J2GQj8sO9QI6WxcNZ8to4sgBjw%3A1713679595795&ei=66wkZvuJMIzT1e8Pnaug2A4&ved=0ahUKEwi7q8P80dKFAxWMafUHHZ0VCOsQ4dUDCBA&uact=5&oq=Green+Chemistry+research+paper+by+pravin+cholake&gs_lp=Egxnd3Mtd2l6LXNlcnAiMEdyZWVuIENoZW1pc3RyeSByZXNlYXJjaCBwYXBlciBieSBwcmF2aW4gY2hvbGFrZTIHECEYoAEYCjIHECEYoAEYCkjBYFAAWJVccAB4AJABAZgB7AOgAd40qgEKMC42LjE3LjQuMbgBA8gBAPgBAvgBAZgCGKAC2C7CAggQABiABBiiBMICBRAhGKABwgIEECEYCsICBRAhGJ8FwgIGEAAYFhgewgILEAAYgAQYhgMYigXCAgQQIxgnmAMAkgcKMC41LjE0LjQuMaAH5YQB&sclient=gws-wiz-serp
14	Green Chemistry	Y. S. Temak	Pharmaceutics	International Journal of Creative Research Thoughts	ISSN: 2320-2882	https://www.google.com/search?q=Green+Chemistry+research+paper+by+pravin+cholake&sca_esv=9c9edb63dbab71a4&sca_upv=1&rlz=1C1VDKB_en-GBIN1070IN1070&biw=1366&bih=641&sxsrf=ACQVn09_J2GQj8sO9QI6WxcNZ8to4sgBjw%3A1713679595795&ei=66wkZvuJMIzT1e8Pnaug2A4&ved=0ahUKEwi7q

Back to Paper Publication

						8P80dKFAxWMafUHHZ0VCOsQ4dUDCBA&uact=5&oq=Green+Chemistry+research+paper+by+pravin+cholake&gs_lp=Egxnd3Mtd2l6LXNlcnAiMEdyZWVuIENoZW1pc3RyeSByZXNlYXJjaCBwYXBlciBieSBwcmF2aW4gY2hvbGFrZTIHECEYoAEYCjIHECEYoAEYCkjBYFAAWJVccAB4AJABAZgB7AOgAd40qgEKMC42LjE3LjQuMbgBA8gBAPgBAvgBAZgCGKAC2C7CAggQABiABBiiBMICBRAhGKABwgIEECEYCsICBRAhGJ8FwgIGEAAYFhgewgILEAAYgAQYhgMYigXCAgQQIxgnmAMAkgcKMC41LjE0LjQuMaAH5YQB&sclient=gws-wiz-serp
15	An Overview on Artificial Blood	Ms. S. B. Kshirsagar	Pharmaceutical Chemistry	International Journal In Pharmaceutical Sciences	Vol 1, Issue 1, 42-48	https://scholar.google.co.in/scholar?q=An+Overview+on+Artificial+Blood+by+s.+b.+kshirsagar&hl=en&as_sdt=0&as_vis=1&oi=scholart

Back to Paper Publication

Academic Year 2022-2023

Sr. no	Title of the paper	Name of Staff	Department	Name of the journal	ISSN number	Link
1	A Review: Etiology, Pathogenesis And Treatment Of Peptic Ulcer	Dr. H. V. Kamble	Pharmacology	World Journal of Pharmaceutical Research	ISSN 2277 – 7105	https://scholar.google.co.in/scholar?q=A+Review:+Etiology,+Pathogenesis+And+Treatment+Of+Peptic+Ulcer+by+H.V.+Kamble&hl=en&as_sdt=0&as_vis=1&oi=scholart
2	Formulation and Evaluation of Sustained Release Ibuprofen Tablet	Dr. H. V. Kamble	Pharmacology	International Journal of Scientific Research and Engineering Development	Volume 6 Issue 2	https://scholar.google.co.in/scholar?q=Formulation+and+Evaluation+of+Sustained+Release+Ibuprofen+Tablet++by+H.V.+Kamble&hl=en&as_sdt=0&as_vis=1&oi=scholart
3	A Comprehensive Review On Clinical Trials	Dr. H. V. Kamble	Pharmacology	World Journal of Pharmaceutical Research World Journal of Pharmaceutical Research	ISSN 2277 – 7105	https://scholar.google.co.in/scholar?q=A+Comprehensive+Review+On+Clinical+Trials+by+H.V.+Kamble&hl=en&as_sdt=0&as_vis=1&oi=scholart
4	Face Care Cosmetic-A Review On Herbal Face Toner	Dr. H. V. Kamble	Pharmacology	World J Pharm Pharm Sci	ISSN 2277 – 7105	https://www.wjpr.net/abstract_show/21922

5	Novel Drug Delivery System : Review Article	Dr. H. V. Kamble	Pharmacology	WJPR	ISSN: 2277-7105	https://www.wjpr.net/abstract_show/22172
6	Face Care Cosmetic-A Review On Herbal Face Toner	Dr. V. M. Satpute	Pharmaceutics	World J Pharm Pharm Sci	ISSN 2277 – 7105	https://www.wjpr.net/abstract_show/21922
7	Chloroquine and Hydroxychloroquine in Coronavirus Disease-19: The Real Savior or a False-positive Treatment	Dr. V. M. Satpute	Pharmaceutics	JETIR	ISSN 2349-5162	https://www.google.com/search?q=Chloroquine+and+Hydroxychloroquine+in+Coronavirus+Disease-19%3A+The+Real+Savior+or+a+False-positive+Treatment+research+paper+by+dr.v.m.satpute%3AJETIR&sca_esv=d143a126214438bd&sca_upv=1&rlz=1C1VDKB_en-GBIN1070IN1070&sxsrf=ACQVn08BcAMGn4TbxWJyDzkQHRBmakjxuw%3A1713681337038&ei=ubMkZsz7AY6Yvr0P5MGs0A8&ved=0ahUKEwiMvOi62NKFAxUOjK8BHeQgC_oQ4dUDCBA&uact=5&oq=Chloroquine+and+Hydroxychloroquine+in+Coronavirus+Disease-19%3A+The+Real+Savior+or+a+False-positive+Treatment+research+paper+by+dr.v.m.satpute%3AJETIR&gs_lp=Egxnd3Mtd2l6LXNlcnAikgFDaGxvcm9xdWluZSBhbmQgSHlkcm94eWNobG9yb3F1aW5lIGluIENvcm9uYXZpcnVzIERpc2Vhc2UtMTk6IFRoZSBSZWFsIFNhdmlvciBvciBhIEZhbHNlLXBvc2l0aXZlIFRyZWF0bWVudCByZXNlYXJjaCBwYXBlciBieSBkci52Lm0uc2F0cHV0ZTpKRVRJUkgAUABYAHAAeACQAQCYAQCgAQCqAQC4AQPIAQD4AQGYAgCgAgCYAwCSBwCgBwA&sclient=gws-wiz-serp

8	Formulation and Evaluation of Sustained Release Ibuprofen Tablet	Dr. S. A. Waghmare	Pharmaceutical Chemistry	International Journal of Scientific Research and Engineering Development	Volume 6 Issue 2	https://scholar.google.co.in/scholar?q=Formulation+and+Evaluation+of+Sustained+Release+Ibuprofen+Tablet++by+H.V.+Kamble&hl=en&as_sdt=0&as_vis=1&oi=scholart
9	A Comprehensive Review On Clinical Trials	Dr. S. A. Waghmare	Pharmaceutical Chemistry	World Journal of Pharmaceutical Research World Journal of Pharmaceutical Research	ISSN 2277 – 7105	https://scholar.google.co.in/scholar?q=A+Comprehensive+Review+On+Clinical+Trials+by+H.V.+Kamble&hl=en&as_sdt=0&as_vis=1&oi=scholart
10	Solid Lipid Nanoparticle (Sln)	Dr. S. A. Waghmare	Pharmaceutical Chemistry	International Research Journal of Modernization in Engineering Technology and Science	ISSN: 2582-5208	https://www.researchgate.net/publication/283600259_Solid_lipid_nanoparticles_A_promising_drug_delivery_system
11	Face Care Cosmetic-A Review On Herbal Face Toner	**Mr. S. A. Waghmare**	Pharmaceutical Chemistry	World J Pharm Pharm Sci	ISSN 2277 – 7105	https://www.wjpr.net/abstract_show/21922
12	Novel Drug Delivery System : Review Article	**Mr. S. A. Waghmare**	Pharmaceutical Chemistry	WJPR	ISSN: 2277-7105	https://www.wjpr.net/abstract_show/22172
13	Novel Maos of Novel Cinnoline Derivatives	Ms. Bhalerao Monali	Pharmaceutical Chemistry	Eur.Chem. Bull	ISSN 2063-5346	https://www.google.com/search?q=Development+and+validation+of+chromatographic+method+for+analysis+of+some+polycyclic

		Ramchandra				+hydrocarbons+on+soybean+oil+by+ms.+monali+r.+bhalerao&rlz=1C1VDKB_en-GBIN1070IN1070&oq=Development+and+validation+of+chromatographic+method+for+analysis+of+some+polycyclic+hydrocarbons+on+soybean+oil+by+ms.+monali+r.+bhalerao&gs_lcrp=EgZjaHJvbWUyBggAEEUYOTIHCAEQIRiPAjIHCAIQIRiPAtIBCTIwMDUzajBqN6gCCLACAQ&sourceid=chrome&ie=UTF-8
14	Development and validation of chromatographic method for analysis of some polycyclic hydrocarbons on soybean oil	Ms. Bhalerao Monali Ramchandra	Pharmaceutical Chemistry	Eur.Chem. Bull	ISSN 2063-5346	https://www.google.com/search?q=Development+and+validation+of+chromatographic+method+for+analysis+of+some+polycyclic+hydrocarbons+on+soybean+oil+by+ms.+monali+r.+bhalerao&rlz=1C1VDKB_en-GBIN1070IN1070&oq=Development+and+validation+of+chromatographic+method+for+analysis+of+some+polycyclic+hydrocarbons+on+soybean+oil+by+ms.+monali+r.+bhalerao&gs_lcrp=EgZjaHJvbWUyBggAEEUYOTIHCAEQIRiPAjIHCAIQIRiPAtIBCTIwMDUzajBqN6gCCLACAQ&sourceid=chrome&ie=UTF-8
15	A Review on Beta Vulgaris (Beet Root)	Mr. A. S. Gaikwad	Pharmaceutical Chemistry	PENSEE	ISSN NO: 0031-4773	https://www.researchgate.net/publication/374813414_A_Review_on_Beta_Vulgaris_Beet_Roo
16	Microsponge Drug Delivery System - A Review	Mr. A. S. Gaikwad	Pharmaceutical Chemistry	PENSEE	ISSN NO: 0031-4773	https://www.researchgate.net/publication/374813521_Microsponge_drug_delivery_Scopus

17	Development And Validation Of RP HPLC Method For Simultaneous Estimation Of Zaltoprofen And Paracetamol In Bulk And Tablet Formulation	Mr. A. S. Gaikwad	Pharmaceutical Chemistry	International journal Of Current Research	ISSN: 0975-833	https://www.researchgate.net/publication/374813419_DEVELOPMENT_AND_VALIDATION_OF_RP_ZALTOPROFEN_AND_PARACETAMOL
18		Ms. M. T. Salve	Pharmaceutical Chemistry	Indian Journal of Pharmaceutical Education and Research	Vol 57, Issue 3(Suppl), Ju	https://www.ijper.org/article/2084
19	Novel Drug Delivery System : Review Article	Ms. S. S. Khillare	Pharmaceutics	WJPR	ISSN: 2277-7105	https://www.wjpr.net/abstract_show/22172
20	A Review on Nanotechnology in Cancer Treatment	Mr. M. M.Garje	Pharmaceutical Chemistry	IJSRT	Vol 7: Isue 9	https://www.google.com/search?q=A+Review+on+Nanotechnology+in+Cancer+Treatment+by+mr.m.m.+garje%3AIJCRT&sca_esv=d143a126214438bd&sca_upv=1&rlz=1C1VDKB_en-GBIN1070IN1070&biw=1366&bih=641&sxsrf=ACQVn08bbuYmF_F1tdObiUbhEUGOJQqEzg%3A1713682995582&ei=M7okZsmQI73fseMPxqWQeA&ved=0ahUKEwiJ9tXR3tKFAxW9b2wGHcYSBA8Q4dUDCBA&uact=5&oq=A+Review+on+Nanotechnology+in+Cancer+Treatment+by+mr.m.m.+garje%3AIJCRT&gs_lp=Egxnd3Mtd2l6LXNlcnAiRUEgUmV2aWV3IG9uIE5hbm90ZWNobm9sb2d5IGluIENhbmNlciBUcmVhdG1lbnQgYnkgbXIubS5tLiBnYXJqZTpJSkNSVEgAUABYAHAAeACQAQCYAQCgAQCqAQC4AQPIAQD4AQGYAgCgAgCYA

						wCSBwCgBwA&sclient=gws-wiz-serp

Back to Paper Publication

From the beginning, the goal of Loknete Shri Dadapatil Pharate College of Pharmacy (LSDPCOP), Mandavgan Pharata, has been to give our rural students access to cutting edge educational resources so they may successfully compete with the outside world. The main goal is to "Nurturing Pharmacy Professionals through Quality Education" so that they can effectively serve the profession. In the end, we are the company that carries out our social obligations in addition to offering top-notch instruction to create competent, professional chemists. Moral education is vital, in our opinion, because it is at this age that children take their first steps towards adulthood, and it is crucial that they do so properly.

Their views, convictions, and concepts are shaped by these moral principles, which also aid in their development as morally robust and unwavering individuals. The students at Loknete Shri Dadapatil rmacy (LSDPCOP), Mandavgan Pharata, are constantly being instilled with s as well as social duties and responsibilities. This has been accomplished by creating and delivering Life Skills courses to our pupils. Life skills are those proficiencies that enable a person to be adaptive and optimistic in the face of life's challenges. The goal is the development of one's personality through self-awareness, interpersonal relationships, introspection on both the concrete and abstract, change-leading and change-generating, and adherence to tried-and-true beliefs. By exposing students to the ideas that underpin both personal and professional success and assisting them in acquiring the skills necessary to put these ideas into practice in their daily lives and jobs, this course aims to maximise their potential and increase their employability. In order to attract superior talent, STCOP is placing greater emphasis on teaching life skills, including professional, leadership, and management abilities, as well as communication skills. We believe that mastering life skills is the key to success.

Back to Paper Publication

The goal of the life skills course is to instill in the students a sense of self-awareness and empathy as well as professional abilities in the areas of interpersonal skills, high-performing teams, leadership potential, communication and presentation skills, problem-solving techniques, decision-making, inspiring creativity and innovation for both professional and personal excellence, stress management, time management, conflict resolution, and human values instillation. As facilitators, mentors, and trainers, the faculty members and students must actively participate in the course. Students are motivated and enrolled in the courses they choose for this reason.

Oral presentations, spontaneous speaking, quizzes, debates, case studies, creative thinking, team-building exercises, field trips, projects, psychometric analysis, and other activities have all been included into the courses. Upon successfully finishing the course, they receive a course completion certificate.